



## MAXWELL UNDERGROUND MINE PROJECT

### **APPENDIX A**

#### **Biodiversity Development Assessment Report**





**MAXWELL UNDERGROUND MINE PROJECT  
MINE ENTRY AREA MODIFICATION**



**BIODIVERSITY DEVELOPMENT ASSESSMENT REPORT**



Prepared by  
Dr Colin Driscoll  
BAAS17004

August 2021



5 August 2021

**Maxwell Underground Mine Project  
Mine Entry Area Modification  
Biodiversity Development Assessment Report  
Statement of Currency**

As required by Section 6.15 (1) of the NSW *Biodiversity Conservation Act 2016* this letter certifies that the Maxwell Underground Mine Project Mine Entry Area Modification Biodiversity Assessment Report has been prepared on the basis of the requirements of (and information provided under) the biodiversity assessment method as at 5 August 2021.

Yours Faithfully  
HUNTER ECO



Dr Colin Driscoll  
BAAS17004

HUNTER ECO · ABN 25 112 984 240  
PO Box 1047, Toronto, NSW 2283 M 0438 773 029 E cd\_enviro@bigpond.com

## TABLE OF CONTENTS

<b>DEFINITIONS .....</b>	<b>7</b>
<b>EXECUTIVE SUMMARY .....</b>	<b>10</b>
<b>1 INTRODUCTION.....</b>	<b>12</b>
1.1 BACKGROUND.....	12
1.2 MODIFICATION DESCRIPTION .....	14
1.3 GENERAL DESCRIPTION OF THE SUBJECT LAND .....	14
1.4 STRUCTURE OF THIS ASSESSMENT .....	14
1.5 INFORMATION SOURCES USED IN THIS ASSESSMENT.....	17
1.5.1 Field Survey Reports.....	17
1.5.2 Published Databases.....	17
1.5.3 Local Data .....	17
1.5.4 BAM-C.....	17
<b>2 LANDSCAPE FEATURES .....</b>	<b>18</b>
2.1 TOPOGRAPHY, HYDROLOGICAL SETTING, GEOLOGY AND SOILS.....	18
2.2 NATIVE VEGETATION COVER .....	18
2.3 IBRA BIOREGIONS AND SUB-REGIONS.....	19
2.4 RIVERS AND STREAMS .....	19
2.5 WETLANDS.....	19
2.6 HABITAT CONNECTIVITY .....	19
2.7 KARST, CAVES, CREVICES, CLIFFS, ROCKS AND OTHER GEOLOGICAL FEATURES .....	19
2.8 AREAS OF OUTSTANDING BIODIVERSITY VALUE .....	22
2.9 NSW MITCHELL LANDSCAPE .....	22
<b>3 NATIVE VEGETATION .....</b>	<b>23</b>
3.1 EXISTING INFORMATION ON NATIVE VEGETATION.....	23
3.2 SUPPLEMENTRY SURVEYS.....	23
3.3 NATIVE VEGETATION EXTENT .....	23
3.4 AREAS WITHOUT NATIVE VEGETATION .....	26
3.5 PLANT COMMUNITY TYPES.....	26
3.5.1 Plant Community Types and Classes .....	26
3.5.2 Evidence to Support Plant Community Types .....	26
3.5.3 Plant Community Types Percent Cleared Value .....	27
3.5.4 Threatened Ecological Communities .....	29
3.5.5 Evidence to Support Threatened Ecological Communities .....	29
3.6 VEGETATION INTEGRITY ASSESSMENT .....	29
3.6.1 Vegetation Zones.....	29
3.6.2 Patch Size.....	30
3.6.3 Vegetation Integrity Plots.....	30
3.6.4 Vegetation Integrity Score.....	30
<b>4 THREATENED SPECIES .....</b>	<b>32</b>
4.1 ECOSYSTEM CREDIT SPECIES - HABITAT SUITABILITY ASSESSMENT .....	32
4.1.1 Ecosystem Species from the BAM-C.....	32
4.1.2 Exclusion of Ecosystem Species.....	34
4.1.3 Addition of Ecosystem Species.....	36
4.2 SPECIES CREDIT SPECIES - HABITAT SUITABILITY ASSESSMENT.....	36

4.2.1	Species Credit Species from the BAM-C .....	37
4.2.2	Exclusion of Species Credit Species .....	38
4.2.3	Addition of Species Credit Species .....	44
4.3	SPECIES CREDIT SPECIES – ASSESSMENT METHOD.....	45
4.3.1	Species Assumed to be Present.....	45
4.3.2	Species Important Habitat Mapping .....	45
4.3.3	Species for Survey .....	45
4.3.4	Export Reports.....	45
4.4	SPECIES CREDIT SPECIES – SURVEY .....	45
4.4.1	Survey Results .....	54
4.4.2	Species Polygons .....	59
<b>5</b>	<b>PRESCRIBED BIODIVERSITY IMPACT ENTITIES .....</b>	<b>60</b>
5.1	KARST, CAVES, CREVICES, CLIFFS, ROCKS AND OTHER GEOLOGICAL FEATURES .....	60
5.2	HUMAN MADE STRUCTURES AND NON-NATIVE VEGETATION .....	60
5.3	HABITAT CORRIDORS OR LINKAGES.....	60
5.4	WATERBODIES OR HYDROLOGICAL PROCESSES THAT SUSTAIN THREATENED SPECIES .....	60
5.5	THREATENED SPECIES AT RISK FROM VEHICLE STRIKE .....	60
<b>6</b>	<b>AVOID AND MINIMISE IMPACTS .....</b>	<b>61</b>
<b>7</b>	<b>EVALUATION OF POTENTIAL IMPACTS .....</b>	<b>62</b>
7.1	DIRECT IMPACTS ON NATIVE VEGETATION AND HABITAT .....	62
7.1.1	Clearance of Native Vegetation.....	62
7.1.2	Cumulative Impacts .....	62
7.2	INDIRECT IMPACTS ON NATIVE VEGETATION AND HABITAT .....	62
7.2.1	Inadvertent Impacts on Adjacent Habitat or Vegetation .....	64
7.2.2	Impacts on Adjacent Habitat or Vegetation from a Change in Land Use Pattern (Increased Human Activity) .....	64
7.2.3	Reduced Viability of Adjacent Habitat Due to Edge Effects .....	64
7.2.4	Reduced Viability of Adjacent Habitat Due to Noise, Dust or Light Spill .....	64
7.2.5	Transport of Weeds and Pathogens from the Site to Adjacent Vegetation .....	64
7.2.6	Increased Risk of Fauna Starvation, Exposure and Loss of Shade or Shelter .....	65
7.2.7	Loss of Breeding Habitats .....	65
7.2.8	Trampling of Threatened Flora Species .....	65
7.2.9	Inhibition of Nitrogen Fixation and Increased Soil Salinity.....	65
7.2.10	Fertiliser Drift .....	65
7.2.11	Rubbish Dumping .....	65
7.2.12	Wood Collection .....	65
7.2.13	Bush Rock Removal and Disturbance .....	65
7.2.14	Increase in Predatory Species Populations.....	66
7.2.15	Increase in Pest Animal Populations .....	66
7.2.16	Increased Risk of Fire.....	66
7.2.17	Disturbance to Specialist Breeding and Foraging Habitat .....	66
7.2.18	Groundwater Dependent Vegetation.....	66
7.3	PRESCRIBED BIODIVERSITY IMPACTS.....	66
7.3.1	Karst, Caves, Crevices, Cliffs, Rocks and Other Geological Features .....	67
7.3.2	Human Made Structures .....	67
7.3.3	Non-Native Vegetation.....	67
7.3.4	Habitat Connectivity.....	67

7.3.5	Fauna Movement .....	68
7.3.6	Water Quality, Water Bodies and Hydrological Processes that Sustain Threatened Species and Threatened Ecological Communities .....	68
7.3.7	Wind Turbines .....	68
7.3.8	Vehicle Strike .....	68
7.4	IMPACTS ON THREATENED SPECIES AND COMMUNITIES UNDER THE NSW FISHERIES MANAGEMENT ACT 1994 .....	68
7.5	STATE ENVIRONMENTAL PLANNING POLICY (KOALA HABITAT PROTECTION) .....	69
7.6	COMMONWEALTH ENVIRONMENT PROTECTION AND BIODIVERSITY CONSERVATION ACT 1999 .....	69
7.6.1	Background.....	69
7.6.2	Box-Gum TEC.....	70
7.6.3	Striped Legless Lizard .....	72
7.6.4	Other Species.....	74
7.6.5	Mitigation Measures Relevant to MNES .....	75
7.6.6	Conclusion .....	75
<b>8</b>	<b>MEASURES TO MITIGATE AND MANAGE IMPACTS .....</b>	<b>79</b>
<b>9</b>	<b>IMPACT SUMMARY.....</b>	<b>83</b>
9.1	SERIOUS AND IRREVERSIBLE IMPACTS .....	83
9.1.1	BAM Requirements .....	83
9.1.2	BC Regulation Requirements.....	87
9.1.3	Conclusion .....	88
9.2	IMPACTS REQUIRING/NOT REQUIRING AN OFFSET .....	88
9.3	AREAS NOT REQUIRING ASSESSMENT .....	88
9.4	BIODIVERSITY CREDITS.....	88
<b>10</b>	<b>CONCLUSION .....</b>	<b>93</b>
<b>11</b>	<b>REFERENCES.....</b>	<b>94</b>

**TABLES**

Table 1	Native Vegetation Extent in the Buffer Area
Table 2	Plant Community Type Data
Table 3	Threatened Ecological Communities
Table 4	Vegetation Integrity Score Detail
Table 5	Ecosystem Species from the BAM-C
Table 6	Ecosystem Species Excluded from the BAM-C
Table 7	Species Credit Species for Assessment
Table 8	Flora Species Credit Species – Geographic Constraints
Table 9	Species Credit Species - Habitat Constraints
Table 10	Species Credit Species Degraded Habitat
Table 11	Species Credit Species Requiring Survey and Timing
Table 12	Species Credit Species Presence
Table 13	Native Vegetation Clearance Summary
Table 14	Impact Mitigation Measures Relevant to Threatened Species and Communities listed under the EPBC Act

Table 15	Summary of Conservation Advice, Recovery Plans or Threat Abatement Plans for EPBC Listed Species in the Project Area
Table 16	Measures to Mitigate and Manage Impacts
Table 17	Modification Ecosystem Credit Requirements
Table 18	Modification Species Credit Requirements

## FIGURES

Figure 1	Regional Location
Figure 2	Approved General Arrangement
Figure 3	Proposed General Arrangement
Figure 4	The Topography of the Mine Entry Components in Relation to the Approved Development
Figure 5	Site Map
Figure 6	Location Map
Figure 7a	Vegetation Mapping – Mine Entry Area
Figure 7b	Vegetation Mapping – Maxwell Infrastructure
Figure 8	Threatened Flora Species/Endangered Populations
Figure 9	Threatened Reptiles and Amphibians
Figure 10	Threatened Birds
Figure 11	Threatened Mammals
Figure 12	Threatened Bats
Figure 13	Pink-tailed Legless Lizard Species Polygon
Figure 14	Striped Legless Lizard Species Polygon
Figure 15	Squirrel Glider Species Polygon
Figure 16	Southern Myotis Species Polygon
Figure 17	Rehabilitation Domains
Figure 18	Serious and Irreversible Impact Entities
Figure 19	Areas Requiring Offset
Figure 20	Areas not Requiring Assessment
Figure 21	Areas not Requiring Offset

## ATTACHMENTS

Attachment A	Maxwell Project Baseline Fauna Survey Report
Attachment B	Vegetation Integrity Data
Attachment C	Expert Report Expected Presence of Threatened Terrestrial Orchids ( <i>Diuris tricolor</i> , <i>Prasophyllum petilum</i> , <i>Pterostylis chaetophora</i> ) Maxwell Project
Attachment D	Threatened Flora and Fauna Species Known or Predicted to occur in the Locality
Attachment E	BAM Biodiversity Credit Report - Biodiversity Assessment Development Footprint
Attachment F	BAM Biodiversity Credit Report (Like for Like)
Attachment G	BAM Biodiversity Credit Report (Variations)

**PLATES**

Plate 1 Proposed MEA Extensions – PCT 1606 (in Derived Native Grassland Form)

Plate 2 Proposed MEA Extensions - PCT 1692

Plate 3 Proposed Water Treatment Facility Location - PCT 1692



## DEFINITIONS

**BAM:** *the Biodiversity Assessment Method.*

**BC Act:** *the NSW Biodiversity Conservation Act 2016.*

**BC Regulation:** *the NSW Biodiversity Conservation Regulation 2017.*

**Benchmarks:** *the quantitative measures that represent the 'best attainable' condition, which acknowledges that native vegetation within the contemporary landscape has been subject to both natural and human-induced disturbance. Benchmarks are defined for specified variables for each PCT. Vegetation with relatively little evidence of modification generally has minimal timber harvesting (few stumps, coppicing, cut logs), minimal firewood collection, minimal exotic weed cover, minimal grazing and trampling by introduced or overabundant native herbivores, minimal soil disturbance, minimal canopy dieback, no evidence of recent fire or flood, no high-frequency burning, and evidence of recruitment of native species.*

**Biodiversity Assessment Method Calculator:** *the online computer program that provides decision support to assessors and proponents by applying the BAM and referred to as the BAM-C. The BAM-C contains biodiversity data from the BioNet Vegetation Classification and the Threatened Biodiversity Data Collection that the assessor is required to use in a BAM assessment. The BAM-C applies the equations used in the BAM, including those to determine the number and class of biodiversity credits required to offset the impacts of a development, or created at a biodiversity stewardship site. It is published by the Department.*

**Biodiversity credit report:** *the report produced by the BAM-C that sets out the number and class of biodiversity credits required to offset the remaining adverse impacts on biodiversity values at a development site or on land to be biodiversity certified. For biodiversity stewardship sites, the biodiversity credit report sets out the number and class of biodiversity credits that are created at that site.*

**Biodiversity Development Assessment Report (BDAR):** *a report prepared by an accredited person in relation to proposed development or activity that would be authorised by a planning approval, or proposed clearing that would be authorised by a vegetation clearing approval, that:*

- (a) *assesses in accordance with the BAM the biodiversity values of the land subject to the proposed development, activity or clearing;*
- (b) *assesses in accordance with the BAM the impact of proposed development, activity or clearing on the biodiversity values of that land;*
- (c) *sets out the measures that the proponent of the proposed development, activity or clearing proposes to take to avoid or minimise the impact of the proposed development, activity or clearing; and*
- (d) *specifies in accordance with the BAM the number and class of biodiversity credits that are required to be retired to offset the residual impacts on biodiversity values of the actions to which the biodiversity offsets scheme applies.*

**Biodiversity Offsets:** *the gain in biodiversity values achieved from the implementation of management actions on areas of land, to compensate for losses to biodiversity values from the impacts of development.*

**Biodiversity Stewardship Agreement:** *means a biodiversity stewardship agreement made under Division 2 of Part 5 of the BC Act.*

**Biodiversity Stewardship Site:** *means the land that is designated by a biodiversity stewardship agreement to be a biodiversity stewardship site for the purposes of the BC Act.*

**Biodiversity Stewardship Site Assessment Report (BSSAR):** *the report that must be prepared in accordance with the BAM and submitted as part of an application for a biodiversity stewardship agreement.*

**Broad Condition State:** areas of the same Plant Community Type that are in relatively homogenous condition. Broad condition is used for stratifying areas of the same Plant Community Type into a vegetation zone for the purpose of determining the vegetation integrity score.

**Class of biodiversity credit:** biodiversity credits that share the same attributes (refer to Subsection 10.2 of the BAM 2020).

**Development Footprint:** the area of land that is directly impacted on by a proposed development, including access roads and areas used to store construction materials. The term development footprint is also taken to include clearing footprint, except where the reference is to a small area development or a major project development.

**Ecosystem credits:** a measurement of the value of threatened ecological communities, threatened species habitat for species that can be reliably predicted to occur with a PCT, and PCTs generally. Ecosystem credits measure the loss in biodiversity values at a development, activity, clearing or biodiversity certification site and the gain in biodiversity values at a biodiversity stewardship site.

**EPBC Act:** the Commonwealth Environment Protection and Biodiversity Conservation Act 1999.

**High threat weed cover:** plant cover composed of vascular plants that, if not controlled, will invade and outcompete native plant species. Also referred to as high threat weeds or high threat exotic vegetation. Plants considered to be high threat weeds are listed on the high threat weeds list published in the BAM-C

**Native Vegetation Cover:** the percentage of native vegetation cover on the subject land and the surrounding buffer area. Cover estimates are based on the cover of native woody and non-woody vegetation. Native vegetation cover includes regrowth, derived native grasslands and plantations that are comprised of plants native to New South Wales

**Plant Community Type (PCT):** a NSW plant community type identified using the Plant Community Type classification system.

**Retirement of Credits:** the action taken whereby biodiversity credits created for a biobanking agreement or a biodiversity stewardship agreement are used to offset the impacts of development, clearing or biodiversity certification.

**Sensitivity to Loss:** a component of the biodiversity risk weighting for an entity that considers the increased threat posed to an entity from offsetting the loss of habitat or population.

**Serious and irreversible impact:** impacts likely to contribute significantly to the risk of a threatened species or ecological community becoming extinct in accordance with the principles set out in clause 6.7(2) of the BC regulation

**Serious and Irreversible Impacts (SAII):** impacts likely to contribute significantly to the risk of extinction of a threatened species or ecological community in NSW.

**Site-based Development:** a development other than a linear-shaped development, or a multiple fragmentation impact development.

**Species Credit Species:** threatened species or components of species habitat that are identified in the Threatened Species Data Collection as requiring assessment for species credits.

**Species Credits:** the class of biodiversity credits created or required for the impact on threatened species that cannot be reliably predicted to use an area of land based on habitat surrogates. Species that require species credits are listed in the Threatened Biodiversity Data Collection.

**Subject Land:** is land subject to a development, activity, clearing, biodiversity certification or a biodiversity stewardship proposal. It excludes the assessment area which surrounds the subject land (i.e. the area of land in the 1500 m buffer zone around the subject land or 500 m buffer zone for linear proposals). In the case of a biodiversity certification proposal, subject land includes the biodiversity certification assessment area.

**Vegetation Class:** *a level of classification of vegetation communities, as defined in Keith (2004).*

**Vegetation Formation:** *a broad level of vegetation classification as defined in Keith (2004). There are 16 vegetation formations and sub-formations in NSW.*

**Vegetation Integrity (VI):** *the condition of native vegetation assessed for each vegetation zone against the benchmark for the Plant Community Type.*

**Vegetation Integrity (VI) Score:** *the quantitative measure of vegetation condition calculated in accordance with Equation 23 or Equation 24.*

**Vegetation Zone:** *a relatively homogenous area of native vegetation on a development site, land to be biodiversity certified or a biodiversity stewardship site that is the same Plant Community Type and broad condition state.*

## EXECUTIVE SUMMARY

Maxwell Ventures (Management) Pty Ltd, a wholly owned subsidiary of Malabar Resources Limited (Malabar), is seeking to modify Development Consent SSD 9526 (the Modification) for the approved Maxwell Underground Mine Project (the Project). The Project is east-southeast of Denman and south-southwest of Muswellbrook in New South Wales (NSW).

A Biodiversity Development Assessment Report (BDAR) was prepared by Hunter Eco for the Project in July 2019 and Development Consent SSD 9526 was granted in December 2020.

This BDAR was prepared by Dr Colin Driscoll (Hunter Eco) to assess the likely biodiversity impacts from a Modification to the Project. The Modification is located wholly within the approved Development Application Area and would comprise the following components:

- repositioning of the underground portal;
- realignment of a small section of the proposed access road at the Maxwell Infrastructure site to utilise an existing haul road and minimise impacts to established mine rehabilitation;
- repositioning of an approved clean water diversion for the MEA;
- repositioning of the water treatment facility from the MEA to the Maxwell Infrastructure; and
- other minor works and ancillary infrastructure components within existing/approved surface development areas (e.g. works associated with the reconfiguration of the MEA, pumps/pipelines associated with the water treatment facility).

### **Native Vegetation**

This BDAR assesses some relatively minor modifications to the approved Project layout, using the extensive information from the previous BDAR along with supplementary sampling. The total amount of native vegetation to be disturbed for the Modification is approximately 4.4 ha (i.e. 2.7 % of the native vegetation approved to be disturbed for the Project).

The revised site access road alignment follows an existing haul road and does not contain native vegetation or threatened species habitat, as such, it is not assessed in this BDAR. The native vegetation in the extension to the portal entry and extension to the clean water diversion bank consists of derived native grassland or re-growth Buloke woodland (also known as Bull Oak). The majority of the water treatment facility location has been disturbed by past mining activities with small fragmented groups of Buloke trees.

Despite the degraded nature of the vegetation present compared to the woodland/forest vegetation that were once present, the 3.8 ha of PCT 1606 (in derived native grassland form) meets the criteria for the *White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions* listed as a threatened ecological community (TEC) under the *NSW Biodiversity Conservation Act 2016* (BC Act) and the *White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland* listed as a TEC under the *Commonwealth Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) (collectively referred to as the Box-Gum TEC).

### **Threatened Species**

Targeted surveys for threatened flora were conducted for the Project by Hunter Eco in 2017, 2018 and 2019 and fauna surveys were conducted by Future Ecology in 2017 and 2018. Additional targeted flora surveys for the Modification were conducted by Hunter Eco in June 2021.

Two 'species credit species' (as defined by the *Threatened Biodiversity Data Collection*) are relevant to the Modification as they are known to be present in habitat adjoining the Biodiversity Assessment Development Footprint, namely the:

- Striped Legless Lizard (*Delmar impar*); and
- Squirrel Glider (*Petaurus norfolcensis*).

No threatened flora species or populations listed under the BC Act or EPBC Act were recorded in the Modification Biodiversity Assessment Development Footprint.

### **Measures to Avoid, Minimise, Mitigate and Manage Impacts**

The extension to the portal entry involves some additional disturbance of derived native grassland (Box-Gum TEC), but avoids woodland/forest vegetation communities. Similarly, the extension to the clean water diversion bank involves some additional disturbance of derived native grassland (Box-Gum TEC), and also largely avoids woodland/forest vegetation communities to the south, except for a partial impact (0.5 ha) on a patch of re-growth Buloke woodland (mapped as PCT 1692).

The Modification would also minimise the impacts to the existing rehabilitation at the Maxwell Infrastructure through the realignment of the site access road along an existing internal haul road, without native vegetation. The proposed location of the water treatment facility has been disturbed by past mining activities with small fragmented groups of Buloke trees (0.1 ha).

The Modified Project would not directly impact species, or their habitats, listed under the NSW *Fisheries Management Act 1994*. The Modified Project is also unlikely to indirectly impact these species, or the habitats because the impact of the Modification on catchment excision (and loss of flows) in Saddlers Creek would be negligible and would not be measurable.

Any impacts on protected matters listed under the EPBC Act would be localised and negligible on a regional, state and national scale. It is concluded that there is unlikely to be significant residual impact on any threatened species and communities listed under the EPBC Act as a result of the Modification.

### **Conclusion**

The credit calculation has determined the offset requirement for clearance of native vegetation (PCT 1606 [Box-Gum TEC] and PCT 1692) (ecosystem credit requirement) and the offset requirement for clearance of habitat for the Striped Legless Lizard and Squirrel Glider (species credit requirements).

## 1 INTRODUCTION

Maxwell Ventures (Management) Pty Ltd, a wholly owned subsidiary of Malabar Resources Limited (Malabar), is seeking to modify Development Consent SSD 9526 (the Modification) for the approved Maxwell Underground Mine Project (the Project). The Project is east-southeast of Denman and south-southwest of Muswellbrook in New South Wales (NSW) (Figure 1).

This Biodiversity Development Assessment Report (BDAR) was prepared by Dr Colin Driscoll (Hunter Eco), who is an accredited assessor under the NSW *Biodiversity Conservation Act, 2016* (BC Act) (assessor accreditation BAAS17004).

### 1.1 BACKGROUND

Development Consent SSD 9526 for the Project was granted by the Independent Planning Commission (IPC) on 22 December 2020. The Project was subsequently approved under the Commonwealth *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) in March 2021 (EPBC 2018/8287).

The Project is an underground mining operation that is approved to operate for 26 years (until 2047). It involves the extraction of run-of-mine (ROM) coal from four seams within the Wittingham Coal Measures, using the following underground mining methods:

- Underground bord and pillar mining with partial pillar extraction in the Whynot Seam; and
- Underground longwall extraction in the Woodlands Hill Seam, Arrowfield Seam and Bowfield Seam.


The substantial existing Maxwell Infrastructure will be used for handling, processing and transportation of coal for the life of the Project. Maxwell Infrastructure includes an existing coal handling and preparation plant (CHPP), train load-out facilities and other infrastructure and services (including water management infrastructure, administration buildings, workshops and services). The approved MEA is in a natural valley and will support underground mining and coal handling activities and provide for personnel and materials access.

ROM coal brought to the surface at the MEA will be transported to the Maxwell Infrastructure area. Early ROM coal will be transported via internal roads during the construction and commissioning of a covered overland conveyor system in accordance with Condition A8 of Development Consent SSD 9526. Subsequently, ROM coal will be transported to the Maxwell Infrastructure area via the covered overland conveyor system.

The existing product coal stockpile area at Maxwell Infrastructure will be extended to allow for better management of different product coal blends. An additional ROM stockpile will also be developed adjacent to the CHPP to cater for delivery of ROM coal via the covered overland conveyor.

The Project area comprises the following main domains:


- Maxwell Underground – comprising the approved area of underground mining operations and the mine entry area (MEA) within Mining Lease Application (MLA) 597.
- Maxwell Infrastructure – the area within Coal Lease (CL) 229, ML 1531 and CL 395 comprising the substantial existing infrastructure (including the CHPP) and previous mining areas.
- The transport and services corridor between the Maxwell Underground and Maxwell Infrastructure – the area within CL 229, MLA 597 and MLA 598 comprising the proposed site access road, a covered, overland conveyor, power supply and other ancillary infrastructure and services.
- The realignment of Edderton Road.



SHM-20-04 MOD WEA, ROAD, 200A



Source: NSW Spatial Services (2021)



- LEGEND**
- Mining Operation
  - Proposed Mining Operation
  - Railway
  - Local Government Boundary
  - State Forest
  - National Parks and Wildlife Service Estate
  - Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Indicative Extent of Underground Development

**MALABAR**  
**MAXWELL UNDERGROUND MINE PROJECT**  
 Regional Location

**Figure 1**

A detailed description of the Project is provided in the main document of the EIS. The approved Project general arrangement is shown on Figure 2.

## 1.2 MODIFICATION DESCRIPTION

The Modification would comprise the following components:

- repositioning of the underground portal;
- realignment of a small section of the approved access road at the Maxwell Infrastructure site to utilise an existing haul road and minimise impacts to established mine rehabilitation;
- repositioning of an approved clean water diversion for the MEA;
- repositioning of the water treatment facility from within the approved MEA to Maxwell Infrastructure; and
- other minor works and ancillary infrastructure components within existing/approved surface development areas (e.g. works associated with the reconfiguration of the MEA, pumps/pipelines associated with the water treatment facility).

## 1.3 GENERAL DESCRIPTION OF THE SUBJECT LAND

The Subject land associated with the Modification (including operational and construction footprints) is approximately 14.2 hectares (ha) in size (Figure 3). This total area comprises approximately 9.3 ha for the revised site access road alignment, approximately 4.3 ha for MEA Extensions (the extension to the portal entry and extension to the clean water diversion bank), and approximately 0.6 ha for the water treatment facility. The Subject land is herein referred to as the Biodiversity Assessment Development Footprint.


The revised site access road alignment follows an existing haul road and does not contain native vegetation or threatened species habitat, as such, it is not assessed in this BDAR. The original native vegetation in the extension to the portal entry and extension to the clean water diversion bank consists of derived native grassland or re-growth Buloke woodland (also known as Bull Oak). The majority of the water treatment facility location has been disturbed by past mining activities and is devoid of native vegetation, with small fragmented groups of Buloke trees.

## 1.4 STRUCTURE OF THIS ASSESSMENT

This BDAR is structured as follows:

- |           |  |
|-----------|--|
| Section 2 | Provides landscape features relevant to the Modification.                          |
| Section 3 | Describes native vegetation.   |
| Section 4 | Describes the threatened species assessment.                                       |
| Section 5 | Outlines the prescribed biodiversity impact entities.                              |
| Section 6 | Provides an assessment of how potential impacts would be avoided and/or minimised. |
| Section 7 | Provides an evaluation of potential impacts.                                       |
| Section 8 | Describes the measures taken to mitigate and manage impacts.                       |
| Section 9 | Presents a summary of the assessment and credit requirements.                      |






- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved CHPP Reject Emplacement Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Approved 66 kV Power Supply
  - Ausgrid 66 kV Power Supply Extension

Source: NSW Spatial Services (2021); MSEC (2019)  
 Orthophoto Mosaic: 2020, 2019

**MALABAR**  
 MAXWELL UNDERGROUND MINE PROJECT  
 Approved General Arrangement

**Figure 2**



- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint

Source: NSW Spatial Services (2021);  
MSEC (2019); AECOM (2019)  
Orthophoto Mosaic: 2020, 2019

**MALABAR**  
**MAXWELL UNDERGROUND MINE PROJECT**  
**Proposed General Arrangement**

**Figure 3**

## 1.5 INFORMATION SOURCES USED IN THIS ASSESSMENT

### 1.5.1 Field Survey Reports

Flora surveys were conducted encompassing the approved surface development areas and the Biodiversity Assessment Development Footprint associated with the Modification by Hunter Eco in 2017, 2018 and 2019 (Hunter Eco 2019a) and fauna surveys were conducted by Future Ecology in 2017 and 2018 (Future Ecology 2019) (Attachment A).

Supplementary surveys were undertaken by Hunter Eco in June 2021 to verify Plant Community Types (PCTs), collect additional Vegetation Integrity (VI) field plots from within the Biodiversity Assessment Development Footprint associated with the Modification and conduct additional targeted surveys for threatened flora species.

### 1.5.2 Published Databases

Published databases used in this assessment include:

- *Threatened Biodiversity Data Collection (TBDC)* (DPIE 2021a)<sup>1</sup>;
- *BioNet Vegetation Classification* (DPIE 2021b);
- *BioNet Atlas* (DPIE 2021c)<sup>2</sup>; and
- *Directory of Important Wetlands of Australia* (Department of the Environment and Energy [DEE] 2018a).

A full reference list of all information sources used in this BDAR is provided in Section 11.

### 1.5.3 Local Data

It was not necessary to use local data or deviate from the DPIE databases (DPIE 2021a, 2021b).

### 1.5.4 BAM-C

BAM-C Version 1.3.0.00 (last updated 29 March 2021) was used in this assessment.

As described above, the revised site access road alignment follows an existing haul road and does not contain native vegetation or habitat for threatened species. Therefore, the revised site access road alignment was not assessed in the BAM-C.

The MEA Extensions (extension to the portal entry and clean water diversion bank) were assessed in one BAM-C calculation. The proposed water treatment facility was assessed in a separate BAM-C calculation as it is more distant (approximately 9 kilometres [km] from the extension areas) and lies in a very different, highly modified, landscape.

---

<sup>1</sup> This website is titled 'Profiles'.


<sup>2</sup> This website is titled 'Species Sightings Search'.

## 2 LANDSCAPE FEATURES

This section provides a description of the landscape features relevant to the Biodiversity Assessment Development Footprint in accordance with the BAM (DPIE 2020).

### 2.1 TOPOGRAPHY, HYDROLOGICAL SETTING, GEOLOGY AND SOILS

Figure 4 shows that the modified MEA is located at the foot of a ridge and the clean water diversion is located further up the same slope. Both areas do not intersect drainage lines which are ephemeral first order streams (Strahler stream ordering, Department of Primary Industries – Water 2017]) that flow north west into Saddlers Creek which flows into the Hunter River.



**Figure 4 The Topography of the Mine Entry Components in Relation to the Approved Development**

The Modification overlies Permian age sedimentary geology consisting of coal seams, claystone tuff, siltstone and sandstone. The Australian Soil Classification (Isbell 2016) map shows soil type in the MEA extensions to be Vertosol. Soil type at the water treatment plant area is no longer original having been disturbed by past mining activities.

### 2.2 NATIVE VEGETATION COVER

The proposed MEA Extensions consist entirely of native vegetation, predominantly derived native grassland (3.8 ha) along with woodland (0.5 ha). The proposed water treatment facility location consists of a fragment of remnant woodland (0.1 ha) with the remainder being a combination of bare earth and old mine workings stabilised with exotic vegetation.

A site-based assessment method described in the BAM, whereby a 1,500 m buffer is placed around the development site was applied to the Modification separately for the proposed MEA Extensions and proposed water treatment facility location components. The extent of native vegetation cover and patch size has been assessed (Table 1).

**Table 1**  
**Native Vegetation Extent in the Buffer Area**

Component	Native Vegetation Extent in the Buffer Area (%)	Patch Size (ha)
Proposed MEA Extensions	99.7	>100
Proposed Water Treatment Facility Location	7	0.1

Areas not shown as native vegetation on the Site Map (Figure 5) are cleared of native vegetation. These cleared areas are further identified in Section 3.4. There were no notable differences between mapped vegetation extent and aerial imagery.

### 2.3 IBRA BIOREGIONS AND SUB-REGIONS

In accordance with the BAM (DPIE 2020), the Site Map is shown on Figure 5 and the Location Map for the Biodiversity Assessment Development Footprint is shown on Figure 6.

The Modification area lies within the Sydney Basin Interim Biogeographic Regionalisation for Australia (IBRA) Bioregion, Hunter IBRA sub-region. The IBRA regional boundaries (Department of the Environment [DotE] 2012) do not occur near the Biodiversity Assessment Development Footprint and hence are not shown on Figures 5 and 6.

### 2.4 RIVERS AND STREAMS

Rivers and streams (and riparian buffer distances based on Strahler stream ordering [Department of Primary Industries – Water 2017]) are shown on Figure 5. The Hunter River, Saltwater Creek and Ramrod Creek are downstream of the Modification. The upper reaches of Saddlers Creek are immediately north of the underground mining area with the creek continuing south-west to the Hunter River.

### 2.5 WETLANDS


There are no important or local wetlands on or, adjacent to the Biodiversity Assessment Development Footprint (after DEE 2018a). The closest important wetland is too far away (over 50 km) to be shown on Figure 6.

### 2.6 HABITAT CONNECTIVITY

Native vegetation extent and habitat connectivity were determined by site survey and current aerial photography (Nearmap 2019). Connectivity of woodland/forest habitat was assessed where gaps between discrete patches were 100 metres (m) or less and native grassland habitat where gaps were 30 m or less. The woodland/forest habitat is fragmented due to past land clearance.

### 2.7 KARST, CAVES, CREVICES, CLIFFS, ROCKS AND OTHER GEOLOGICAL FEATURES

There are no karst, caves, cliffs or other areas of geological significance on, or in the vicinity of, the Modification.




- LEGEND**
- Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Subject Land
  - Project Cadastral Boundary
  - Existing Conservation/Offset Area
  - Habitat Connectivity Area
  - Non-native Vegetation
  - Strahler Stream Order**
  - 1st Order (Riparian Corridor Width 10 m)
  - 2nd Order (Riparian Corridor Width 20 m)
  - 3rd Order (Riparian Corridor Width 30 m)
  - 4th Order (Riparian Corridor Width 40 m)
  - Mitchell Landscapes**
  - Chv Central Hunter Alluvial Plains
  - Chf Central Hunter Foothills
  - Est/War Estuary/Water Added
  - Hrb Hunter River Basalts
  - Uhc Upper Hunter Channels and Floodplain

Source: NSW Spatial Services (2021); MSEC (2019)  
 Orthophoto Mosaic: 2020, 2019



**Figure 5**



- LEGEND**
- Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Subject Land
  - Assessment Area (1500 m Buffer Area)
  - Woodland within the Buffer
  - Derived Native Grassland within the Buffer
  - Habitat Connectivity Area outside the Buffer
  - Local Government Area
  - Strahler Stream Order
  - 1st Order (Riparian Corridor Width 10 m)
  - 2nd Order (Riparian Corridor Width 20 m)
  - 3rd Order (Riparian Corridor Width 30 m)
  - 4th Order (Riparian Corridor Width 40 m)
  - Mitchell Landscapes**
  - Chv Central Hunter Alluvial Plains
  - Chf Central Hunter Foothills
  - Est/Wat Estuary/Water Added
  - Hrb Hunter River Basalts
  - Uhc Upper Hunter Channels and Floodplain

Source: NSW Spatial Services (2021); MSEC (2019)  
Orthophoto Mosaic: 2020, 2019

Figure 6

## 2.8 AREAS OF OUTSTANDING BIODIVERSITY VALUE

There are no Areas of Outstanding Biodiversity Value listed under the NSW *Biodiversity Conservation Regulation, 2017* (BC Regulation) associated with the Modification.

## 2.9 NSW MITCHELL LANDSCAPE

The Modification components lie entirely within the Central Hunter Foothills Mitchell landscape (Mitchell 2002).



### 3 NATIVE VEGETATION

This section provides a description of the native vegetation relevant to the Biodiversity Assessment Development Footprint associated with the Modification.

#### 3.1 EXISTING INFORMATION ON NATIVE VEGETATION

This BDAR has built upon the extensive flora and vegetation survey and assessment works completed for the *Maxwell Project Baseline Flora Report* (Hunter Eco 2019a) and *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b). A desktop assessment of the Modification disturbance areas was conducted to determine what information was already available from the baseline flora report (Hunter Eco 2019a) and any extra data that were needed to be collected by field survey.

#### 3.2 SUPPLEMENTARY SURVEYS


Supplementary flora surveys were undertaken by Dr Colin Driscoll (Hunter Eco) on 28 June 2021. The purpose of the supplementary flora surveys was to verify PCTs, collect additional Vegetation Integrity (VI) plots and conduct additional targeted surveys for threatened flora species.

A total of four VI plots were sampled from within the Biodiversity Assessment Development Footprints (Figures 7a and 7b). Section 3.5.3 describes the VI plots used in the BAM-C. A description of the surveys undertaken for threatened flora species is provided in Section 4.4.

#### 3.3 NATIVE VEGETATION EXTENT

The proposed extension to the portal entry is entirely grassland derived from PCT 1606 as mapped by Hunter Eco (2019). Scattered regrowth *Acacia salicina* is present but no eucalypts. The grasses are dominated by *Dichanthium sericeum* along with scattered patches of *Aristida ramosa*, *Bothriochloa biloba*, *Sporobolus creber*, *Eulalia aurea*, *Chloris divaricata* and *Digitaria divaricatissima*. Herbs consists of *Asperula conferta*, *Chrysocephalum semipapposum*, *Vittadinia* species and *Calotis lappulacea*. Twiners are *Glycine clandestina*, *Desmodium varians* and *Oxytes brachypoda*. Weeds are sparse and included *Carthamus lanatus* and *Senecio madagascariensis*. The grass cover is relatively open varying from a few centimetres to under 1 m tall; small herbs are readily visible without needing to part the grasses.

The proposed extension to the clean water diversion bank consists partly of derived grassland similar in content and structure to that of the proposed extension to the portal entry, along with a patch of PCT 1692 Buloke woodland. The woodland is dominated by *Allocasuarina luehmannii* with scattered *Acacia salicina*; there are no eucalypts. The ground cover is sparse and consists of low shrubs *Cryptandra amara*, *Eremophila debilis* and *Maireana microphylla*. Grasses and grass-like species are *Rytidosperma caespitosum*, *Aristida ramosa*, *Chloris ventricosa*, *Eragrostis alveiformis*, *Bothriochloa decipiens*, *Eulalia aurea*, *Lomandra multiflora*, *Lomandra filiformis* subsp. *coriacea* and *Lomandra filiformis* subsp. *filiformis*. Herbs consists of *Veronica plebeia*, *Brunoniella australis*, *Einadia nutans*, *Enchylaena tomentosa* and *Glossocardia bidens*. Twiners are *Desmodium varians* and *Convolvulus graminetinus*. Weeds are sparse and included *Galenia pubescens*, and *Senecio madagascariensis*.



**LEGEND**

- Exploration Licence Boundary
- Mining Lease Application Boundary
- Approved Surface Development Area
- Approved Extent of Conventional Subsidence from Underground Mining
- Biodiversity Assessment Development Footprint
- Vegetation Integrity Plot Location
- Survey Track
- Dry Sclerophyll Forests (Shrub/grass sub-formation)
- White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)<sup>1</sup>
- 2a White Box - Ironbark - Red Gum Shrubby Forest - Derived Native Grassland (PCT1606)<sup>1</sup>
- Dry Sclerophyll Forests (Shrubby sub-formation)
- 3 Slaty Box Shrubby Woodland (PCT1655)
- Forested Wetlands
- 4 Swamp Oak Forest (PCT1731)
- Grassy Woodlands
- 6 Bull Oak Grassy Woodland (PCT1692)
- 7 Yellow Box - Apple Grassy Woodland (PCT1693)
- 9 Ironbark - Grey Box Grassy Woodland (PCT1691)
- 9a Ironbark - Grey Box Grassy Woodland - Derived Native Grassland (PCT1691)
- Other
- C Waterbody/Dam

<sup>1</sup> Listed BC Act, White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions


<sup>2</sup> Listed BC Act, Hunter Valley Footslopes Slaty Gum Woodland in the Sydney Basin Bioregion

<sup>3</sup> Listed BC Act, Central Hunter Grey Box - Ironbark Woodland in the NSW North Coast and Sydney Basin Bioregions

Source: Hunter Eco (2021); MSEC (2019); NSW Spatial Services (2021)  
 Orthophoto: ESRI World Imagery (2021)

MALABAR  
 MAXWELL UNDERGROUND MINE PROJECT  
 Vegetation Mapping  
 - Mine Entry Area

Figure 7a



SHW-20-04/NOO/MEA\_BDAR\_20/68

- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Biodiversity Assessment Development Footprint
  - Vegetation Integrity Plot Location
  - Dry Sclerophyll Forests (Shrub/grass sub-formation)
  - 1 Red Gum - Ironbark - Apple Shrubby Woodland (PCT1607)
  - 1a Red Gum - Ironbark - Apple Shrubby Woodland - Derived Native Grassland (PCT1607)
  - 2a White Box - Ironbark - Red Gum Shrubby Forest - Derived Native Grassland (PCT1606)<sup>1</sup>
  - Forested Wetlands
  - 5 Hunter Lowland Red Gum Forest (PCT1598)<sup>2</sup>

- 6 Grassy Woodlands
- Bull Oak Grassy Woodland (PCT1692)
- 11 Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)<sup>3</sup>
- 11a Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland - Derived Native Grassland (PCT1604)
- Other**
- C Waterbody/Dam
- D Woodland Rehabilitation
- E Pasture Rehabilitation
- F Infrastructure/Cleared Land

<sup>1</sup> Listed BC Act, White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions

<sup>2</sup> Listed BC Act, Hunter Lowland Redgum Forest

<sup>3</sup> Listed BC Act, Central Hunter Ironbark - Spotted Gum - Grey Box Forest in the NSW North Coast and Sydney Basin Bioregions

Source: Hunter Eco (2021); NSW Spatial Services (2021) Orthophoto Mosaic: 2020

**MALABAR**  
MAXWELL UNDERGROUND MINE PROJECT  
Vegetation Mapping  
- Maxwell Infrastructure

**Figure 7b**

The Water Treatment Plant consists partly of three fragments of PCT 1692 Buloke woodland remaining after past clearing for an electricity transmission line (ETL), the remainder being bare ground access and work areas and old mine disturbance areas. The woodland is dominated by *Allocasuarina luehmannii* and one isolated *Eucalyptus blakelyi*. Native grass species are *Austrostipa verticillata* and *Aristida ramosa* both at only 0.1% cover. Other scattered native ground cover consists of *Enchylaena tomentosa*, *Maireana microphylla*, *Carex inversa*, *Eremophila debilis*, *Lomandra confertifolia* subsp. *rubiginosa*, and *Dichondra repens* each recorded at 0.1% cover. Weeds are abundant comprising over 100% of cover and included *Galenia pubescens*, *Panicum antidotale*, *Melinis repens*, *Hyparrhenia hirta*, *Gomphocarpus fruticosus*, *Bidens pilosa*, *Verbena officinalis*, *Senecio madagascariensis*, *Chloris gayana*, *Asphodelus fistulosus*, *Lysimachia arvensis*, *Ricinus communis* and *Carthamus lanatus*. Thirty-six metres of fallen logs are present as a result of clearing for the ETL, however these are covered by matted weeds. The old mine disturbance areas had been sown with exotic grasses, primarily *Panicum antidotale*. There are also a small number of exotic *Acacia saligna* shrubs.

### 3.4 AREAS WITHOUT NATIVE VEGETATION

As described above, the revised site access road alignment follows an existing haul road and does not contain native vegetation or habitat for threatened species (Figure 7b). Similarly, the majority of the water treatment facility location has been disturbed by past mining activities and is devoid of native vegetation or habitat for threatened species (Figure 7b).

### 3.5 PLANT COMMUNITY TYPES

#### 3.5.1 Plant Community Types and Classes

Plant Community Types (PCT) on the Modification land and surrounding area in accordance with the BAM (DPIE 2020) and *BioNet Vegetation Classification* (DPIE 2021b) (Figures 7a and 7b) (Table 2). The PCTs are assigned to a vegetation class in Table 2. Table 2 also includes the Percent Cleared Values from the *BioNet Vegetation Classification* (DPIE 2021b) and the Vegetation Integrity (VI) Scores (Section 3.6.4).

The Biodiversity Assessment Development Footprint associated with the Modification is approximately 14.2 hectares (ha) in size comprising (Figures 7a and 7b):

- 0.6 ha of fragmented (i.e. not continuous) native woodland/forest vegetation;
- 3.8 ha of derived native grassland; and
- 9.8 ha is infrastructure/cleared land.

#### 3.5.2 Evidence to Support Plant Community Types

Hunter Eco (2019a) justifies the PCT and vegetation zone mapping (including the species relied upon for identification of PCTs). The Biodiversity Assessment Development Footprints are small fragments of much larger areas that were classified and mapped by Hunter Eco (2019) in and around the Project development footprint. The derived native grassland in the Biodiversity Assessment Development Footprint (Plate 1) is part of approximately 1600 ha of continuous derived native grassland classified as PCT 1606 by Hunter Eco (2019) based on 237 paddock trees 80% of which were identified as White Box (outside of the Biodiversity Assessment Development Footprint). The woodland fragments in the Biodiversity Assessment Development Footprint were unambiguously identified as PCT 1692, the only PCT having Buloke (*Allocasuarina luehmannii*) as the dominant upper stratum species (Plates 2 and 3).

The location of vegetation integrity (site condition) plots used in the BAM-C for the Biodiversity Assessment Development Footprint are shown on Figures 7a and 7b.

**3.5.3 Plant Community Types Percent Cleared Value**

The BAM (DPIE 2020) defines ‘Percent Cleared Value’ as the percentage of a PCT that has been cleared as a proportion of its pre-1750 extent, as identified in the *BioNet Vegetation Classification* (DPIE 2021b). Percent cleared values for each PCT are shown in Table 2.



**Plate 1 Proposed MEA Extensions – PCT 1606  
(in derived native grassland form)**



**Plate 2 Proposed MEA Extensions - PCT 1692**



**Plate 3 Proposed Water Treatment Facility Location - PCT 1692**

**Table 2  
Plant Community Type Data**

Vegetation Zone	PCT	PCT Name	Class	Generic Name	Area (ha)	Percent Cleared	Sensitivity Class <sup>^</sup>	VI Score <sup>~</sup>
<b>Proposed MEA Extensions</b>								
<i>Dry Sclerophyll Forests (Shrub/grass sub-formation)</i>								
1	1606	White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter - DNG <sup>1</sup>	North-west Slopes Dry Sclerophyll Woodlands	2a. White Box - Ironbark - Red Gum Shrubby Forest (DNG)	3.8	29%	High	31.4
<i>Grassy Woodlands</i>								
2	1692	Bull Oak Grassy Woodland of the central Hunter Valley	Coastal Valley Grassy Woodlands	6. Bull Oak Grassy Woodland	0.5	53%	High	48.5
<b>Proposed Water Treatment Facility Location</b>								
<i>Grassy Woodlands</i>								
1	1692	Bull Oak Grassy Woodland of the central Hunter Valley*	Coastal Valley Grassy Woodlands	6. Bull Oak Grassy Woodland	0.1	53%	High	38.2
<b>Total Woodland/Forest</b>					0.6	-		-
<b>Total Derived Native Grassland</b>					3.8	-		-
<b>Total</b>					4.4	-		-

<sup>1</sup> Listed as Box-Gum TEC under the BC Act and EPBC Act.

<sup>^</sup> The sensitivity class is set for each PCT by DPIE (2021b).

<sup>~</sup> BAM-C.

### 3.5.4 Threatened Ecological Communities

Threatened ecological communities and associated PCT within the Biodiversity Assessment Development Footprint are listed in Table 3 and shown on Figures 7a and 7b.

**Table 3**  
**Threatened Ecological Communities**

Threatened Ecological Community	Conservation Status*	Associated PCT	Area (ha)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions listed under the BC Act	CEEC	1606	3.8
White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland listed as a TEC under the EPBC Act.			

\* Threatened ecological community status under the BC Act and EPBC Act (current as at August 2021).

CEEC = Critically Endangered Ecological Community.

### 3.5.5 Evidence to Support Threatened Ecological Communities

The TBDC (DPIE 2021a) associates PCT 1606 woodland or derived native grassland with the Critically Endangered Ecological Community *White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions* referred to as the Box-Gum threatened ecological community (TEC). As noted in Section 3.5.2, Hunter Eco (2019) assessed PCT 1606 on the grounds of species composition in the wider landscape based in part on the dominant presence of White Box, a key component of Box-Gum TEC.

The TBDC associates PCT 1692 with:

- *Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions; or,*
- *Central Hunter Grey Box—Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions.*

Neither of these TEC are a match for PCT 1692 in the Modification as there are no Red Gum, Grey Box or Ironbark present in the canopy in or near the Modification.

## 3.6 VEGETATION INTEGRITY ASSESSMENT

### 3.6.1 Vegetation Zones

Two vegetation zones (i.e. areas of native vegetation that are the same PCT and similar broad condition states) are mapped for the proposed MEA Extensions and one vegetation zone for the proposed water treatment facility location (Table 2; Figures 7a and 7b). Vegetation condition states recognised are: woodland and derived native grassland.

### 3.6.2 Patch Size

Patch size for the vegetation zones associated with proposed MEA Extensions is >100 ha (Figures 5 and 6). Patch size for the vegetation zones associated with the proposed water treatment facility location is 0.1 ha consisting only of the native vegetation within the Biodiversity Assessment Development Footprint as it is separated from the next nearest native vegetation by >100 m.

### 3.6.3 Vegetation Integrity Plots

The baseline flora surveys collected data from four plots located to sample the vegetation zones present with the number of plots meeting the requirements per area of each zone (Attachment B). For the MEA two additional plots (one each from PCT 1606 and 1692) collected in the immediate vicinity of the Biodiversity Assessment Development Footprints during earlier surveys (Hunter Eco 2019) were included in the BAM-C data input in order to provide a more representative sample of the habitat.

### 3.6.4 Vegetation Integrity Score

The BAM-C was used to determine the VI Scores for each vegetation zone (Table 4). According to the BAM-C, all of the vegetation zones have a VI Score requiring an offset (Table 4). This is because the VI Scores are greater than 17 for PCTs associated with species habitat and greater than 15 for PCTs representative of a critically endangered ecological community (i.e. Vegetation Zone 1 of the Proposed MEA Extensions).



**Table 4  
Vegetation Integrity Score Detail**

Vegetation Zone	PCT	PCT-condition class	Composition Condition Score	Structure Condition Score	Function Condition Score	Trees with Hollows	Vegetation Integrity Score	Threshold for Requiring an Offset	Offset Required?
<b>Proposed MEA Extensions</b>									
1	1606*	Derived Native Grassland	53.4	26.8	21.7	0	31.4	≥15	Yes
2	1692	Moderate	72.6	35.9	43.8	0	48.5	≥17	Yes
<b>Proposed Water Treatment Facility Location</b>									
1	1692	Low	27	35.3	58.6	0	38.2	≥17	Yes

\* Listed as Box-Gum TEC under the BC Act and EPBC Act.

## 4 THREATENED SPECIES

Threatened species that are 'ecosystem credit species' and/or 'species credit species' are pre-determined by DPIE in the BAM-C and TBDC (DPIE 2021a). The BAM (DPIE 2020) states:

*'Ecosystem credit species' are threatened species whose occurrence can generally be predicted by vegetation surrogates and/or landscape features, or that have a low probability of detection using targeted surveys. The TBDC identifies the threatened species assessed for ecosystem credits. A targeted survey is not required to identify or confirm the presence of ecosystem credit species.*

...

*'Species credit species' are threatened species for which vegetation surrogates and/or landscape features cannot reliably predict the likelihood of their occurrence or components of their habitat. These species are identified in the TBDC. A targeted survey or an expert report is required to confirm the presence of these species on the subject land. Alternatively, for a development, activity, clearing or biodiversity certification proposal only, the proponent may elect to assume the species is present. (This option must not be applied to proposed biodiversity stewardship sites).*

...

*In some circumstances the TBDC may identify a threatened species that requires assessment for ecosystem credits and species credits (referred to as dual credit species). For dual credit species, part of the habitat is assessed as a species credit (e.g. breeding habitat or land mapped on an important habitat map for a species). The remaining habitat components for the species are assessed as an ecosystem credit (e.g. foraging habitat). Dual credit species are generally: a. highly mobile species that rely on particular habitat components for breeding, such as maternity caves for bats, tree hollows for some large forest owls or cockatoos, or b. species for which particular areas in the landscape are important for their survival, such as selected beaches for migratory shorebirds.*

### 4.1 ECOSYSTEM CREDIT SPECIES - HABITAT SUITABILITY ASSESSMENT

In accordance with the BAM (DPIE 2020), assessing habitat suitability for an ecosystem credit species involves the following steps:

Step 1: Identify threatened species for assessment; and

Step 2: Assess the habitat constraints and vagrant species on the Biodiversity Assessment Development Footprint.

These steps are applied below.

#### 4.1.1 Ecosystem Species from the BAM-C

A total of 33 ecosystem credit species for assessment are listed in Table 5 from the BAM-C. All 33 species listed in Table 5 were identified in BAM-C for the proposed MEA Extensions, while 14 were identified in BAM-C for the proposed water treatment facility location.

**Table 5**  
**Ecosystem Species from the BAM-C**

Scientific Name	Common Name	Conservation Status <sup>1</sup>		Class of Credit <sup>2</sup>	Sensitivity to Gain Class	Proposed MEA Extensions	Proposed Water Treatment Facility Location
		BC Act	EPBC Act				
<b>Birds</b>							
<i>Lophoictinia isura</i>	Square-tailed Kite	V	-	S/E	Moderate	•	-
<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle	V	-	S/E	High	•	•
<i>Hieraaetus morphnoides</i>	Little Eagle	V	-	S/E	Moderate	•	-
<i>Calyptorhynchus lathami</i>	Glossy Black-Cockatoo	V	-	S/E	High	•	•
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo	V	-	S/E	Breeding: High Foraging: Moderate	•	-
<i>Glossopsitta pusilla</i>	Little Lorikeet	V	-	E	High	•	•
<i>Neophema pulchella</i>	Turquoise Parrot	V	-	E	High	•	-
<i>Lathamus discolor</i>	Swift Parrot	E	CE	S/E	Moderate	•	•
<i>Tyto novaehollandiae</i>	Masked Owl	V	-	S/E	High	•	-
<i>Ninox strenua</i>	Powerful Owl	V	-	S/E	High	•	-
<i>Ninox connivens</i>	Barking Owl	V	-	S/E	High	•	-
<i>Hirundapus caudacutus</i>	White-throated Needletail	-	V	E	High	•	•
<i>Climacteris picumnus victoriae</i>	Brown Treecreeper (eastern subspecies)	V	-	E	High	•	-
<i>Chthonicola sagittata</i>	Speckled Warbler	V	-	E	High	•	•
<i>Melithreptus gularis gularis</i>	Black-chinned Honeyeater (eastern subspecies)	V	-	E	Moderate	•	-
<i>Anthochaera phrygia</i>	Regent Honeyeater	CE	CE	S/E	High	•	-
<i>Grantiella picta</i>	Painted Honeyeater	V	V	E	Moderate	•	-
<i>Melanodryas cucullata cucullata</i>	Hooded Robin (south-eastern form)	V	-	E	Moderate	•	•
<i>Petroica phoenicea</i>	Flame Robin	V	-	E	Moderate	•	•
<i>Petroica boodang</i>	Scarlet Robin	V	-	E	Moderate	•	•
<i>Pomatostomus temporalis temporalis</i>	Grey-crowned Babbler (eastern subspecies)	V	-	E	Moderate	•	•

Scientific Name	Common Name	Conservation Status <sup>1</sup>		Class of Credit <sup>2</sup>	Sensitivity to Gain Class	Proposed MEA Extensions	Proposed Water Treatment Facility Location
		BC Act	EPBC Act				
<i>Daphoenositta chrysoptera</i>	Varied Sittella	V	-	E	Moderate	•	-
<i>Stagonopleura guttata</i>	Diamond Firetail	V	-	E	Moderate	•	-
<b>Mammals</b>							
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	V	E	E	High	•	•
<i>Phascolarctos cinereus</i>	Koala	V	V	S/E	High	•	-
<i>Petaurus australis</i>	Yellow-bellied Glider	V	-	E	High	•	-
<i>Pteropus poliocephalus</i>	Grey-headed Flying-fox	V	V	S/E	High	•	•
<i>Saccolaimus flaviventris</i>	Yellow-bellied Sheath-tail-bat	V	-	E	High	•	-
<i>Micronomus norfolkensis</i>	Eastern Coastal Free-tailed Bat	V	-	E	High	•	•
<i>Miniopterus orianae oceanensis</i>	Large Bentwing-bat	V	-	S/E	Breeding: Very High Foraging: High	•	•
<i>Nyctophilus corbeni</i>	Corben's Long-eared Bat	V	V	E	High	•	-
<i>Falsistrellus tasmaniensis</i>	Eastern False Pipistrelle	V	-	E	High	•	-
<i>Scoteanax rueppellii</i>	Greater Broad-nosed Bat	V	-	E	High	•	-

<sup>1</sup> Threatened fauna species status under the BC Act and/or EPBC Act (current as at August 2021).

V = Vulnerable; E = Endangered; CE = Critically Endangered.

<sup>2</sup> Biodiversity credit class under the TBDC (DPIE 2021a) (current as at August 2021).  
E = Ecosystem; S = Species.

#### 4.1.2 Exclusion of Ecosystem Species

Table 6 details ecosystem species excluded from the BAM-C based on habitat constraints or PCT association.

**Table 6**  
**Ecosystem Species Excluded from the BAM-C**

Scientific Name	Common Name	Conservation Status <sup>1</sup>		Class of Credit <sup>2</sup>	Proposed MEA Extensions	Proposed Water Treatment Facility Location
		BC Act	EPBC Act			
<b>Birds</b>						
<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle	V	-	S/E	Not associated with PCT 1692 because MEA >1 km from Saddler's Creek	
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo	V	-	S/E	Woodland species not associated with PCT 1606 DNG; potentially associated with PCT 1692	N/A
<i>Calyptorhynchus lathamii</i>	Glossy Black-Cockatoo	V	-	S/E	Woodland species not associated with PCT 1606 DNG; potentially associated with PCT 1692 and Buloke is a feed tree species	Potentially associated with 1692 because it is a woodland species and Buloke is a feed tree species
<i>Glossopsitta pusilla</i>	Little Lorikeet	V	-	E	Woodland species not associated with PCT1606 DNG; potentially associated with PCT 1692	Potentially associated with 1692 because it is a woodland species
<i>Lathamus discolor</i>	Swift Parrot	E	CE	S/E	No foraging habitat	No foraging habitat
<i>Climacteris picumnus victoriae</i>	Brown Treecreeper (eastern subspecies)	V	-	E	Woodland species not associated with PCT 1692 or DNG form of PCT 1606	N/A
<i>Anthochaera phrygia</i>	Regent Honeyeater	CE	CE	S/E	No foraging habitat	N/A
<i>Grantiella picta</i>	Painted Honeyeater	V	V	E	Woodland species not associated with DNG form of PCT 1606 or PCT 1692 due to low mistletoe count	N/A

Scientific Name	Common Name	Conservation Status <sup>1</sup>		Class of Credit <sup>2</sup>	Proposed MEA Extensions	Proposed Water Treatment Facility Location
		BC Act	EPBC Act			
<b>Mammals</b>						
<i>Phascolarctos cinereus</i>	Koala	V	V	S/E	Not associated with DNG form of PCT 1606, no eucalypts present in PCT 1692	N/A (Isolated and degraded habitat)
<i>Petaurus australis</i>	Yellow-bellied Glider	V	-	E	Not associated with DNG form of PCT 1606, no hollow-bearing trees present in PCT 1692	N/A (No hollow-bearing trees).
<i>Pteropus poliocephalus</i>	Grey-headed Flying-fox	V	V	S/E	Not associated with DNG form of PCT 1606, no potential feed trees present in PCT 1692	No potential feed trees present in PCT 1692

<sup>1</sup> Threatened fauna species status under the BC Act and/or EPBC Act (current as at August 2021).  
V = Vulnerable; E = Endangered; CE = Critically Endangered.

<sup>2</sup> Biodiversity credit class under the TBDC (DPIE 2021a) (current as at August 2021).  
E = Ecosystem; S = Species.

### 4.1.3 Addition of Ecosystem Species

No ecosystem species were added to the BAM-C.

## 4.2 SPECIES CREDIT SPECIES - HABITAT SUITABILITY ASSESSMENT

Assessing habitat suitability for a species credit species involves the following steps:

- Step 1: Identify species credit species for assessment.
- Step 2: Assess the habitat constraints and vagrant species for species credit species on the Biodiversity Assessment Development Footprint.
- Step 3: Further assessment of candidate species credit species.
- Step 4: Determine the presence of a candidate species credit species.
- Step 5: Determine the area or count, and location of suitable habitat for a species credit species (a species polygon).
- Step 6: Determine the habitat condition within the species polygon for species assessed by area.

#### 4.2.1 Species Credit Species from the BAM-C

A total of 38 species credit species from the BAM-C are listed in Table 7 for assessment. All 38 species listed in Table 7 were identified in BAM-C for the proposed MEA Extensions, while 13 were identified in BAM-C for the proposed water treatment facility location.

**Table 7  
Species Credit Species for Assessment**

Scientific Name	Common Name	Conservation Status <sup>1</sup>		Class of Credit <sup>2</sup>	Proposed MEA Extensions	Proposed Water Treatment Facility Location
		BC Act	EPBC Act			
<b>Flora</b>						
<i>Acacia pendula</i> – endangered population	<i>Acacia pendula</i> population in the Hunter catchment	E	-	S	•	•
<i>Cryptostylis hunteriana</i>	Leafless Tongue Orchid	V	V	S	•	-
<i>Cymbidium canaliculatum</i> – endangered population	<i>Cymbidium canaliculatum</i> population in the Hunter Catchment	E	-	S	•	•
<i>Cynanchum elegans</i>	White-flowered Wax Plant	E	E	S	•	-
<i>Diuris tricolor</i>	Pine Donkey Orchid	V/EP	-	S	•	-
<i>Eucalyptus glaucina</i>	Slaty Red Gum	V	V	S	•	•
<i>Monotaxis macrophylla</i>	Large-leafed Monotaxis	E	-	S	•	-
<i>Ozothamnus tessellatus</i>	Ozothamnus tessellatus	V	V	S	•	-
<i>Pomaderris bodalla</i>	Bodalla Pomaderris	V	-	S	•	-
<i>Pomaderris queenslandica</i>	Scant Pomaderris	E	-	S	•	-
<i>Thesium australe</i>	Austral Toadflax	V	V	S	•	-
<b>Amphibians</b>						
<i>Litoria aurea</i>	Green and Golden Bell Frog	E	V	S	•	•
<b>Reptiles</b>						
<i>Aprasia parapulchella</i>	Pink-tailed Legless Lizard	V	V	S	•	•
<i>Delma impar</i>	Striped Legless Lizard	V	V	S	•	•
<i>Hoplocephalus bitorquatus</i>	Pale-headed Snake	V	-	S	•	-
<b>Birds</b>						
<i>Lophoictinia isura</i>	Square-tailed Kite	V	-	S/E	•	-
<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle	V	-	S/E	•	•
<i>Hieraetus morphnoides</i>	Little Eagle	V	-	S/E	•	-
<i>Burhinus grallarius</i>	Bush Stone-curlew	E	-	S	•	-
<i>Calyptorhynchus lathami</i>	Glossy Black-Cockatoo	V	-	S/E	•	•
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo	V	-	S/E	•	-

Scientific Name	Common Name	Conservation Status <sup>1</sup>		Class of Credit <sup>2</sup>	Proposed MEA Extensions	Proposed Water Treatment Facility Location
		BC Act	EPBC Act			
<i>Lathamus discolor</i>	Swift Parrot	E	CE	S/E	•	•
<i>Tyto novaehollandiae</i>	Masked Owl	V	-	S/E	•	-
<i>Ninox strenua</i>	Powerful Owl	V	-	S/E	•	-
<i>Ninox connivens</i>	Barking Owl	V	-	S/E	•	-
<i>Anthochaera phrygia</i>	Regent Honeyeater	CE	CE	S/E	•	-
<b>Mammals</b>						
<i>Phascogale tapoatafa</i>	Brush-tailed Phascogale	V	-	S	•	•
<i>Planigale maculata</i>	Common Planigale	V	-	S	•	-
<i>Phascolarctos cinereus</i>	Koala	V	V	S/E	•	-
<i>Cercartetus nanus</i>	Eastern Pygmy-possum	V	-	S	•	-
<i>Petauroides volans</i>	Greater Glider	-	V	S	•	-
<i>Petaurus norfolcensis</i>	Squirrel Glider	V	-	S	•	-
<i>Petrogale penicillata</i>	Brush-tailed Rock-wallaby	E	V	S	•	-
<i>Pteropus poliocephalus</i>	Grey-headed Flying-fox	V	V	S/E	•	•
<i>Miniopterus orianae oceanensis</i>	Large Bentwing-bat	V	-	S/E	•	•
<i>Chalinolobus dwyeri</i>	Large-eared Pied Bat	V	V	S	•	-
<i>Myotis macropus</i>	Southern Myotis	V	-	S	•	•
<i>Vespadelus troughtoni</i>	Eastern Cave Bat	V	-	S	•	-

<sup>1</sup> Threatened flora species status under the BC Act and/or EPBC Act (current as at August 2021).

V = Vulnerable; E = Endangered; CE = Critically Endangered; EP = Endangered Population.

<sup>2</sup> Biodiversity credit class under the TBDC (DPIE 2021a) (current as at August 2021).

E = Ecosystem; S = Species.

## 4.2.2 Exclusion of Species Credit Species

### 4.2.2.1 Geographical Constraints

Geographic constraints are identified in the TBDC (DPIE 2021a) for some species credit species (Table 8). None of the geographic constraints are relevant to the Biodiversity Assessment Development Footprint.

### 4.2.2.2 Habitat Constraints and Vagrancy

Habitat constraints are identified in the TBDC (DPIE 2021a) for some species credit species, and the absence of identified habitat precludes the species from further assessment (Table 9).



**Table 8**  
**Flora Species Credit Species – Geographic Constraints**

Scientific Name	Common Name	Geographic Constraint within the Hunter Sub-zone in the BAM Calculator	Assessment
<i>Acacia pendula</i> – endangered population	<i>Acacia pendula</i> population in the Hunter catchment	Within Hunter River catchment	Not a relevant constraint.
<i>Cryptostylis hunteriana</i>	Leafless Tongue Orchid	None	-
<i>Cymbidium canaliculatum</i> – endangered population	<i>Cymbidium canaliculatum</i> in the Hunter Catchment	Must be within Hunter catchment as defined by Australia’s River Basins (Geoscience Australia 1997)	Not a relevant constraint.
<i>Cynanchum elegans</i>	White-flowered Wax Plant	None	-
<i>Diuris tricolor</i>	Pine Donkey Orchid	None	-
<i>Diuris tricolor</i> – endangered population	Pine Donkey Orchid population in the Muswellbrook local government area	Muswellbrook local government area	Not a relevant constraint.
<i>Eucalyptus glaucina</i>	Slaty Red Gum	None	-
<i>Monotaxis macrophylla</i>	Large-leafed Monotaxis	None	-
<i>Ozothamnus tessellatus</i>	-	None	-
<i>Pomaderris bodalla</i>	Bodalla Pomaderris	None	-
<i>Pomaderris queenslandica</i>	Scant Pomaderris	None	-
<i>Thesium australe</i>	Austral Toadflax	None	-

Shaded species are species that have geographical constraints within the Hunter sub-zone.

**Table 9**  
**Species Credit Species - Habitat Constraints**

Common Name	Credit Class	Habitat Constraints identified in the TBDC (DPIE 2021a)	Assessment Prior to the Surveys
<b>Flora</b>			
<i>Acacia pendula</i> population in the Hunter catchment	Species	<i>None.</i>	-
Leafless Tongue Orchid	Species	<i>None.</i>	-
<i>Cymbidium canaliculatum</i> in the Hunter Catchment	Species	<i>Epiphytic in a range of eucalypts, Acacia and Angophora; Cut stumps or logs on ground.</i>	No suitable habitat in PCT 1606 DNG. Potential habitat in PCT 1692.
White-flowered Wax Plant	Species	<i>None.</i>	-
Pine Donkey Orchid	Species	<i>None.</i>	-
Pine Donkey Orchid population in the Muswellbrook local government area	Species	<i>None.</i>	-
Slaty Red Gum	Species	<i>None.</i>	-
Large-leafed Monotaxis	Species	<i>Species is a short-lived annual, and will not be present unless a recent disturbance/fire event has occurred and triggered germination.</i>	Not associated with PCT 1606 DNG or PCT 1692
<i>Ozothamnus tessellatus</i>	Species	<i>None.</i>	-
Bodalla Pomaderris	Species	<i>None.</i>	-
Scant Pomaderris	Species	<i>None.</i>	-
<i>Thesium australe</i>	Species	<i>None.</i>	-
<b>Amphibians</b>			
Green and Golden Bell Frog	Species	<i>Semi-permanent/ephemeral wet areas (within 1km of wet areas). Swamps (within 1km of swamp). Waterbodies (within 1km of waterbody).</i>	Dams (waterbodies) present
<b>Reptiles</b>			
Pink-tailed Legless Lizard	Species	<i>Rocky areas or within 50 m of rocky areas.</i>	Habitat absent
Striped Legless Lizard	Species	<i>None.</i>	-

Common Name	Credit Class	Habitat Constraints identified in the TBDC (DPIE 2021a)	Assessment Prior to the Surveys
Pale-headed Snake	Species	<i>None.</i>	-
<b>Birds</b>			
Square-tailed Kite	Species/Ecosystem	* <i>Breeding constraint: Other (Nest trees).</i>	Habitat absent, no large trees
White-bellied Sea-Eagle	Species/Ecosystem	* <i>Breeding constraint: Other (Living or dead mature trees within suitable vegetation within 1km of a rivers, lakes, large dams or creeks, wetlands and coastlines).</i> * <i>Foraging constraint: Waterbodies (Within 1km of a rivers, lakes, large dams or creeks, wetlands and coastlines).</i>	Habitat absent, no large trees
Little Eagle	Species/Ecosystem	* <i>Breeding constraint: Other (Nest trees - live (occasionally dead) large old trees within vegetation).</i>	Habitat absent, no large trees
Bush Stone-curlew	Species	<i>Fallen/standing dead timber including logs.</i>	Habitat potentially present
Glossy Black-Cockatoo	Species/Ecosystem	* <i>Breeding constraint: Hollow-bearing trees (Living or dead tree with hollows greater than 15cm diameter and greater than 5m above ground).</i>	Habitat absent, no suitable tree hollows
Gang-gang Cockatoo	Species/Ecosystem	* <i>Breeding constraint: Hollow-bearing trees (Eucalypt tree species with hollows greater than 9 cm diameter).</i>	Habitat absent, no suitable tree hollows
Swift Parrot	Species/Ecosystem	* <i>Breeding constraint: Other (As per mapped important areas – contact OEH for information).</i> <i>Foraging constraint: none.</i>	Not a mapped important area
Masked Owl	Species/Ecosystem	* <i>Breeding constraint: Hollow-bearing tree (Living or dead trees with hollows greater than 20cm diameter).</i>	Habitat absent, no suitable tree hollows
Powerful Owl	Species/Ecosystem	* <i>Breeding constraint: Hollow-bearing tree (Living or dead trees with hollows greater than 20cm diameter).</i>	Habitat absent, no suitable tree hollows
Barking Owl	Species/Ecosystem	* <i>Breeding constraint: Hollow-bearing tree (Living or dead trees with hollows greater than 20 cm diameter and greater than 4m above the ground).</i>	Habitat absent, no suitable tree hollows
Regent Honeyeater	Species/Ecosystem	* <i>Breeding constraint: Other (As per mapped areas; contact OEH).</i>	Not a mapped important area
<b>Mammals</b>			
Brush-tailed Phascogale	Species	<i>Hollow-bearing trees.</i>	Habitat absent, no suitable tree hollows
Common Planigale	Species	<i>None.</i>	-
Koala	Species/Ecosystem	* <i>Breeding constraint: Other (Areas identified via survey as important habitat).</i>	Habitat absent, no eucalypts are present in the proposed MEA Extensions and only one was present

Common Name	Credit Class	Habitat Constraints identified in the TBDC (DPIE 2021a)	Assessment Prior to the Surveys
			in the proposed water treatment facility location, identified as Blakely's Red Gum ( <i>Eucalyptus blakelyi</i> ).
Eastern Pygmy-possum	Species	<i>None.</i>	-
Greater Glider	Species	<i>Hollow-bearing trees.</i>	Habitat absent, no suitable tree hollows
Squirrel Glider	Species	<i>None</i>	-
Brush-tailed Rock-wallaby	Species	<i>Other (Land within 1 km of rocky escarpments, gorges, steep slopes, boulder piles, rock outcrops or clifflines).</i>	Habitat absent
Grey-headed Flying-fox	Species/Ecosystem	<i>* Breeding constraint: Other (Breeding camps).</i>	Breeding habitat absent
Large Bentwing-bat	Species/Ecosystem	<i>* Breeding constraint: Caves (Cave, tunnel, mine, culvert or other structure known or suspected to be used for breeding).</i>	Habitat absent
Large-eared Pied Bat	Species	<i>Cliffs (Within two kilometres of rocky areas containing caves, overhangs, escarpments, outcrops, or crevices, or within two kilometres of old mines or tunnels).</i>	Habitat absent
Southern Myotis	Species	<i>Hollow-bearing trees (Within 200 m of riparian zone). Other (Bridges, caves or artificial structures within 200 m of riparian zone).</i>	Habitat absent

Shaded species are species that have habitat constraints as identified in the TBDC (DPIE 2021a) such that these species are not likely to occur and are therefore excluded from further assessment.

\* Habitat Constraints not in the BAM-C.

## 4.2.2.3 Degraded Habitat

A candidate species credit species is considered unlikely to occur on the Biodiversity Assessment Development Footprint (or specific vegetation zones) if after carrying out a field assessment of the habitat constraints or microhabitats on the Biodiversity Assessment Development Footprint, the assessor determines that the habitat is substantially degraded to the point that the species is unlikely to utilise the Biodiversity Assessment Development Footprint (or specific vegetation zones) (DPIE 2020) (Table 10).

**Table 10**  
**Species Credit Species Degraded Habitat**

Scientific Name	Common Name	Conservation Status <sup>1</sup>		Class of Credit <sup>2</sup>	PCT Association	Microhabitat is likely to be Substantially Degraded?
		BC Act	EPBC Act			
<b>Flora</b>						
<i>Acacia pendula</i> – endangered population	<i>Acacia pendula</i> population in the Hunter catchment	E	-	S	PCT 1606 and 1692	No
<i>Cryptostylis hunteriana</i>	Leafless Tongue Orchid	V	V	S	PCT 1606	Yes, the PCT 1606 present is derived native grassland. This species grows in eucalypt woodland.
<i>Cymbidium canaliculatum</i> – endangered population	<i>Cymbidium canaliculatum</i> in the Hunter Catchment	E	-	S	PCT 1606 and 1692	No
<i>Cynanchum elegans</i>	White-flowered Wax Plant	E	E	S	PCT 1606	Yes, the PCT 1606 present is derived native grassland. This species grows in woodland and scrub.
<i>Diuris tricolor</i>	Pine Donkey Orchid	V/EP	-	S	PCT 1606	Refer Section 4.3.4
<i>Eucalyptus glaucina</i>	Slaty Red Gum	V	V	S	PCT 1692	No
<i>Ozothamnus tessellatus</i>	-	V	V	S	PCT 1606	Yes, the PCT 1606 present is derived native grassland. This species grows in eucalypt woodland.
<i>Pomaderris bodalla</i>	Bodalla Pomaderris	V	-	S	PCT 1606	Yes, the PCT 1606 present is derived native grassland. All known populations of this species grow in open forest or woodland.
<i>Pomaderris queenslandica</i>	Scant Pomaderris	E	-	S	PCT 1606	Yes, the PCT 1606 present is derived native grassland. This species is found in moist eucalypt forest or sheltered woodlands.
<i>Thesium australe</i>	Austral Toadflax	V	V	S	PCT 1606	No

Scientific Name	Common Name	Conservation Status <sup>1</sup>		Class of Credit <sup>2</sup>	PCT Association	Microhabitat is likely to be Substantially Degraded?
		BC Act	EPBC Act			
<b>Amphibians</b>						
<i>Litoria aurea</i>	Green and Golden Bell Frog	E	V	S	PCT 1606 and 1692	The dams present are not suitable for the Green and Golden Bell Frog.
<b>Reptiles</b>						
<i>Hoplocephalus bitorquatus</i>	Pale-headed Snake	V	-	S	PCT 1606 and 1692	No trees with hollows.
<i>Delma impar</i>	Striped Legless Lizard	V	V	S	PCT 1606 and 1692	No for the Proposed MEA Extensions but PCT 1692 at the proposed water treatment facility location is not considered potential habitat for this species as it is highly disturbed and provides no suitable shelter (rocks, dumped material or dried cow manure).
<b>Birds</b>						
<i>Burhinus grallarius</i>	Bush Stone-curlew	E	-	S	PCT 1606 and 1692	No
<b>Mammals</b>						
<i>Planigale maculata</i>	Common Planigale	V	-	S	PCT 1606 and 1692	No
<i>Cercartetus nanus</i>	Eastern Pygmy-possum	V	-	S	PCT 1606 and 1692	No trees with hollows, paddock trees are not important habitat for this species.
<i>Petaurus norfolcensis</i>	Squirrel Glider	V	-	S	PCT 1606	No

1 Threatened species status under the BC Act and/or EPBC Act (current as at August 2021). V = Vulnerable; E = Endangered; CE = Critically Endangered; EP = Endangered Population.

2 Biodiversity credit class under the TBDC (DPIE 2021a) (current as at August 2021). E = Ecosystem; S = Species.

### 4.2.3 Addition of Species Credit Species

No Species Credit Species were added to the BAM-C. It is noted that DPIE has previously queried the additional consideration of *Prasophyllum petilum* and *Pterostylis chaetophora* for the Project. However, the analysis by Dr Stephen Bell (Bell 2020) (Attachment C) concludes these two species are not relevant to the locality and/or vegetation types in the Biodiversity Assessment Development Footprint associated with the Modification.

#### 4.2.3.1 Review of Databases

The following databases were reviewed for any nearby potentially occurring threatened species records (including species credit species):

- *Birdlife Australia Atlas Database* (Birdlife Australia 2018);
- *BioNet Atlas* (DPIE 2021c);
- *Protected Matters Search Tool* (DEE 2018b); and

- *Atlas of Living Australia* (Atlas of Living Australia [ALA] 2021).

Attachment D provides a summary of the threatened species records in the locality from survey records or database records (threatened species shaded in Attachment D are species with records in the Project study area). Threatened species records are shown on Figures 8 to 12.

### 4.3 SPECIES CREDIT SPECIES – ASSESSMENT METHOD

#### 4.3.1 Species Assumed to be Present

No Species Credit Species are assumed to be present.

#### 4.3.2 Species Important Habitat Mapping

No DPIE Important Habitat Mapping is relevant to the Biodiversity Assessment Development Footprint.

#### 4.3.3 Species for Survey

Relevant species credit species were surveyed as per Section 4.4.

#### 4.3.4 Export Reports

Dr Stephen Bell of Eastcoast Flora Survey prepared an Expert Report on the Pine Donkey Orchid (*Diuris tricolor*) for the Project. The analysis by Dr Stephen Bell (Bell 2020) (Attachment C) concludes that the vegetation units defined by Hunter Eco (2019) as PCT 1691 derived native grassland and PCT 1604 provide the only potential habitat for *Diuris tricolor*. No PCT 1691 derived native grassland or PCT 1604 would be cleared for the Modification.

It is noted that the TBDC (DPIE 2021a) indicates that the species may be associated with PCT 1606, however it also indicates that the endangered population in the Muswellbrook Local Government Area (i.e. the listing relevant to the Biodiversity Assessment Development Footprint) is not associated with PCT 1606.


On the basis of the analysis by Dr Stephen Bell (Bell 2020) (Attachment C), the information in the TBDC (DPIE 2021a) and site observations, the Pine Donkey Orchid is not considered to be a relevant species for survey for the Modification.

### 4.4 SPECIES CREDIT SPECIES – SURVEY

#### **Threatened Flora**

Dr Colin Driscoll (Hunter Eco) (2019) undertook targeted surveys for flora species credit species for the Project. Supplementary flora surveys for the Modification were undertaken by Dr Colin Driscoll on 28 June 2021 (Table 11). Surveys were conducted according to the requirements in the Department of Planning, Industry & Environment (DPIE) (2020) *Surveying Threatened Plants and Their Habitats: NSW Survey Guide for the Biodiversity Assessment Method*.

Based on Table 10, only *Acacia pendula*, *Thesium australe*, *Eucalyptus glaucina* and *Cymbidium canaliculatum* required surveying. These were surveyed as follows.



- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
- Threatened Species**
- *Diuris tricolor*
  - *Eucalyptus glaucina*
  - *Eucalyptus nicholii*
- Endangered Populations**
- Cymbidium canaliculatum* population in the Hunter Catchment
  - *Acacia pendula* population in the Hunter Catchment #
  - *Diuris tricolor* Fitzg., the Pine Donkey Orchid, in the Muswellbrook Local Government Area
  - *Eucalyptus camaldulensis* in the Hunter Catchment

# Note *Acacia pendula* is also listed as a threatened ecological community under the BC Act (Hunter Valley Weeping Myall Woodland) and the EPBC Act (Weeping Myall [*Acacia pendula*] Woodland).

Note: OEH (2019) also contains records for the Cumberland Ecology and Hunter Eco records of *Cymbidium canaliculatum* population in the Hunter Catchment, but the co-ordinates are less precise and are therefore not shown on this figure.


Reference:  
 8. Cumberland Ecology (2015)  
 13. DPIE (2020)  
 14. Hunter Eco (2019)  
 21. Hunter Eco (2020)  
 Note: There are no references 1 - 7, 9 - 12 and 15 - 20 on this figure.

Source: NSW Spatial Services (2021); MSEC (2019)  
 Orthophoto Mosaic: 2020, 2019

**MALABAR**  
**MAXWELL UNDERGROUND MINE PROJECT**  
**Threatened Flora Species/  
 Endangered Populations**

**Figure 8**






- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
  - Threatened Species
  - Pink-tailed Legless Lizard
  - Striped Legless Lizard

Reference:  
 13. DPIE (2020)  
 15. Future Ecology (2019)  
 Note: There are no references 1 - 12 and 14 on this figure.

Source: NSW Spatial Services (2021); MSEC (2019)  
 Orthophoto Mosaic: 2020, 2019

**MALABAR**  
 MAXWELL UNDERGROUND MINE PROJECT  
 Threatened Reptiles and Amphibians

**Figure 9**




- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extend of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
- Threatened Species**
- Black Falcon
  - Square-tailed Kite
  - White-bellied Sea-Eagle
  - Spotted Harrier
  - Little Eagle
  - Little Lorikeet
  - Swift Parrot
  - Barking Owl
  - White-throated Needletail
  - Brown Treecreeper (eastern subspecies)
  - Speckled Warbler
  - Black-chinned Honeyeater (eastern subspecies)
  - Painted Honeyeater
  - Hooded Robin (south-eastern form)
  - Flame Robin
  - Scarlet Robin
  - Grey-crowned Babbler (eastern subspecies)
  - Varied Sittella
  - Dusky Woodswallow
  - Diamond Firetail

- Reference:**
- |                               |                                     |
|-------------------------------|-------------------------------------|
| 1. Ecotone (2000)             | 12. Birdlife Australia (2018)       |
| 2. Umwelt (2005)              | 13. DPIE (2020)                     |
| 6. Cumberland Ecology (2009a) | 14. Hunter Eco (pers. comm. (2019)) |
| 7. Cumberland Ecology (2010)  | 15. Future Ecology (2019)           |
| 8. Cumberland Ecology (2015)  | 17. Eco Logical Australia (2015)    |


Note: There are no references 3 - 5, 9 - 11 and 16 on this figure.

Source: NSW Spatial Services (2021); MSEC (2019)  
Orthophoto Mosaic: 2020, 2019

**MALABAR**  
**MAXWELL UNDERGROUND MINE PROJECT**  
**Threatened Birds**



**Figure 10**



- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
- Threatened Species**
- ▲ Spotted-tailed Quoll
  - ▲ Koala
  - ▲ Squirrel Glider

**Reference:**


- 5. Umwelt (2007b)
- 6. Cumberland Ecology (2009a)
- 13. DPIE (2020)
- 15. Future Ecology (2019)
- 16. Eco Logical Australia (2014)
- 18. Eco Logical Australia (2016a)
- 19. Eco Logical Australia (2016b)

*Note: There are no references 1 - 4, 8 - 12, 14 and 17 on this figure.*

**Source:** NSW Spatial Services (2021); MSEC (2019)  
 Orthophoto Mosaic: 2020, 2019



**Figure 11**



- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
- Threatened Species**
- ▽ Grey-headed Flying-fox
  - ▽ Yellow-bellied Shearwater-bat
  - ▽ Northern Free-tailed Bat
  - ▽ Eastern Coastal Free-tailed Bat
  - ▽ Large Bent-winged Bat
  - ▽ Corben's Long-eared Bat
  - ▽ Large-eared Pied Bat
  - ▽ Little Bent-winged Bat
  - ▽ Eastern False Pipistrelle
  - ▽ Southern Myotis
  - ▽ Greater Broad-nosed Bat
  - ▽ Eastern Cave Bat

**Reference:**

1. Ecotone (2000)
4. Umwelt (2007a)
6. Cumberland Ecology (2009a)
8. Cumberland Ecology (2015)
11. Niche (2012)
13. DPIE (2020)
15. Future Ecology (2019)
16. Eco Logical Australia (2014)
17. Eco Logical Australia (2015)
18. Eco Logical Australia (2016a)
19. Eco Logical Australia (2016b)
20. Eco Logical Australia (2017)

*Note: There are no references 2, 3, 5, 7, 9, 10, 12 and 14 on this figure. (Some records by Niche (2012) were possible/probable records).*

**Source:** © NSW Spatial Services (2021); MSEC (2019) Orthophoto Mosaic: 2019, 2020

**MAXWELL UNDERGROUND MINE PROJECT**  
**Threatened Bats**



**Figure 12**

*Acacia pendula*

In June 2021, 10 m wide parallel transects within PCT 1606 (in derived native grassland form) were conducted in the Proposed MEA Extensions (Figure 7a). This exceeds the 40 m spacing requirement for open vegetation (DPIE 2020). Given the small area of PCT 1692 in the Proposed MEA Extensions all *Acacia* trees were inspected; only *Acacia salicina* was present. Given the proposed water treatment facility location was small (0.1 ha), the entire occurrence of PCT 1692 was inspected and every *Acacia* sp. tree present was documented with *Acacia pendula* absent.

*Cymbidium canaliculatum*

In June 2021, 10 m wide parallel transects within PCT 1606 (in derived native grassland form) were conducted in the Proposed MEA Extensions (Figure 7a). A 10 m wide transect survey was appropriate because, as noted in Section 3.3, the grassland was open habitat. Given the small area of PCT 1692, all trees in the patches of PCT 1692 were inspected. No eucalypts are present in the proposed MEA Extensions and only one was present in the proposed water treatment facility location, identified as Blakely's Red Gum (*Eucalyptus blakelyi*).

*Thesium australe*

*Thesium australe* is a low sprawling plant generally associated with Kangaroo Grass (*Themeda triandra*). In June 2021, 10 m wide parallel transects within PCT 1606 (in derived native grassland form) were conducted in the proposed MEA Extensions targeting the parasitic host *Themeda triandra* (Figure 7a). A 10 m wide transect survey was appropriate because, as noted in Section 3.3, the grassland was open habitat. Although this timing was outside of the survey months for *Thesium australe*, this timing was suitable to detect the parasitic host *Themeda triandra*, if present. This approach was consistent with the advice from DPIE on the Project (dated August 2020).

*Eucalyptus glaucina*

Given the small area of PCT 1692, all trees in the patches of PCT 1692 were inspected. One Blakely's Red Gum (*Eucalyptus blakelyi*) was present in the proposed water treatment facility location.

**Threatened Fauna**

Future Ecology (2019) (Attachment A) undertook targeted surveys for candidate fauna species credit species for the Project. Months in which targeted surveys were undertaken are shaded in Table 11.

*Bush Stone-curlew*

No Bush Stone-curlew are known to occur in the locality (Figure 10). The potential habitat for the Bush Stone-curlew in the Biodiversity Assessment Development Footprint is marginal, and there is more extensive area of similar potential habitat in the surrounding locality that is not known to be used by the species.

Numerous nocturnal bird surveys incorporating call-playback and spotlighting were carried out at various sites at various times of the year for the Bush Stone-curlew in 2018 (Future Ecology 2019). Survey sites included locations within 1 km of the MEA Extensions and approximately 2 km from the proposed water treatment facility location.

No Bush Stone-curlew were found, despite the extensive targeted surveys by Future Ecology (2019).

### Common Planigale

No Common Planigale are known to occur in the locality (Figure 11). The potential habitat for the Common Planigale in the Biodiversity Assessment Development Footprint is marginal, and there is more extensive area of similar potential habitat in the surrounding locality that is not known to be used by the species.

A range of survey methods were used by Future Ecology (2019) (Attachment A) to target the species in the Project study area:

- a total of 5,004 hair tube trap nights and 501 camera trap nights were completed over the study area from January to June and November to December 2018;
- a total of 16 cage and 280 arboreal Elliott B trap nights were carried out in January and November 2018;
- a total of 1,855 nest box nights were carried out from January to June 2018;
- numerous nocturnal spotlighting surveys were carried out throughout the study area;
- predator scats were collected and analysed whenever detected;
- a total of 72 trap nights of pitfall trapping took place; and
- a total of 400 Elliott A terrestrial trap nights were carried out.

No Common Planigale were found, despite the extensive targeted surveys by Future Ecology (2019) (Attachment A).

### Squirrel Glider

The Squirrel Glider has been recorded in the locations shown on Figure 11. Future Ecology (2019) (Attachment A) recorded the species during targeted surveys involving a total of 5,004 hair tube trap nights and 501 camera trap nights in the Project study area from January to June and November to December 2018. A total of 16 cage and 280 arboreal Elliott B trap nights were carried out in January and November 2018. A total of 1,855 nest box nights were carried out from January to June 2018.

Numerous nocturnal spotlighting surveys were carried out throughout the study area. Predator scats were collected and analysed whenever detected.

### Striped Legless Lizard

The Striped Legless Lizard has been recorded in the locations shown on Figure 9. Future Ecology (2019) (Attachment A) recorded the species during targeted surveys involving several methods, namely pitfall trapping, artificial shelter sites and active searches for both living specimens and sloughs (shed skins).

Pitfall trapping was undertaken over 24 trap nights, using six traps for four consecutive nights, at three sites in November 2018. In addition, four grids of 50 artificial shelter sites (total of 200) were installed in November 2018 and checked several times through December 2018.

Diurnal and nocturnal active searches were carried out in November/December 2018. Further, diurnal habitat searches were also undertaken in January 2018, and included careful observations under cow pats (dung) in open grassy areas with a good cover of native grasses and herbs.

**Table 11**  
**Species Credit Species Requiring Survey and Timing**

Scientific Name	Common Name	Conservation Status <sup>1</sup>		Class of Credit <sup>2</sup>	Survey Months for Each Species												
		BC Act	EPBC Act		Jan	Feb	March	April	May	June	July	August	Sep	Oct	Nov	Dec	
<b>Flora</b>																	
<i>Acacia pendula</i> – endangered population	<i>Acacia pendula</i> population in the Hunter catchment	E	-	S	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
<i>Cymbidium canaliculatum</i> – endangered population	<i>Cymbidium canaliculatum</i> in the Hunter Catchment	E	-	S	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	-	Yes	Yes	Yes	Yes
<i>Eucalyptus glaucina</i>	Slaty Red Gum	V	-	S	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
<i>Thesium australe</i> <sup>4</sup>	Austral Toadflax	V	V	S	Yes	Yes	-	-	-	*Note	-	-	-	-	-	Yes	Yes
<b>Fauna</b>																	
<i>Delma impar</i>	Striped Legless Lizard	V	V	S	-	-	-	-	-	-	-	-	-	Yes	Yes	Yes	Yes
<i>Burhinus grallarius</i>	Bush Stone-curlew	E	-	S	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
<i>Planigale maculata</i>	Common Planigale	V	-	S	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
<i>Petaurus norfolcensis</i>	Squirrel Glider	V	-	S	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes

Note: Months in which surveys for the species are to be conducted in accordance with the TBDC (DPIE 2021a) are denoted with 'Yes'. The shaded month is the month in which targeted surveys were undertaken for the relevant species.

\* Although, this timing was outside of the survey month for *Thesium australe*, this timing is suitable to detect the parasitic host *Themeda triandra*.

1 Threatened species status under the BC Act and/or EPBC Act (current as at August 2021). V = Vulnerable; E = Endangered; CE = Critically Endangered.

2 Biodiversity credit class under the TBDC (DPIE 2021a) (current as at August 2021). E = Ecosystem; S = Species.

4 A 10 m transect survey targeting the Austral Toadflax host species Kangaroo Grass was conducted in open derived native grassland of PCT 1606 in the MEA and none was found negating the need for later targeted surveys for Austral Toadflax.

#### 4.4.1 Survey Results

##### ***Threatened Flora***

No threatened flora species were recorded. No eucalypts are present in the proposed MEA Extensions and only one was present in the proposed water treatment facility location, identified as Blakely's Red Gum (*Eucalyptus blakelyi*). The parasitic host of *Thesium australe* (i.e. *Themeda triandra*) was not found.

##### ***Threatened Fauna***

The following species credit fauna species were relevant to the Modification (Figures 13 to 16):

- Pink-tailed Legless Lizard (listed as vulnerable under the BC Act and EPBC Act);
- Striped Legless Lizard (listed as vulnerable under the BC Act and EPBC Act);
- Squirrel Glider (listed as vulnerable under the BC Act); and
- Southern Myotis (listed as vulnerable under the BC Act).

No habitat is present in the Biodiversity Assessment Development Footprints for the Pink-tailed Legless Lizard and Southern Myotis (Figures 13 and 16). However, the Biodiversity Assessment Development Footprints contain potential habitat for the Squirrel Glider and Striped Legless Lizard. The Squirrel Glider has a biodiversity risk weighting of '2', and the Striped Legless Lizard has a biodiversity risk weighting of '1.5'. Neither of these species are Potential SAI Entities.

##### *Striped Legless Lizard*

One specimen of Striped Legless Lizard was recorded by Future Ecology (2019) (Attachment A) approximately 1 km south east of the MEA (Figure 14). A total of 26 observations of Striped Legless Lizard were recorded within the Project study area, 16 of which were live specimens and 10 were sloughs. Most observations were scattered throughout the Maxwell Underground and along or near the proposed transport and services corridor between Maxwell Infrastructure and Maxwell Underground. The Striped Legless Lizard was recorded under rocks, dumped material and dried cow manure.


As described in Table 10, PCT 1692 at the proposed water treatment facility location is not considered potential habitat for this species as it is highly disturbed and provides no suitable shelter (rocks, dumped material or dried cow manure).

##### *Squirrel Glider*

The Squirrel Glider was observed by Future Ecology (2019) (Attachment A) twice just outside of the Maxwell Underground to the east, twice in the north of the Maxwell Underground and twice (of one individual) just south of the MEA Extensions (Figure 15). It is not clear how many other Squirrel Gliders have been recorded by other groups in the past, however this number is expected to be relatively numerous (Umwelt 2006, 2007; Cumberland Ecology 2009a, 2010; DPIE 2021b). There have also been several observations of this species outside of the Project study area by other groups.

Similar to the Striped Legless Lizard, PCT 1692 at the proposed water treatment facility location is not considered potential habitat for the Squirrel Glider.






- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
- SPECIES POLYGON MAPPING**
- Dry Sclerophyll Forests (Shrub/grass sub-formation)
  - 2. White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)
  - 2a. White Box - Ironbark - Red Gum Shrubby Forest - Derived Native Grassland (PCT1606)
- Other**
- Rocky Area
  - Rocky Area 50 m Zone

Reference:  
 15. Future Ecology (2019)  
 Note: There are no references 1 - 14 on this figure.

Source: NSW Spatial Services (2021); MSEC (2019)  
 Orthophoto Mosaic: 2020, 2019

**MALABAR**  
 MAXWELL UNDERGROUND MINE PROJECT  
 Pink-tailed Legless Lizard  
 Species Polygon

**Figure 13**




- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
  - Striped Legless Lizard
- SPECIES POLYGON MAPPING**
- Dry Sclerophyll Forests (Shrub/grass sub-formation)
  - 2. White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)
  - 2a. White Box - Ironbark - Red Gum Shrubby Forest - Derived Native Grassland (PCT1606)
  - Dry Sclerophyll Forests (Shrubby sub-formation)
  - 3. Slaty Box Shrubby Woodland (PCT1655)
  - 3a. Slaty Box Shrubby Woodland - Derived Native Grassland (PCT1655)
  - Grassy Woodlands
  - 6. Bull Oak Grassy Woodland (PCT1692)
  - 7. Yellow Box - Apple Grassy Woodland (PCT1693)
  - 7a. Yellow Box - Apple Grassy Woodland - Derived Native Grassland (PCT1693)
  - 9. Ironbark - Grey Box Grassy Woodland (PCT1691)
  - 9a. Ironbark - Grey Box Grassy Woodland - Derived Native Grassland (PCT1691)
  - 11. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)

*Reference:*  
 13. DPIE (2020)  
 15. Future Ecology (2019)  
*Note: There are no references 1 - 12 and 14 on this figure.*

*Source: NSW Spatial Services (2021); MSEC (2019) Orthophoto Mosaic, 2020, 2019*

  
**MAXWELL UNDERGROUND MINE PROJECT**  
**Striped Legless Lizard**  
**Species Polygon**

**Figure 14**



- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
  - ▲ Squirrel Glider
- SPECIES POLYGON MAPPING**
- Dry Sclerophyll Forests (Shrub/grass sub-formation)
  - 2. White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)
  - Dry Sclerophyll Forests (Shrubby sub-formation)
  - 3. Slaty Box Shrubby Woodland (PCT1655)
  - Forested Wetlands
  - 5. Hunter Lowland Red Gum Forest (PCT1598)
  - Grassy Woodlands
  - 8. Fuzzy Box Woodland (PCT201)
  - 11. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)
- Other**
- Paddock Trees that may provide Potential Habitat

**Reference:**


- 5. Umwelt (2007b)
- 6. Cumberland Ecology (2009a)
- 13. DPIE (2020)
- 15. Future Ecology (2019)
- 16. Eco Logical Australia (2014)
- 18. Eco Logical Australia (2016a)
- 19. Eco Logical Australia (2016b)

*Note: There are no references 1 - 4, 7 - 12, 14 and 17 on this figure.*

**Source:** NSW Spatial Services (2021); MSEC (2019)  
 Orthophoto Mosaic: 2020, 2019

**MALABAR**  
**MAXWELL UNDERGROUND MINE PROJECT**  
**Squirrel Glider**  
**Species Polygon**

**Figure 15**



**LEGEND**

- Exploration Licence Boundary
- Mining and Coal Lease Boundary
- Mining Lease Application Boundary
- Approved Surface Development Area
- Approved Extent of Conventional Subsidence from Underground Mining
- Biodiversity Assessment Development Footprint
- Existing Conservation/Offset Area
- ▼ Southern Myotis

**SPECIES POLYGON MAPPING**

**Grassy Woodlands**

- 6. Bull Oak Grassy Woodland (PCT1692)
- 9. Ironbark - Grey Box Grassy Woodland (PCT1691)
- 11. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)

**Other**

- Waterbody/Dam
- Waterbody/Dam 200 m Buffer

**Reference:**

1. Ecotone (2000)
6. Cumberland Ecology (2009a)
8. Cumberland Ecology (2015)
13. DPIE (2020)
20. Eco Logical Australia (2017)

*Note: There are no references 2 - 5, 7, 9 - 12, 14 - 19 on this figure.*

**Source:** NSW Spatial Services (2021); MSEC (2019) Orthophoto Mosaic: 2020, 2019

**MALABAR**  
 MAXWELL UNDERGROUND MINE PROJECT  
 Southern Myotis  
 Species Polygon

**Figure 16**

4.4.2 Species Polygons

Species Polygons for the Striped Legless Lizard and Squirrel Glider were prepared by Future Ecology (2019) (Attachment A) as shown on Figures 14 and 15. The area of habitat in the Biodiversity Assessment Development Footprint is quantified in Table 12. The VI Score for each vegetation zone associated with the Species Polygons for the Striped Legless Lizard and Squirrel Glider are listed in Table 12.

**Table 12  
Species Credit Species Presence**

Generic Name	PCT	VI Score~	Area of Habitat (ha)	
			Striped Legless Lizard	Squirrel Glider
<b>Proposed MEA Extensions</b>				
2a. White Box - Ironbark - Red Gum Shrubby Forest PCT 1606 (DNG)	1606	31.4	3.8	1.1
6. Bull Oak Grassy Woodland PCT 1692	1692	48.5	0.5	0
<b>Proposed Water Treatment Facility Location</b>				
6. Bull Oak Grassy Woodland PCT 1692	1692	38.2	0	0
<b>Total</b>			<b>4.3</b>	<b>1.1</b>

~ Vegetation Integrity Score – refer to Table 2.

## 5 PRESCRIBED BIODIVERSITY IMPACT ENTITIES

Prescribed biodiversity impact entities are discussed below.

### 5.1 KARST, CAVES, CREVICES, CLIFFS, ROCKS AND OTHER GEOLOGICAL FEATURES

There are no karst, caves, crevices or cliffs that provide habitat for threatened species in the Biodiversity Assessment Development Footprint. Rocky areas providing potential habitat for the Pink-tailed Legless Lizard and Striped Legless Lizard are present in the Project area as mapped by Future Ecology (2019) (Attachment A). Rocky areas are shown on Figure 5. Individual rocks were mostly less than 50 cm long and wide and deep.

### 5.2 HUMAN MADE STRUCTURES AND NON-NATIVE VEGETATION

There are no human made structures that provide habitat for threatened species in the Biodiversity Assessment Development Footprint. There are no areas of non-native vegetation that provide habitat for threatened species in the Biodiversity Assessment Development Footprint.

### 5.3 HABITAT CORRIDORS OR LINKAGES

No defined habitat corridors or linkages would be impacted. The revised site access road alignment follows an existing haul road and does not contain native vegetation. The original native vegetation in the extension to the portal entry and extension to the clean water diversion bank consists of derived native grassland or re-growth Buloke woodland. The majority of the water treatment facility location has been disturbed by past mining activities with small fragmented groups of Buloke trees.

It is recognised that the derived native grassland may provide habitat linkages for some species (e.g. Striped Legless Lizard) (Section 7.6.3).

### 5.4 WATERBODIES OR HYDROLOGICAL PROCESSES THAT SUSTAIN THREATENED SPECIES

No water bodies or hydrological processes that are known to sustain threatened species would be impacted by the Modification.

### 5.5 THREATENED SPECIES AT RISK FROM VEHICLE STRIKE

The Modification would not change the risk on threatened species from vehicle strike. The revised site access road alignment follows an existing haul road and does not contain native vegetation.

## 6 AVOID AND MINIMISE IMPACTS

The location of the Project has been selected based upon the presence of coal seams able to be economically mined within Malabar's existing tenements and the extensive geological and geotechnical data available within the target area in EL 5460. Malabar is committed to developing the Project solely as an underground mining operation. Underground mining methods significantly reduce environmental impacts, including vegetation and habitat disturbance, in comparison to open cut mining methods. In addition to the use of underground mining methods, Project elements have been located and designed to avoid or minimise impacts to vegetation and habitat disturbance as described in the *Maxwell Project – Biodiversity Development Assessment Report* (Hunter Eco, 2019).

The Modification would primarily comprise the following components:

- repositioning of the underground portal;
- realignment of a small section of the proposed access road at the Maxwell Infrastructure site to utilise an existing haul road;
- repositioning of approved clean water diversions for the MEA; and
- repositioning of the water treatment facility to the Maxwell Infrastructure.

The reconfiguration of the MEA, specifically the extension to the portal entry outside the approved surface development area, would allow for safer commencement of the underground portal and would optimise the timing and costs associated with underground mine development. The extension to the portal entry involves some additional disturbance of derived native grassland (Box-Gum TEC), but avoids woodland/forest vegetation communities (Figure 7a).

The extension to the clean water diversion bank involves some additional disturbance of derived native grassland (Box-Gum TEC), and also largely avoids woodland/forest vegetation communities to the south, except for a partial impact (0.5 ha) on a patch of re-growth Buloke woodland (mapped as PCT 1692) (Figure 7a).

The clearance of Box-Gum TEC for the Modification is unavoidable, as it covers a large area surrounding the Project surface facilities such that the Modification could not be moved in a way that would avoid or reduce the amount lost.

The Modification would also minimise the impacts to the existing rehabilitation at the Maxwell Infrastructure through the realignment of the site access road along an existing internal haul road, without native vegetation.

Repositioning of the approved clean water diversion and water treatment facility provide material environment benefits through improved water management outcomes (e.g. reducing environmental risk associated with pumping brine from the MEA to the Maxwell Infrastructure).

The proposed location of the water treatment facility is on previously disturbed land at the Maxwell Infrastructure. Malabar considered moving the proposed location of the water treatment facility to avoid PCT 1692 (0.1 ha), however, the water treatment facility was located close to existing electrical infrastructure that is needed for the facility. The majority of the water treatment facility location has been disturbed by past mining activities with small fragmented groups of Buloke trees.

There are no alternative modes or technologies relevant to the Modification.

## 7 EVALUATION OF POTENTIAL IMPACTS

### 7.1 DIRECT IMPACTS ON NATIVE VEGETATION AND HABITAT

#### 7.1.1 Clearance of Native Vegetation

After applying the measures to avoid and/or minimise impacts on biodiversity values (Section 6), the Modification would result in the clearance of approximately 4.4 ha of native vegetation within the Biodiversity Assessment Development Footprint (Table 13). The total amount of native vegetation to be disturbed for the Modification is 2.7 % of the native vegetation approved to be disturbed by the Project (i.e. 161.6 ha).

**Table 13**  
**Native Vegetation Clearance Summary**

PCT	Generic Name	Total Area (ha)	Change in VI Score
1606	2a. White Box - Ironbark - Red Gum Shrubby Forest (DNG) <sup>1</sup>	3.8	-31.4
1692	6. Bull Oak Grassy Woodland	0.6	-48.5 and -38.2
<b>Total Native Vegetation</b>		4.4	

<sup>1</sup> Listed as Box-Gum TEC under the BC Act and EPBC Act.

Threatened species associated with the Biodiversity Assessment Development Footprint are discussed in Section 4.

Surface disturbance areas associated with the Biodiversity Assessment Development Footprint would be rehabilitated and revegetated when the surface facilities are no longer required or at the end of the mine life where no further ongoing beneficial use is identified (Figure 17). The final land use of the surface disturbance areas would include areas for agricultural land use and native vegetation, and would be developed in consultation with relevant stakeholders.

#### 7.1.2 Cumulative Impacts


As described in the *Maxwell Project – Biodiversity Development Assessment Report* (Hunter Eco 2019), the surface development for the Project would involve direct disturbance of approximately 25.6 ha of fragmented native woodland/forest and 136 ha of derived native grassland, including areas that would be impacted from ponding within the Maxwell Underground area. The total amount of native vegetation to be disturbed for the Project is approximately 161.6 ha.

The Project was assessed under the BAM which requires biodiversity offsets resulting in no net loss in biodiversity. The Modification is also assessed under the BAM and would also require a biodiversity offset resulting in no net loss in biodiversity. The biodiversity offset is required to be a greater area of land, multiple times the size of the Biodiversity Assessment Development Footprint, which will be conserved and managed to achieve a gain in biodiversity values.

### 7.2 INDIRECT IMPACTS ON NATIVE VEGETATION AND HABITAT

Native vegetation adjacent to the Biodiversity Assessment Development Footprint is shown on Figures 7a and 7b and threatened fauna records adjacent to the Biodiversity Assessment Development Footprint are shown on Figures 8 to 12. Potential indirect impacts on habitat and vegetation listed in the BAM (DPIE 2020) are assessed below in relation to the Modification. The Modification is unlikely to result in an increase or adverse indirect impacts on native vegetation and habitat for threatened species.






- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
- Secondary Domains**
- A Biodiversity Offset
  - B Water Management
  - C Pasture
  - D Woodland
  - E Buffer Land
  - F Remediated Underground Mining Area

Source: MSEC (2019); NSW Spatial Services (2021)  
 Orthophoto Mosaic: 2020, 2019

**MALABAR**  
 MAXWELL UNDERGROUND MINE PROJECT  
 Rehabilitation Domains



**Figure 17**

The Modification would not result in any changes to subsidence impacts described in the *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b).

### **7.2.1 Inadvertent Impacts on Adjacent Habitat or Vegetation**

As described in the *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b), a vegetation clearance protocol would be adopted (Section 8) to minimise the risk of inadvertent impacts on adjacent habitat or native vegetation in the short-term during construction or operation of the Project (and Modification).

### **7.2.2 Impacts on Adjacent Habitat or Vegetation from a Change in Land Use Pattern (Increased Human Activity)**

No adverse impacts are likely to result on habitat and vegetation adjacent to the Biodiversity Assessment Development Footprint as the Modification would not increase human activity in the locality.

### **7.2.3 Reduced Viability of Adjacent Habitat Due to Edge Effects**

There would be no material edge effects from the revised site access road alignment or water treatment facility location. In regard to the proposed MEA Extensions, there would be a slight increase in the native vegetation that would be adjacent to the Modified Project (compared to the approved surface development area). However, the viability of the adjacent habitat is unlikely to be reduced due to edge effects because the adjacent habitat is either derived native grassland or regrowth Buloke trees.

### **7.2.4 Reduced Viability of Adjacent Habitat Due to Noise, Dust or Light Spill**

The *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b) describes that the Project is unlikely to reduce the viability of any adjacent habitat due to noise, dust or light spill during construction or operation. From field observations, there is no evidence of dust from currently approved operations having impacted surrounding vegetation. It is unlikely that any flora species would be adversely impacted either directly or indirectly by any dust increase generated as a result of the Project. Lighting is used at the existing Maxwell Infrastructure and would be used for the Project. Night-lighting of the Project surface facilities would be kept to a practicable minimum.

The Modification would not increase night-lighting. The scale of dust and noise-generating activities would not materially increase compared to the activities assessed for the Project. For example, the site access road would be sealed during the first year of mining operations. Dust suppression would occur along the site access road, prior to it being sealed.

### **7.2.5 Transport of Weeds and Pathogens from the Site to Adjacent Vegetation**

The *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b) describes that weeds are relatively evenly distributed across the wider area which means that it is unlikely that there would be any dispersal of weeds from the Project that were not already present in the surrounds. The potential impacts from the Project to surrounding native vegetation associated with introduced flora is likely to be low. The Modification is unlikely to increase the risk of weeds or pathogens transporting from the site to adjacent vegetation.

### **7.2.6 Increased Risk of Fauna Starvation, Exposure and Loss of Shade or Shelter**

The *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b) describes that sufficient connectivity would remain around the Biodiversity Assessment Development Footprint such that no fauna are likely to become isolated as a result of the Project. The Modification would not increase the risk of fauna become isolated and subject to starvation, exposure and loss of shade or shelter.

### **7.2.7 Loss of Breeding Habitats**

The *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b) describes that the Project is not likely to indirectly impact fauna breeding habitat resources (such as trees with hollows and bush rock). Similarly, the Modification is unlikely to indirectly impact fauna breeding habitat resources.

### **7.2.8 Trampling of Threatened Flora Species**

The *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b) describes that no threatened flora species have been located in areas at risk of trampling during construction or operation of the Project. Similarly, no threatened flora species have been located in areas adjacent to the Biodiversity Assessment Development Footprint.

### **7.2.9 Inhibition of Nitrogen Fixation and Increased Soil Salinity**

The Project (and Modified Project) would not inhibit nitrogen fixation or increase soil salinity.

### **7.2.10 Fertiliser Drift**

The Project (and Modified Project) would not involve the use of fertiliser, except in small quantities to assist with revegetation works. Agricultural and other land management activities (which may involve use of fertilizer) would continue on Malabar-owned properties irrespective of the Project (and Modified Project).

### **7.2.11 Rubbish Dumping**

The Project (and Modified Project) would not involve rubbish dumping. Rubbish generated by the Project (and Modified Project) would be disposed of appropriately in designated areas.

### **7.2.12 Wood Collection**

Collection of wood from surrounding native vegetation (for fires or other activities) would not be permitted for the Project (and Modified Project). Agricultural and other land management activities would continue on Malabar-owned properties irrespective of the Project (and Modified Project).

### **7.2.13 Bush Rock Removal and Disturbance**

Removal or disturbance of bush rock from surrounding native vegetation would not be permitted for the Project (and Modified Project). Agricultural and other land management activities would continue on Malabar-owned properties irrespective of the Project (and Modified Project).

### 7.2.14 Increase in Predatory Species Populations

The Project (and Modified Project) is unlikely to increase predatory species populations (such as Cat and Red Fox). Agricultural activities would continue to occur both inside and outside the Project area (and Modified Project), including control of pest animal populations.

### 7.2.15 Increase in Pest Animal Populations

The *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b) describes that the Project is unlikely to increase pest animal populations. The Project (and Modified Project) would continue pest management activities (e.g. wild dogs and rabbits).

### 7.2.16 Increased Risk of Fire

The Project (and Modified Project) is unlikely to increase fire risk. Bushfire risk would continue to be managed in accordance with Malabar's existing Bushfire Management Procedure, which applies to Maxwell Infrastructure and Maxwell Underground. Bushfire management measures include the maintenance of fire breaks and access tracks, regular inspections of electricity transmission easements, restricted vehicle movements, and the prohibition of smoking in fire prone areas or the lighting of fires or fireworks.

### 7.2.17 Disturbance to Specialist Breeding and Foraging Habitat

The BAM (DPIE 2020) does not define 'specialist breeding and foraging habitat', although gives the example of 'beach nesting for shorebirds'. No specialist breeding and foraging habitat occurs in the Biodiversity Assessment Development Footprint.

### 7.2.18 Groundwater Dependent Vegetation

The Modification does not involve any potential impacts on groundwater.

## 7.3 PRESCRIBED BIODIVERSITY IMPACTS

The BC Regulation identifies actions that are prescribed as impacts to be assessed under the Biodiversity Offsets Scheme. Prescribed Biodiversity Impacts are as follows:

- (a) *the impacts of development on the following habitat of threatened species or ecological communities:*
  - (i) *karst, caves, crevices, cliffs and other geological features of significance,*
  - (ii) *rocks,*
  - (iii) *human made structures,*
  - (iv) *non-native vegetation,*
- (b) *the impacts of development on the connectivity of different areas of habitat of threatened species that facilitates the movement of those species across their range,*
- (c) *the impacts of development on movement of threatened species that maintains their lifecycle,*
- (d) *the impacts of development on water quality, water bodies and hydrological processes that sustain threatened species and threatened ecological communities (including from subsidence or upsidence resulting from underground mining or other development),*
- (e) *the impacts of wind turbine strikes on protected animals,*
- (f) *the impacts of vehicle strikes on threatened species of animals or on animals that are part of a threatened ecological community.*

These impacts are assessed below in relation to the Modification.

### 7.3.1 Karst, Caves, Crevices, Cliffs, Rocks and Other Geological Features

As described in Section 2.7, there are no karst, caves, cliffs or other areas of geological significance on, or in the vicinity of, the Biodiversity Assessment Development Footprint. The Modification would avoid clearance of any rocky areas that provide potential habitat for the Pink-tailed Legless Lizard (or any other threatened species) as shown on Figure 13.

### 7.3.2 Human Made Structures

There are no human made structures that provide habitat for threatened species that would be adversely impacted by the Modification.

### 7.3.3 Non-Native Vegetation

There are no areas of non-native vegetation that provide habitat for threatened species that would be adversely impacted by the Modification.

### 7.3.4 Habitat Connectivity

The BAM (DPIE 2020) states the following assessment requirements for habitat connectivity:

*Assessment of the impacts of the proposal on connectivity of habitat of threatened entities must:*

- (a) describe the nature, extent and duration of short-term and long-term impacts*
- (b) predict the consequences of impacts for the persistence of the threatened entities identified in Subsection 6.1.3, taking into consideration mobility, abundance, range and other relevant life history factors*
- (c) justify predictions of impacts with relevant literature and other published sources of information.*

Habitat connectivity areas are shown on Figure 5 and woodland areas within the assessment buffer areas are shown on Figure 6. There are no defined woodland corridors in the Biodiversity Assessment Development Footprint, however, it is possible that the woodland areas facilitate the movement of species in the landscape. It is recognised that the derived native grassland may provide habitat linkages for some species (e.g. Striped Legless Lizard) (Section 7.6.3).

All threatened species and communities known to occur in the Biodiversity Assessment Development Footprint are likely to benefit from the current level of connectivity, in particular species that are known to use habitat linkages, such as the Squirrel Glider, and species that are unlikely to cross roads (e.g. Striped Legless Lizard [Dorrough et al. 1999]).

Fragmentation as a result of the Modification would be negligible because the MEA Extension areas are small (combined approximately 3.8 ha) and adjoin the approved Project disturbance area, and sufficient connectivity would remain around the Biodiversity Assessment Development Footprint (Figure 5) such that no threatened species are likely to become isolated as a result of the Modification.

The surface disturbance areas associated with the Biodiversity Assessment Development Footprint would be rehabilitated and revegetated when the surface facilities are no longer required or at the end of the mine life where no further ongoing beneficial use is identified. The conceptual final rehabilitation plan is shown on Figure 17. Revegetation would aim to increase the continuity of woodland vegetation by establishing links between woodland vegetation.

### 7.3.5 Fauna Movement

Movement patterns key to the life cycle of a threatened species could include seasonal movements between foraging and breeding habitats. The Modification is not likely to impact a well-defined movement pattern for any particular threatened species.

As described in Section 7.3.4, all threatened species and communities known to occur in the Biodiversity Assessment Development Footprint are likely to benefit from the current level of connectivity. However, despite the impact to habitat connectivity, sufficient connectivity would remain around the Biodiversity Assessment Development Footprint (Figure 5) such that no threatened species are likely to become isolated as a result of the Modification.

### 7.3.6 Water Quality, Water Bodies and Hydrological Processes that Sustain Threatened Species and Threatened Ecological Communities

The Modification would not impact water quality, water bodies or hydrological processes that are known to sustain a threatened species or TEC.

As part of the Modification, there would be a small increase in surface development area at the MEA. This results in a small increase in the catchment excision of Saddlers Creek, and a subsequent increase in the loss of catchment runoff (WRM 2021).

The Modification would increase the catchment excision of Saddlers Creek by 4 ha. The total pre-development catchment area of Saddlers Creek is 9,714 ha and, therefore, the Modification would increase the loss of catchment (and hence loss of catchment runoff) by approximately 0.04% (WRM 2021).

The impact of the Modification on catchment excision (and loss of flows) in Saddlers Creek would be negligible and would not be measurable (WRM 2021).

### 7.3.7 Wind Turbines

No wind turbines are planned for the Project (or Modified Project).

### 7.3.8 Vehicle Strike

As described in Section 5.5, the Modification would not change the risk on threatened species from vehicle strike. The revised site access road alignment follows an existing haul road and does not contain native vegetation.

## 7.4 IMPACTS ON THREATENED SPECIES AND COMMUNITIES UNDER THE NSW FISHERIES MANAGEMENT ACT 1994

An aquatic ecology assessment was prepared for the Project by Eco Logical (2019) in consideration of the requirements under the FM Act and the *Threatened Species Assessment Guidelines: the Assessment of Significance* (DPI, 2008).

Two species, the Purple-spotted Gudgeon (listed as an endangered species under the FM Act) and Darling River Hardyhead in the Hunter Catchment (listed as an endangered population under the FM Act), have modelled distributions along the Hunter River, adjacent to the Project. However, neither of these species were not found during the 2018 field surveys by Eco Logical (2019). Eco Logical (2019) concluded that it is unlikely that the Project would directly or indirectly impact these species, or the habitats that support them, as the Project would have a negligible impact on the flow frequency in the Hunter River and habitat features would not be impacted. Further, no threatened ecological communities listed under the FM Act potentially occur in the Project area and surrounds.

The Modified Project would not directly impact these species, or the habitats. It is also considered that the Modified Project is unlikely to indirectly impact these species, or the habitats because the impact of the Modification on catchment excision (and loss of flows) in Saddlers Creek would be negligible and would not be measurable (WRM 2021) (Section 7.3.6).

## **7.5 STATE ENVIRONMENTAL PLANNING POLICY (KOALA HABITAT PROTECTION)**

The State Environmental Planning Policy (SEPP) (Koala Habitat Protection) 2021 was made and commenced on 17 March 2021. However, at this stage, for all RU1, RU2 and RU3 zoned land outside of the Sydney Metropolitan Area and the Central Coast, the Koala Habitat Protection SEPP 2020 continues to apply.

The land associated with the MEA Extensions is zoned RU1 Rural, and therefore the Koala Habitat Protection SEPP 2020 applies. The land associated with the proposed water treatment facility location is zoned SP2 Infrastructure and neither SEPP applies.

Habitat for the Koala is absent with the MEA Extensions, as no eucalypts are present. Therefore, the Koala Habitat Protection SEPP is not relevant to the Modification.

## **7.6 COMMONWEALTH ENVIRONMENT PROTECTION AND BIODIVERSITY CONSERVATION ACT 1999**

### **7.6.1 Background**

The Project was subsequently approved under the EPBC Act in March 2021 (EPBC 2018/8287). The controlling provisions for the Project were "listed threatened species and communities" (sections 18 and 18A of the EPBC Act) and "a water resource, in relation to coal seam gas development and large coal mining development" (sections 24D and 24E of the EPBC Act).

This BDAR assesses some relatively minor modifications to the approved Project layout. The total amount of native vegetation to be disturbed for the Modification is approximately 4.4 ha (i.e. 2.7 % of the native vegetation approved to be disturbed).

The Australian Government has endorsed the NSW Biodiversity Offsets Scheme. This means any NSW proponent who needs an EPBC Act approval can use the NSW Biodiversity Offsets Scheme to assess and meet their biodiversity offset requirements. The relevant threatened species and communities listed under the EPBC Act are incorporated into the BAM-C and are discussed below.

This section provides a detailed analysis of the nature and extent of the impacts relevant to the following species and communities are assessed below:

- Box-Gum TEC;
- Striped Legless Lizard;
- White-throated Needle-tail
- Spotted-tailed Quoll; and
- Corben's Long-eared Bat.

In addition to those species assessed in the BAM-C, the Pink-tailed Legless Lizard, Grey-headed Flying-fox and Koala are listed as 'known to occur' within the Biodiversity Assessment Development Footprints in the EPBC Act Protected Matters Search (Department of Agriculture, Water and the Environment [DAWE] 2021a). However, Table 9 describes that these species have been excluded from further assessment in this BDAR due to the absence of suitable habitat available in the Biodiversity Assessment Development Footprints.

The assessments below are in accordance with relevant Commonwealth guidelines and policy statements (DAWE 2021b).

### 7.6.2 Box-Gum TEC

*The Box-Gum TEC listed under the EPBC Act is present within the Biodiversity Assessment Development Footprints as the associated PCT 1606 (in derived native grassland form) present in the MEA Extensions (Figure 7a).*

*An action is likely to have a significant impact on a critically endangered or endangered ecological community if there is a real chance or possibility that it will:*

- a) *reduce the extent of an ecological community*

The Modification would result in the clearance of approximately 3.8 ha of Box-Gum TEC represented by PCT 1606 (in derived native grassland form). The clearance would occur in the short-term for construction of the MEA Extensions. As described in Section 6, the clearance of Box-Gum TEC derived native grassland for the Modification is unavoidable, as it covers a large area surrounding the Project surface facilities such that the Modification could not be moved in a way that would avoid or reduce the amount lost.

- b) *fragment or increase fragmentation of an ecological community, for example by clearing vegetation for roads or transmission lines*

Fragmentation of Box-Gum TEC as a result of the Modification would be negligible because the MEA Extension areas are small (combined approximately 3.8 ha) and adjoin the approved Project disturbance area, there would be no direct or indirect isolation of important areas of Box-Gum TEC as it would remain connected around the Biodiversity Assessment Development Footprint of both the Modification and the approved Project (Figures 7a and 7b).

- c) *adversely affect habitat critical to the survival of an ecological community*

The derived native grassland in the Biodiversity Assessment Development Footprint is not *habitat critical to the survival of the Box-Gum TEC listed under the EPBC Act*. The Box-Gum TEC in and adjacent to the Biodiversity Assessment Development Footprints (Figures 7a and 7b) is mostly located in an agricultural grazing property and as such is subject to a number of existing recognised threats, namely, grazing, habitat fragmentation, weed invasion and lack of fire (TSSC 2006; Rawlings et al. 2010; DECCW 2010).



- d) *modify or destroy abiotic (non-living) factors (such as water, nutrients, or soil) necessary for an ecological community's survival, including reduction of groundwater levels, or substantial alteration of surface water drainage patterns*

The derived native grassland (conforming to the Box-Gum TEC) in the Biodiversity Assessment Development Footprint would be cleared. The Modification would not modify or destroy abiotic factors necessary for the survival of the *Box-Gum TEC listed under the EPBC Act* in the surrounds. The MEA Extensions would not alter water availability (primarily from direct rainfall) or change soil nutrients in the areas surrounding the area of immediate loss.

- e) *cause a substantial change in the species composition of an occurrence of an ecological community, including causing a decline or loss of functionally important species, for example through regular burning or flora or fauna harvesting*

The derived native grassland (conforming to the Box-Gum TEC) in the Biodiversity Assessment Development Footprint would be cleared. The Modification would not cause a change in the species composition in the occurrence of the *Box-Gum TEC listed under the EPBC Act* in the surrounds. As described earlier, the occurrence of the ecological community in and adjacent to the MEA Extensions is in a derived native grassland landscape (of average quality as indicated by a VI score of 31.4 [Table 2]) and is subject to a number of existing threats.

- f) *cause a substantial reduction in the quality or integrity of an occurrence of an ecological community, including, but not limited to:*
- i. *assisting invasive species, that are harmful to the listed ecological community, to become established, or*
  - ii. *causing regular mobilisation of fertilisers, herbicides or other chemicals or pollutants into the ecological community which kill or inhibit the growth of species in the ecological community, or*

The Modification is unlikely to cause a reduction in the quality or integrity of the occurrence of the *Box-Gum TEC listed under the EPBC Act* in the surrounds. Mitigation measures for the ecological community are outlined in Section 8 and include the following:

- implementation of a vegetation clearance protocol;
- weed management; and
- bushfire management.

- g) *interfere with the recovery of an ecological community*

The Modification would not interfere with any of the objectives or actions outlined in the *National Recovery Plan for White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland* (DECCW 2010). As such, the Modification is unlikely to interfere with the recovery of the *Box-Gum TEC listed under the EPBC Act*.

## Conclusion

The loss of approximately 3.8 ha of derived native grassland of the *Box-Gum TEC listed under the EPBC Act* is not considered significant in consideration of the *Matters of National Environmental Significance: Significant impact guidelines 1.1* (DotE 2013).

The impacts on the *Box-Gum TEC listed under the EPBC Act* would be offset in accordance with the NSW Biodiversity Offsets Scheme and would result in the retirement of the required number and class of like-for-like ecosystem credits (Section 9.4).

### 7.6.3 Striped Legless Lizard

The Striped Legless Lizard is the only species credit species listed under the EPBC Act that is relevant to the Biodiversity Assessment Development Footprint (Section 4.4.1).

This species was not recorded within the Biodiversity Assessment Development Footprint associated with the Modification during targeted surveys undertaken in 2018 by Future Ecology (2019) (Attachment A), however four individuals were recorded within approximately 1 km of the MEA Extensions (Figure 9). A total of 26 observations of Striped Legless Lizard were recorded in the broader Project study area, including live specimens (16 individuals) and sloughs (10). Most records were associated with the Maxwell Underground and along or near the transport and services corridor between the Maxwell Underground and Maxwell Infrastructure. The majority of observations were made under dried cow manure in open grassy areas with a good cover of native grasses and herbs, though the species was also recorded with pitfall traps and artificial shelter habitat, and associated with rocks and dumped material. An assessment of the significant impact criteria (DotE 2013) for the Striped Legless Lizard is provided below.

*An action is likely to have a significant impact on a vulnerable species if there is a real chance or possibility that it will:*

- a) *lead to a long-term decrease in the size of an important population of a species*

The *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b) recognised that the local population of the Striped Legless Lizard could be considered an important population (as defined by DSEWPac 2011). However, the Modification is not likely to lead to a long-term decrease in the size of the population of the Striped Legless Lizard because the area of potential habitat to be cleared for the Modification (4.3 ha) is small compared to the wider occurrence of known and potential habitat (Figure 14). The potential habitat comprises of open agricultural grazing land with dried cow manure that may provide shelter, no rocky habitat.

As described in Table 10, PCT 1692 at the proposed water treatment facility location is not considered potential habitat for this species as it is highly disturbed and provides no suitable shelter (rocks, dumped material or dried cow manure).

In the long-term, the surface disturbance areas associated with the Biodiversity Assessment Development Footprint would be rehabilitated and revegetated when surface facilities are no longer required (Section 7.3.4). The population of this species would persist in the surrounding locality due to the amount of known habitat and the occurrence of the species outside the Biodiversity Assessment Development Footprint (Figure 14).

- b) *reduce the area of occupancy of an important population*

The Striped Legless Lizard was not recorded within the Biodiversity Assessment Development Footprint. Given its presence in the surrounds, it is possible that the Striped Legless Lizard utilises potential habitat within the Biodiversity Assessment Development Footprint. However, the area of potential habitat to be cleared for the Modification (4.3 ha) is small compared to the wider occurrence of known and potential habitat (Figure 14).

- c) *fragment an existing important population into two or more populations*

Studies indicate that the Striped Legless Lizard only moves across short distances, having been recorded moving at least 20 m in one day (and up to 50 m over several weeks) (DAWE 2021b). The creation of barriers to lizard movements can cause populations to become fragmented (DAWE 2021b). The Modification could potentially limit movement of the Striped Legless Lizard; however, this species was not recorded within the development footprint or close to the access road

at the Maxwell Infrastructure, and the Modification is unlikely to impact movement patterns for this species (Section 7.3.5; Figure 9). Therefore, based on available data, the population is unlikely to be significantly fragmented.

*d) adversely affect habitat critical to the survival of a species*

The *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b) recognised that potential habitat for the species in the Project area may form part of 'habitat critical to the survival of the species' according to the TSSC (2016). The removal of the habitat for the Modification is not likely to be material to the survival of the species given the area of potential to be removed is small (4.3 ha) and comprised of open agricultural grazing land with dried cow manure that may provide shelter, no rocky habitat.

*e) disrupt the breeding cycle of an important population*

Breeding habitat for the Striped Legless Lizard is determined by the presence of two or more adult individuals or juveniles and a habitat assessment of available resources such as tussocks with high biomass, surface rocks or invertebrate burrows (TSSC 2016). It is possible that the species could breed in the Biodiversity Assessment Development Footprint, area of potential to be removed is small (4.3 ha) and better habitat (rocky areas) occur outside of the Biodiversity Assessment Development Footprint (Figure 14).

*f) modify, destroy, remove or isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline*

In the long-term, the surface disturbance areas associated with the Biodiversity Assessment Development Footprint would be rehabilitated and revegetated when surface facilities are no longer required (Section 7.3.4). The population of this species would persist in the surrounding locality due to the amount of known habitat and occurrence of the species outside the Biodiversity Assessment Development Footprint (Figure 14).

*g) result in invasive species that are harmful to a vulnerable species becoming established in the vulnerable species' habitat*

*h) introduce disease that may cause the species to decline, or*

The Modification is unlikely to indirectly impact the Striped Legless Lizard adjacent to the MEA Extensions as the Modification would include measures to manage environmental weeds spreading from the Biodiversity Assessment Development Footprint, considering that the Striped Legless Lizard is a grassland specialist (DAWE 2021b).

Best practice mitigation measures as outlined in the *Environment Protection and Biodiversity Conservation Act 1999 referral guidelines for the vulnerable striped legless lizard, Delma impar* (DSEWPaC 2011) have been considered for the Striped Legless Lizard, and would include (Section 8):

- the salvage and re-use of material for habitat (e.g. surface rocks);
- weed management; and
- feral animal management (e.g. feral cats).

*i) interfere substantially with the recovery of the species*

The Modification would not interfere with any of the objectives, criteria and actions outlined in the *National Recovery Plan for the Striped Legless Lizard (Delma impar) 1999-2003* (Smith and Robertson 1999). As such, the Modification is unlikely to interfere substantially with the recovery of the Striped Legless Lizard.

## **Conclusion**

The loss of approximately 4.3 ha of potential habitat for the Striped Legless Lizard is not considered significant in consideration of the *Matters of National Environmental Significance: Significant impact guidelines 1.1* (DotE 2013).

The impacts on the Striped Legless Lizard would be offset in accordance with the NSW Biodiversity Offsets Scheme and would result in the retirement of the required number and class of like-for-like species credits (Section 9.4).

### **7.6.4 Other Species**

The following ecosystem species listed under the EPBC Act were retained in the BAM-C (Section 4.1.1):

- White-throated Needle-tail
- Spotted-tailed Quoll; and
- Corben's Long-eared Bat.

These are discussed below.

#### ***White-throated Needletail***

The Modification would not impact the White-throated Needletail as this species is almost exclusively aerial in Australia and no breeding habitat impacted. The Modification is not likely to have a significant impact on the White-throated Needletail in consideration of the *Matters of National Environmental Significance Significant Impact Guidelines 1.1*. (DotE 2013).

This species is classified as an 'Ecosystem Credit Species' and as such has been conservatively included in the credit calculations for the Modification.

#### ***Spotted-tailed Quoll (South-eastern Mainland Population)***

All of the native vegetation in the Biodiversity Assessment Development Footprint (4.4 ha) is potential habitat for the Spotted-tailed Quoll (south-eastern mainland population), and local records of the species occur on Saddlers Creek (Figure 11).

Potential habitat for this species is widespread. The Modification is not likely to have a significant impact on the Spotted-tailed Quoll (south-eastern mainland population) in consideration of the *Matters of National Environmental Significance Significant Impact Guidelines 1.1*. (DotE 2013).

This species is classified as an 'Ecosystem Credit Species' and as such has been conservatively included in the credit calculations for the Modification.

#### ***Corben's Long-eared Bat***

There are no hollow bearing trees in the Biodiversity Assessment Development Footprint. The Modification is not likely to have a significant impact on the Corben's Long-eared Bat in consideration of the *Matters of National Environmental Significance Significant Impact Guidelines 1.1*. (DotE 2013).

This species is classified as an 'Ecosystem Credit Species' and as such has been conservatively included in the credit calculations for the Modification.

### 7.6.5 Mitigation Measures Relevant to MNES

Conservation advice, recovery plans and threat abatement plans for relevant EPBC Act-listed species and communities are considered in Table 14. A reconciliation table of all conservation advice, recovery plans and threat abatement plans for relevant EPBC listed species is provided in Table 15.

### 7.6.6 Conclusion

The impacts of the Modification on biodiversity a local scale would be minimal. Any impacts on protected matters listed under the EPBC Act would be localised and negligible on a regional, state and national scale. It is concluded that there is unlikely to be significant residual impact on any threatened species and communities listed under the EPBC Act as a result of the Modification.

**Table 14**  
**Impact Mitigation Measures Relevant to Threatened Species and Communities listed under the EPBC Act**

Matter	Impact	Mitigation Measure	Techniques	Impact Mitigation Measures/Effectiveness	Basis for the Mitigation Measures
White Box – Yellow Box – Blakely’s Red Gum Grassy Woodland and Derived Native Grassland	Clearance Impacts on Native Vegetation and Habitat	Vegetation Clearance Protocol	Areas to be cleared are delineated to prevent accidental damage to adjoining areas during vegetation clearance activities or other works.	Effective if clearly delineated.	Rawlings et al. (2010), Threatened Species Scientific Committee (TSSC) (2006) and Department of Environment, Climate Change and Water (DECCW) (2010) describe protection of the threatened ecological communities (TEC).
	Indirect Impacts on Native Vegetation and Habitat	Weed Management	Where they have been taken off road, washdown of vehicles and mechanical equipment to minimise seed transport off the site.	Effective when done in a controlled manner.	Rawlings et al. (2010), TSSC (2006) and DECCW (2010) describe weed management of the threatened ecological communities (TEC).
			Identification of weeds requiring control.		
			Mechanical removal of identified weeds and/or the application of approved herbicides.		
Follow-up site inspections to determine the effectiveness of the eradication programs.					
		Bushfire Management	According to the Bushfire Management Procedure.	Effective when applied.	Standard practice.
Striped Legless Lizard	Clearance Impacts on Native Vegetation and Habitat	Vegetation Clearance Protocol	Areas to be cleared are delineated to prevent accidental damage during vegetation clearance activities or other works.	Effective if clearly delineated.	Department of Sustainability, Environment, Water, Population and Communities (SEWPaC) (2011).
	Loss of Individuals	Minimise Loss	Pre-clearance fauna surveys by suitably qualified personnel.	Relocation of captured individuals.	SEWPaC (2011).
			Impacts on fauna are managed during clearing activities by suitably qualified personnel.	Relocation of captured individuals.	

**Table 14 (Continued)**  
**Impact Mitigation Measures Relevant to Threatened Species and Communities listed under the EPBC Act**

<b>Matter</b>	<b>Impact</b>	<b>Mitigation Measure</b>	<b>Techniques</b>	<b>Impact Mitigation Measures/Effectiveness</b>	<b>Basis for the Mitigation Measures</b>
Striped Legless Lizard (Continued)	Loss of Habitat	Mine Site Rehabilitation and Revegetation	Surface disturbance areas associated with the Biodiversity Assessment Development Footprint would be rehabilitated and revegetated (when the surface facilities are no longer required or at the end of the mine life where no further ongoing beneficial use is identified).	Effective when applied.	SEWPaC (2011).
	Salvage and Re-Use of Material for Habitat Enhancement within Mine Site Rehabilitation	Identification of habitat features (e.g. surface rocks) that would be beneficial for habitat enhancement.	Effective when applied.		
	Indirect Impacts on Habitat	Feral Animal Management	Maintain a clean, rubbish-free environment to discourage scavenging and reduce the potential for colonisation of these areas by non-endemic fauna.	Effective if ongoing during development and operational stages.	SEWPaC (2011).
	Uncontrolled Spread of Weeds	Weed Management	Where they have been taken off road, washdown of vehicles and mechanical equipment to minimise seed transport off the site. Identification of weeds requiring control. Mechanical removal of identified weeds and/or the application of approved herbicides. Follow-up site inspections to determine the effectiveness of the eradication programs.	Effective when applied.	SEWPaC (2011).

**Table 15**  
**Summary of Conservation Advice, Recovery Plans or Threat Abatement Plans for EPBC Listed Species in the Project Area**

<b>Matter</b>	<b>Conservation Advice, Recovery Plans and Threat Abatement Plans (EPBC Species Profile and Threats Database)</b>
White Box – Yellow Box – Blakely’s Red Gum Grassy Woodland and Derived Native Grassland	<ul style="list-style-type: none"> <li>• No approved Conservation Advice.</li> <li>• <i>National Recovery Plan for White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland</i> (NSW DECCW, 2010).</li> <li>• <i>Threat abatement plan for the biological effects, including lethal toxic ingestion, caused by cane toads</i> (DSEWPAC, 2011).</li> <li>• <i>Threat abatement plan for predation, habitat degradation, competition and disease transmission by feral pigs (Sus scrofa)</i> (2017) (DEE, 2017).</li> <li>• <i>Threat abatement plan for disease in natural ecosystems caused by Phytophthora cinnamomic</i> (DEE, 2018).</li> </ul>
Striped Legless Lizard	<ul style="list-style-type: none"> <li>• <i>Conservation Advice Delma impar striped legless lizard</i> (TSSC, 2016).</li> <li>• <i>National Recovery Plan for the Striped Legless Lizard (Delma impar) 1999-2003</i> (Smith, W.J.S. &amp; P. Robertson, 1999).</li> <li>• <i>Threat abatement plan for predation by feral cats</i> (DoE, 2015).</li> <li>• <i>Threat abatement plan for competition and land degradation by rabbits</i> (DoEE, 2016).</li> <li>• <i>Threat abatement plan for predation by the European red fox</i> (DEWHA, 2008).</li> </ul>



## 8 MEASURES TO MITIGATE AND MANAGE IMPACTS

The measures to mitigate and manage impacts from the Project as described in the *Maxwell Project Biodiversity Development Assessment Report* (Hunter Eco 2019b) (Table 16) would continue to be applied to the Modified Project.

**Table 16**  
**Measures to Mitigate and Manage Impacts**

Potential Impact	Mitigation/Management Measure	Techniques	Timing/Frequency	Potential Risk of Failure	Likelihood and Consequence of Residual Impacts
Displacement of Fauna	Presence of a Trained Ecological or Licensed Wildlife Handler	Capture and release.	During native vegetation clearance and clearance of rocky areas.	Low.	Low risk of a smaller portion of resident fauna becoming displaced.
Clearance Impacts on Native Vegetation and Habitat	Vegetation Clearance Protocol	Areas to be cleared are delineated to prevent accidental damage during vegetation clearance activities or other works.	During native vegetation clearance and clearance of rocky areas.	Low.	Low risk of a smaller portion of resident fauna becoming displaced or injured.
		Pre-clearance fauna surveys by suitably qualified personnel.	During native vegetation clearance and clearance of rocky areas.	Low.	
		Impacts on fauna are managed during clearing activities by suitably qualified personnel.	During native vegetation clearance and clearance of rocky areas.	Low.	
		Review of environmental impacts that may result from subsidence remediation (threatened flora species and populations, rocky areas that may provide habitat for threatened lizards) and consideration of whether alternative methods of remediation are warranted (e.g. without machinery).	Prior to any remediation of surface cracks.	Low.	Low.

Potential Impact	Mitigation/Management Measure	Techniques	Timing/Frequency	Potential Risk of Failure	Likelihood and Consequence of Residual Impacts
Clearance Impacts on Native Vegetation and Habitat (continued)	Vegetation Clearance Protocol (continued)	Restricting vegetation clearance to the slashing of vegetation where possible along power line easements (i.e. leaving the lower stem and roots in-situ to maximise the potential for natural regrowth).	During vegetation clearance.	Low.	Vegetation clearance is quantified in Table 13 (no reduction has been applied accounting for these measures).
		Lopping of branches, rather than the removal of trees where possible along power line easements.	During vegetation clearance.	Low.	
	Mine Site Rehabilitation and Revegetation	Surface disturbance areas associated with the Biodiversity Assessment Development Footprint would be rehabilitated and revegetated.	Over the life of the Project. Surface facilities used for the Project would be decommissioned when they are no longer required or at the end of the mine life where no further ongoing beneficial use is identified.	Low.	None.
	Salvage and Re-Use of Material for Habitat Enhancement within the Mine Site Rehabilitation	Identification of habitat features (e.g. cleared trees, surface rocks) that would be beneficial for habitat enhancement.	During and after vegetation clearance.	Moderate.	Low.
	Site Induction	Where possible, encourage Malabar personnel to use existing tracks for site access to Project areas to minimise potential disturbance of soils and revegetated areas.	During construction and operational stages.	Low.	Low.
	Access	Use of defined tracks to access sites to minimise the disturbance of soils.	During construction and operational stages.	Low.	Low.

Potential Impact	Mitigation/Management Measure	Techniques	Timing/Frequency	Potential Risk of Failure	Likelihood and Consequence of Residual Impacts
Subsidence Impacts on Native Vegetation and Habitat	Remediation of Surface Cracks	Remediation of mine subsidence effects (e.g. surface cracking and minor erosion).	As required, where impacts are identified as part of the subsidence monitoring program.	Low.	Low.
Indirect Impacts on Native Vegetation and Habitat	Feral Animal Management	Maintain a clean, rubbish-free environment to discourage scavenging and reduce the potential for colonisation of these areas by non-endemic fauna.	During construction and operational stages.	Low.	Low.
	Weed Management	Where they have been off road, washdown of vehicles and mechanical equipment to minimise seed transport off the site.	During construction and operational stages.	Moderate.	Low.
		Identification of weeds requiring control.	Regular site inspections.	Moderate.	
		Mechanical removal of identified weeds and/or the application of approved herbicides.	During construction and operational stages.	Moderate.	
		Follow-up site inspections to determine the effectiveness of the eradication programs.	During construction and operational stages.	Moderate.	
	Bushfire Management	According to the Bushfire Management Procedure.	During construction and operational stages.	Low.	Low.
Vehicle Strike	Fencing	Fencing along the length of the site access road to exclude kangaroos (and cattle).	Installation during construction of the site access road.	Low.	Low.
	Speed Limits	Imposing a maximum 60 km per hour speed limit on internal roads and maximum 80 km per hour speed limit on the sealed site access road.	During the construction and operational stages.	Moderate.	Low.

## 9 IMPACT SUMMARY

### 9.1 SERIOUS AND IRREVERSIBLE IMPACTS


Under the BC Act, a determination of whether an impact is serious and irreversible must be made for 'potential SAI entities' identified in the BAM Credit Calculator. There is one 'potential SAI entity' relevant to the Modification, namely the Box-Gum TEC (Figure 18).

#### 9.1.1 BAM Requirements

The BAM (DPIE 2020) requires the following information to be provided:

##### *9.1.1 Additional impact assessment provisions for threatened ecological communities at risk of an SAI*

1. *The assessor is required to provide further information in the BDAR or BCAR regarding the impacts on each TEC at risk of an SAI. This must include the action and measures taken to avoid the direct and indirect impact on the TEC at risk of an SAI. Where these have been addressed elsewhere the assessor can refer to the relevant sections of the BDAR and BCAR.*
2. *The assessor must consult the TBDC and/or other sources to report on the current status of the TEC including:*
  - a. *evidence of reduction in geographic distribution (Principle 1, clause 6.7(2)(a) BC Regulation) as the current total geographic extent of the TEC in NSW AND the estimated reduction in geographic extent of the TEC since 1970 (not including impacts of the proposal)*
  - b. *extent of reduction in ecological function for the TEC using evidence that describes the degree of environmental degradation or disruption to biotic processes (Principle 2, clause 6.7(2)(b) BC Regulation) indicated by:*
    - i. *change in community structure*
    - ii. *change in species composition*
    - iii. *disruption of ecological processes*
    - iv. *invasion and establishment of exotic species*
    - v. *degradation of habitat, and*
    - vi. *fragmentation of habitat*
  - c. *evidence of restricted geographic distribution (Principle 3, clause 6.7(2)(c) BC Regulation), based on the TEC's geographic range in NSW according to the:*
    - i. *extent of occurrence*
    - ii. *area of occupancy, and*
    - iii. *number of threat-defined locations*
  - d. *evidence that the TEC is unlikely to respond to management (Principle 4, clause 6.7(2)(d) BC Regulation).*
3. *Where the TBDC indicates data is 'unknown' or 'data deficient' for a TEC for a criterion listed in Subsection 9.1.1(2.), the assessor must record this in the BDAR or BCAR.*
4. *In relation to the impacts from the proposal on the TEC at risk of an SAI, the assessor must include data and information on:*
  - a. *the impact on the geographic extent of the TEC (Principles 1 and 3) by estimating the total area of the TEC to be impacted by the proposal:*
    - i. *in hectares, and*
    - ii. *as a percentage of the current geographic extent of the TEC in NSW.*



- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
  - Threatened Ecological Communities
  - 1 White Box - Yellow Box - Blakely's Red Gum Woodland

Source: MSEC (2019); NSW Spatial Services (2021)  
 Orthophoto Mosaic: 2020, 2019

**MALABAR**  
 MAXWELL UNDERGROUND MINE PROJECT  
 Serious and Irreversible Impact Entities

**Figure 18**

*Data and information should include direct impacts (i.e. from clearing) and indirect impacts where partial loss of the TEC is likely as a result of the proposal. The assessor should consider for example, changes to fire regime (frequency, severity), hydrology, pollutants, species interactions (increased competition, changes to pollinators or dispersal), fragmentation, increased edge effects and disease, pathogens and parasites, which are likely to contribute to the loss of flora and/or fauna species characteristic of the TEC*

- b. *the extent that the proposed impacts are likely to contribute to further environmental degradation or the disruption of biotic processes (Principle 2) of the TEC by:
 
    - i. *estimating the size of any remaining, but now isolated, areas of the TEC; including areas of the TEC within 500 m of the development footprint or equivalent area for other types of proposals*
    - ii. *describing the impacts on connectivity and fragmentation of the remaining areas of TEC measured by:
 
      - *distance between isolated areas of the TEC, presented as the average distance if the remnant is retained AND the average distance if the remnant is removed as proposed, and*
      - *estimated maximum dispersal distance for native flora species characteristic of the TEC, and*
      - *other information relevant to describing the impact on connectivity and fragmentation, such as the area to perimeter ratio for remaining areas of the TEC as a result of the development**
    - iii. *describing the condition of the TEC according to the vegetation integrity score for the relevant vegetation zone(s) (Section 4.3). The assessor must also include the relevant composition, structure and function condition scores for each vegetation zone.**
5. *The assessor may also provide new information that demonstrates that the principle identifying that the TEC is at risk of an SAII is not accurate.*

The Box-Gum TEC is a 'potential SAII entity' due to Principle 1 and 2 in the DPIE (2019) *Guidance and Criteria to Assist a Decision Maker to Determine a Serious and Irreversible Impact*.

These are addressed below.

**1. The assessor is required to provide further information in the BDAR or BCAR regarding the impacts on each TEC at risk of an SAII.**

Approximately 3.8 ha of Box-Gum TEC would be cleared, represented by PCT 1606 (in derived native grassland form) currently used for grazing livestock (Vegetation Zone 1 with a VI Score of 31.4) (Table 2). The clearance of Box-Gum TEC for the Modification is unavoidable, as it covers a large area surrounding the Project surface facilities such that the Modification could not be moved in a way that would avoid or reduce the amount lost.

**2. The assessor must consult the TBDC and/or other sources to report on the current status of the TEC**

Principle 1, clause 6.7(2)(a) of the BC Regulation states that:

- (2) *An impact is to be regarded as serious and irreversible if it is likely to contribute significantly to the risk of a threatened species or ecological community becoming extinct because—*
  - (a) *it will cause a further decline of the species or ecological community that is currently observed, estimated, inferred or reasonably suspected to be in a rapid rate of decline*

Considering historical, recent and contemporary clearing in combination, the TSSC estimates that the Box-Gum TEC has experienced a distribution reduction exceeding 90% across its entire range. The current total geographic extent of the TEC in NSW is estimated to be 70,280,000 ha (TSSC 2020).

Principle 2, clause 6.7(2)(b) of the BC Regulation further defines a serious and irreversible impact as contributing to the decline of a TEC because:

- (b) *it will further reduce the population size of the species or ecological community that is currently observed, estimated, inferred or reasonably suspected to have a very small population size*

Box-Gum TEC has experienced a high degree of reduction in ecological function as a result of environmental degradation and disruption to biotic processes. It is estimated that less than 10% of the original distribution of the community has avoided the long-term impacts of pastoralism, and compositional changes associated with grazing and management practices continue to impact remnant areas of Box-Gum TEC. The TSSC (2020) states that the community is subject to a number of threatening processes that have caused severe declines in biotic processes and interactions, including grazing, pasture improvement, dryland salinity, elevated soil nitrogen and prolonged absence of fire. These threats have been associated with the invasion and establishment of exotic species, *Eucalyptus* spp. dieback, habitat fragmentation, and changes in community structure and species composition (TSSC, 2020).

**3. Where the TBDC indicates data is 'unknown' or 'data deficient' for a TEC for a criterion listed in Subsection 9.1.1(2.), the assessor must record this in the BDAR or BCAR.**

The TBDC (DPIE 2021a) does not indicate data is 'unknown' or 'data deficient' for the Box-Gum TEC.

**4. In relation to the impacts from the proposal on the TEC at risk of an SAII, the assessor must include data and information on:**

- a. **the impact on the geographic extent of the TEC (Principles 1 and 3) by estimating the total area of the TEC to be impacted by the proposal:**
- b. **the extent that the proposed impacts are likely to contribute to further environmental degradation or the disruption of biotic processes (Principle 2) of the TEC by:**

Approximately 3.8 ha of Box-Gum TEC would be cleared for the proposed Modification, represented by PCT 1606 (in derived native grassland form) (Vegetation Zone 1 with a VI Score of 31.4) (Table 2). This is a small area considering the current total geographic extent of the TEC in NSW is estimated to be 70,280,000 ha (<0.00000001 %) (TSSC, 2020).

Clearance of native vegetation and the construction of the MEA Extensions is unlikely to cause any change to fire regimes, pollutants, or disease, pathogens and parasites. Potential impacts of clearing of the Box-Gum TEC by the Modification would be fundamentally surficial, leading to no impact on any deep groundwater resources or surface flow patterns, with the latter being managed through the Water Management Plan.

Fragmentation of Box-Gum TEC as a result of the Modification would be negligible because the MEA Extension areas are small (combined 3.8 ha) and adjoin the approved Project disturbance area, and there would be no direct or indirect isolation of important areas of Box-Gum TEC as it would remain connected around the Biodiversity Assessment Development Footprint of both the Modification and the approved Project (Figures 7a and 7b).

The condition of the Box-Gum TEC remaining outside of the Biodiversity Assessment Development Footprint would be at least retained as it is currently. For example, there would be no intrusion by firewood collectors or bush rock collectors. Potential invasion by weed species along the edges of the cleared areas would be managed through the Biodiversity Management Plan, which would also provide controls for the use of herbicides and fertilisers. As such, the MEA Extensions are unlikely to contribute to loss of flora or fauna species characteristic of the Box-Gum TEC.



**5. The assessor may also provide new information that demonstrates that the principle identifying that the TEC is at risk of an SAI is not accurate.**

Not applicable.

### 9.1.2 BC Regulation Requirements

Section 6.7 (2) of the BC Regulation provides principles for the purposes of determining whether an impact on diversity values is a serious and irreversible impact for the purposes of the Biodiversity Offsets Scheme. It states:

- (2) *An impact is to be regarded as serious and irreversible if it is likely to contribute significantly to the risk of a threatened species or ecological community becoming extinct because:*
  - (a) *it will cause a further decline of the species or ecological community that is currently observed, estimated, inferred or reasonably suspected to be in a rapid rate of decline, or*
  - (b) *it will further reduce the population size of the species or ecological community that is currently observed, estimated, inferred or reasonably suspected to have a very small population size, or*
  - (c) *it is an impact on the habitat of the species or ecological community that is currently observed, estimated, inferred or reasonably suspected to have a very limited geographic distribution, or*
  - (d) *the impacted species or ecological community is unlikely to respond to measures to improve its habitat and vegetation integrity and therefore its members are not replaceable.*
- (3) *For the purpose of this clause, a decline of a species or ecological community is a continuing or projected decline in:*
  - (a) *an index of abundance appropriate to the taxon, or*
  - (b) *the geographic distribution and habitat quality of the species or ecological community.*

These are addressed below in consideration of the DPIE (2019) *Guidance and Criteria to Assist a Decision Maker to Determine a Serious and Irreversible Impact*.

***Will the Modification cause a further decline of the species or ecological community that is currently observed, estimated, inferred or reasonably suspected to be in a rapid rate of decline?***

Adherence to the NSW Biodiversity Offset Scheme would result in the retirement of the required number and class of like-for-like biodiversity credits for the derived grassland that conforms to the Box-Gum TEC (Section 9.4).

***Will the Modification further reduce the population size of the species or ecological community that is currently observed, estimated, inferred or reasonably suspected to have a very small population size?***

As described in Section 9.1.1(4), Box-Gum TEC does not have a very small population size with approximately 6,561 ha being mapped within the Hunter sub-region, as well as having a State-wide distribution.

***Will the Modification impact on the habitat of the species or ecological community that is currently observed, estimated, inferred or reasonably suspected to have a very limited geographic distribution?***

Box-Gum TEC is found across NSW so does not have a limited geographic distribution.

***Is the community unlikely to respond to measures to improve its habitat and vegetation integrity and therefore its members are not replaceable?***

The derived grassland that conforms to the Box-Gum TEC in the Biodiversity Assessment Development Footprint is comprised of flora species that readily seed and germinate under suitable conditions. The Box-Gum TEC has been shown to respond well to both natural regeneration where threats such as grazing and fire are managed, and to assisted natural regeneration with supplementary planting of appropriate species (NSW National Parks and Wildlife Service 2002).

### **9.1.3 Conclusion**

The Modification is unlikely to have a serious and irreversible impact on the Box-Gum TEC given the small area of derived grassland that conforms to the Box-Gum TEC in the Biodiversity Assessment Development Footprint (combined 3.8 ha) relative to the wider occurrence of the Box-Gum TEC (Figure 7a). Adherence to the NSW Biodiversity Offset Scheme would result in the retirement of the required number and class of like-for-like biodiversity credits for the Box-Gum TEC (Section 9.4).

## **9.2 IMPACTS REQUIRING/NOT REQUIRING AN OFFSET**

The VI Score for all vegetation zones is sufficiently high that all vegetation zones require an offset (Figures 19 and 21).


## **9.3 AREAS NOT REQUIRING ASSESSMENT**

The revised site access road alignment follows an existing haul road and does not contain native vegetation or threatened species habitat, as such, it does not require assessment in this BDAR (Figures 7b and 20). Similarly, the majority of the water treatment facility location has been disturbed by past mining activities and is devoid of native vegetation or habitat for threatened species (Figure 7b).

## **9.4 BIODIVERSITY CREDITS**

Table 17 provides a summary of the ecosystem credits required for the Modification. The change in VI Score is provided in Table 13 and the future score is assumed to be 'zero'.

Table 18 provides a summary of the species credits required for the Modification.




- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
  - Dry Sclerophyll Forests (Shrub/grass sub-formation)
  - 2a. White Box - Ironbark - Red Gum Shrubby Forest - Derived Native Grassland (PCT1606)
  - Grassy Woodlands
  - 6. Bull Oak Grassy Woodland (PCT1692)

Source: NSW Spatial Services (2021); MSEC (2019)  
 Orthophoto Mosaic: 2020, 2019

**MALABAR**  
 MAXWELL UNDERGROUND MINE PROJECT  
 Areas Requiring Offset

**Figure 19**




- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
  - Infrastructure/Cleared Land

Source: NSW Spatial Services (2021); MSEC (2019)  
 Orthophoto Mosaic: 2020, 2019

**MALABAR**  
 MAXWELL UNDERGROUND MINE PROJECT  
 Areas Not Requiring Assessment

**Figure 20**



- LEGEND**
- Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Mining Lease Application Boundary
  - Approved Surface Development Area
  - Approved Extent of Conventional Subsidence from Underground Mining
  - Biodiversity Assessment Development Footprint
  - Existing Conservation/Offset Area
  - Infrastructure/Cleared Land

Source: NSW Spatial Services (2021); MSEC (2019)  
 Orthophoto Mosaic: 2020, 2019

**MALABAR**  
 MAXWELL UNDERGROUND MINE PROJECT  
 Areas Not Requiring Offset

**Figure 21**

**Table 17**  
**Modification Ecosystem Credit Requirements**

PCT	PCT Name	Area (ha)		Credits Required <sup>1</sup>		
		Proposed MEA Extensions	Proposed Water Treatment Facility Location	Proposed MEA Extensions	Proposed Water Treatment Facility Location	Total
1606	White Box – Narrow-leaved Ironbark – Blakely’s Red Gum Shrubby Open Forest of the Central and Upper Hunter – DNG <sup>2</sup>	3.8	0	75	0	75
1692	Bull Oak Grassy Woodland of the Central Hunter Valley	0.5	0.1	11	2	13
<b>Total</b>		<b>4.3</b>	<b>0.1</b>	<b>86</b>	<b>2</b>	<b>88</b>

<sup>1</sup> Refer to Attachments E, F and G.

<sup>2</sup> Listed as Box-Gum TEC under the BC Act and EPBC Act.

**Table 18**  
**Modification Species Credit Requirements**

Species Credit Species	Area of Habitat (ha)		Credits Required <sup>1</sup>		
	Proposed MEA Extensions	Proposed Water Treatment Facility Location	Proposed MEA Extensions	Proposed Water Treatment Facility Location	Total
Striped Legless Lizard	4.3	0	54	0	55
Squirrel Glider	1.1	0	17	0	17
<b>Total</b>		<b>71</b>	<b>0</b>	<b>72</b>	<b>72</b>

<sup>1</sup> Refer to Attachments E, F and G.

## 10 CONCLUSION

A BDAR was prepared by Hunter Eco for the Project in July 2019 and Development Consent SSD 9526 was granted in December 2020. The total amount of native vegetation to be disturbed for the Project is approximately 161.6 ha.

This BDAR assesses some relatively minor modifications to the approved Project layout, using the extensive information from the previous BDAR along with supplementary sampling. The total amount of native vegetation to be disturbed for the Modification is approximately 4.4 ha (i.e. 2.7 % of the native vegetation approved to be disturbed for the Project).

The revised site access road alignment follows an existing haul road and does not contain native vegetation or threatened species habitat, as such, it is not assessed in this BDAR. The native vegetation in the extension to the portal entry and extension to the clean water diversion bank consists of derived native grassland or re-growth Buloke woodland. The majority of the water treatment facility location has been disturbed by past mining activities with small fragmented groups of Buloke trees.

The credit calculation has determined the offset requirement for clearance of native vegetation (PCT 1606 [Box-Gum TEC] and PCT 1692) (ecosystem credit requirement) and the offset requirement for clearance of habitat for the Striped Legless Lizard and Squirrel Glider (species credit requirements).

## 11 REFERENCES

Atlas of Living Australia (2021) *Atlas of Living Australia Database with Search Co-ordinates -32.3136 150.7728, -32.3136 150.9855, -32.4929 150.9855, -32.4929 150.7728*. Website: <http://www.ala.org.au> [Accessed May 2021].

Bell, S. (2020) *Expert Report Expected Presence of Threatened Terrestrial Orchids (Diuris Tricolor, Prasophyllum Petilum, Pterostylis Chaetophora) Maxwell Project*. Prepared by Eastcoast Flora Survey.

Birdlife Australia (2018) *Birdlife Australia Atlas Database with Search Co-ordinates -32.3136 150.7728, -32.3136 150.9855, -32.4929 150.9855, -32.4929 150.7728*.

Cumberland Ecology (2009a) *Ecological Assessment of Section 75W Modification for Drayton Mine*.

Cumberland Ecology (2009b) *Mt Arthur Coal Consolidation Project Ecological Assessment*. A report prepared by Cumberland Ecology for Hansen Bailey Pty Ltd.

Cumberland Ecology (2010) *Mt Arthur Coal Flora and Fauna Monitoring Program Spring 2010 Ecological Monitoring Report*. A draft report prepared by Cumberland Ecology for Mt Arthur Coal.

Department of Agriculture, Water and the Environment (2021a) *Protected Matters Search Tool*. Accessed 29 July 2021 August 2021. Website:

Department of Agriculture, Water and the Environment (2021b) *Species Profile and Threats Database*. Website: <http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl>.

Department of Environment, Climate Change and Water (2010) *National Recovery Plan for White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland*. Department of Environment, Climate Change and Water New South Wales, Sydney. Website: <http://www.environment.gov.au/biodiversity/threatened/recovery-plans/white-box-yellow-box-blakelys-red-gum-grassy-woodland-and-derived-native-grassland-national>. In effect under the EPBC Act from 22-Mar-2013.

Department of Planning, Industry and Environment (2019) *Guidance and Criteria To Assist A Decision Maker To Determine A Serious And Irreversible Impact*.

Department of Planning, Industry and Environment (2020) *Biodiversity Assessment Method*, NSW Government DPIE Sydney, October 2020.

Department of Planning, Industry and Environment (2021a) *Threatened Biodiversity Data Collection*. Viewed in June 2021. Website: [https://www.environment.nsw.gov.au/atlaspublicapp/UI\\_Modules/ATLAS\\_/AtlasSearch.aspx](https://www.environment.nsw.gov.au/atlaspublicapp/UI_Modules/ATLAS_/AtlasSearch.aspx).

Department of Planning, Industry and Environment (2021b) *BioNet Vegetation Classification*. Website: <http://www.environment.nsw.gov.au/NSWVCA20PRapp/LoginPR.aspx?ReturnUrl=%2fNSWVCA20PRapp%2fdefault.aspx>.

Department of Planning, Industry and Environment (2021c) *BioNet Atlas*. -32.3136 150.7728, -32.3136 150.9855, -32.4929 150.9855, -32.4929 150.7728. Viewed in June 2021. Website: [https://www.environment.nsw.gov.au/atlaspublicapp/UI\\_Modules/ATLAS\\_/AtlasSearch.aspx?who=0b679421-e424-47ea-b672-f30693729a7e](https://www.environment.nsw.gov.au/atlaspublicapp/UI_Modules/ATLAS_/AtlasSearch.aspx?who=0b679421-e424-47ea-b672-f30693729a7e).

Department of Planning, Industry and Environment (2021d) *BioNet Threatened Species Profile Database*. Viewed in June 2021. Website: <https://www.environment.nsw.gov.au/threatenedspeciesapp/>.



Department of Primary Industries (2008) *Threatened Species Assessment Guidelines: the Assessment of Significance*.

Department of Primary Industries – Water (2017) *Strahler Stream Order Mapping*.

Department of Sustainability, Environment, Water, Population and Communities (2011) Environment Protection and Biodiversity Conservation Act 1999 Referral Guidelines for the Vulnerable Striped Legless Lizard, *Delma impar*.

Department of Sustainability, Environment, Water, Population and Communities (2011) *Environment Protection and Biodiversity Conservation Act 1999 Referral Guidelines for the Vulnerable Striped Legless Lizard, Delma impar*.

Department of the Environment (2012) *Interim Biogeographic Regionalisation for Australia (Sub-regions) v.7*. Website: <http://intspat01.ris.environment.gov.au/fed/catalog/search/resource/details.page?uuid=%7BD470D2-FA7E-4792-B524-0250EC3AF5B3%7D>.

Department of the Environment (2013) Matters of National Environmental Significance Significant Impact Guidelines 1.1. Accessed: October 2015. Website: [https://www.environment.gov.au/system/files/resources/42f84df4-720b-4dcf-b262-48679a3aba58/files/nes-guidelines\\_1.pdf](https://www.environment.gov.au/system/files/resources/42f84df4-720b-4dcf-b262-48679a3aba58/files/nes-guidelines_1.pdf)

Department of the Environment and Energy (2017) *Threat abatement plan for predation, habitat degradation, competition and disease transmission by feral pigs (Sus scrofa)*.

Department of the Environment and Energy (2018a) *Directory of Important Wetlands of Australia*. Website: <https://www.environment.gov.au/wetlands>.

Department of the Environment and Energy (2018b) *Protected Matters Search within a 15km buffer of point coordinates -32.415270° and 150.858402°* being the nominal centre of the Project study area. Data Received: July 2018.

Dorrough, J., & Ash, J. E. (1999) *Using past and present habitat to predict the current distribution and abundance of a rare cryptic lizard, Delma impar (Pygopodidae)*. Australian Journal of Ecology, 24(6), 614-624.

Eco Logical Australia (2019) *Maxwell Project – Aquatic Ecology and Stygofauna Assessment*. Prepared for Malabar Coal Limited.

Future Ecology (2019) *Maxwell Project Baseline Fauna Survey Report*. Prepared for Malabar Coal Limited.

Geoscience Australia (1997) *Australia's River Basins*. Bioregional Assessment Source Dataset. Website: <http://data.bioregionalassessments.gov.au/dataset/d4561e86-2d13-4dcb-bf9e-aae75ba4850c>.

Hansen Bailey (2007) *Drayton Mine Extension Flora and Fauna Impact Assessment*.

Hunter Eco (2019a) *Maxwell Project Baseline Flora Report*. Prepared for Malabar Coal Limited.

Hunter Eco (2019b) *Maxwell Project Biodiversity Assessment Report*. Prepared for Malabar Coal Limited.

Isbell (2016) *The Australian Soil Classification*. The National Committee on Soil and Terrain. CSIRO Publishing January 2016.

Keith, D. (2004) *Ocean shores to desert dunes: The native vegetation of New South Wales and the ACT*. Department of Environment and Conservation.

Mine Subsidence Engineering Consultants [MSEC] (2019) *Maxwell Project: Environmental Impact Statement – Subsidence Assessment*.

Mitchell, P. (2002) *NSW Landscapes Mapping: Background and Methodology*. Report prepared for the NSW National Parks and Wildlife Service.

National Parks and Wildlife Service (2002) *White Box-Yellow Box-Blakely's Red Gum (Box-Gum) Woodland Fact-Sheet*.

Nearmap (2019) Photomaps by Nearmap. Website: <https://www.nearmap.com.au/>.

Rawlings, K., Freudenberger, D. and Carr, D. (2010) *A Guide to Managing Box Gum Grassy Woodlands*. Commonwealth of Australia 2010.

Smith, W. J. S., & Robertson, P. (1999) *National Recovery Plan for the Striped Legless Lizard (Delma impar) 1999-2003*

Threatened Species Scientific Committee (2006) *Commonwealth Listing Advice on White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland*. Available from: <http://www.environment.gov.au/biodiversity/threatened/communities/box-gum.html>.

Threatened Species Scientific Committee (2016) *Conservation Advice Delma impar striped legless lizard*. Canberra: Department of the Environment and Energy. Website: <http://www.environment.gov.au/biodiversity/threatened/species/pubs/1649-conservation-advice-16122016.pdf>.

Threatened Species Scientific Community (2020) *White Box Yellow Box Blakely's red gum woodland - Threatened Species Scientific Committee final determination*. Available from: <https://www.environment.nsw.gov.au/-/media/OEH/Corporate-Site/Documents/Animals-and-plants/Scientific-Committee/Determinations/2020/white-box-yellow-box-final-determination-ceec.pdf?la=en&hash=DD6076E55435D715E7E90B1A901EEB83D488563B>

Umwelt Environmental Consultants (2006) *Ecological Assessment Proposed South Pit Extension*.

Umwelt Environmental Consultants (2007) *Ecological Assessment - Proposed Mt Arthur Underground Project*. A report prepared for Mt Arthur Coal. December 2007.

WRM Water & Environment Pty Ltd (2019) *Maxwell Project Surface Water Assessment*.

WRM Water & Environment Pty Ltd (2021) *Maxwell Underground Mine Project Mine Entry Area Modification Surface Water Impact Assessment*.

ATTACHMENT A  
MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT



FUTURE  
ECOLOGY

ABN 20 117 362 311  
546 Bellingen Rd  
Missabotti NSW 2449  
M. 0423 981 786  
E. [garon@futureecology.com.au](mailto:garon@futureecology.com.au)

# MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Prepared for Malabar Coal Limited

**June 2019**

## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

<b>PROJECT NUMBER</b>	2018-01		
<b>PROJECT NAME</b>	Maxwell Project Baseline Fauna Survey Report		
<b>PROJECT ADDRESS</b>	Jerrys Plains NSW		
<b>PREPARED FOR</b>	Malabar Coal Limited		
<b>AUTHOR/S</b>	Garon Staines		
<b>VERSION</b>	<b>Version</b>	<b>Draft/Final</b>	<b>Date to client</b>
	1.0	Draft V1	26/02/2019
		Draft V1B	30/3/2019
		Draft V1C	17/4/2019
		Final V1	25/6/2019

This report should be cited as: '*Future Ecology (2019) Malabar Project Baseline Fauna Survey Report. Prepared for Malabar Coal Limited*'.

### Disclaimer

This report has been prepared by Future Ecology for Malabar Coal Limited and may only be used for the purpose agreed between these parties, as described in this report. The opinions, conclusions and recommendations set out in this report are limited to those set out in the scope of works and agreed between these parties. Future Ecology accepts no responsibility or obligation for any third party that may use this information or for conclusions drawn from this report that are not provided in the scope of works or following changes occurring subsequent to the date that the report was prepared.

The recommendations provided in this report are based on the results from currently accepted and naturally limited ecological survey techniques. Every effort is made and reasonable care taken to detect all threatened species that may have potential to occur in the locality.

Any representation, statement, opinion or advice expressed or implied in this report is made in good faith on the basis that Future Ecology Pty Ltd, its agents and employees are not liable (whether by reason of negligence, lack of care or otherwise) to any person for any damage or loss whatsoever which has occurred or may occur in relation to that person taking or not taking (as the case may be) action in respect of any representation, statement or advice referred to above.

# Abbreviations

ABBR./TERM	DESCRIPTION
mm/cm/m/km/ha	millimetres/centimetres/metres/kilometres/hectares
CEEC	Critically Endangered Ecological Community
CHPP	Coal Handling and Preparation Plant
CL	Coal Lease
DA	Development Application
DBH	Diameter at breast height
DEE	Department of the Environment and Energy
EEC	Endangered Ecological Community
EIS	Environmental Impact Statement
EL	Exploration Licence
EP&A Act	NSW <i>Environmental Planning and Assessment Act, 1979</i>
EPBC Act	Commonwealth <i>Environment Protection and Biodiversity Conservation Act, 1999</i>
IBRA	Interim Biogeographical Regionalisation of Australia
LGA	Local Government Area
masl	metres above sea level
MEA	mine entry area
ML	Mining Lease
NSW	New South Wales
OEH	Office of Environment and Heritage - NSW
BC Act	NSW <i>Biodiversity Conservation Act, 2016</i>
*	Denotes exotic species

# Contents

Executive Summary .....	1
1 Introduction and Project Description .....	1
1.1 Purpose of Report .....	4
1.2 Site Description .....	4
1.2.1 Study Area .....	4
1.2.2 Land Use .....	4
1.2.3 Regional Setting .....	5
1.2.4 Landform and Hydrology .....	5
1.2.5 Vegetation .....	5
1.2.6 Summary of Previous Threatened Species Recorded in the Study Area .....	5
2 Methods.....	11
2.1 Literature and Database Review.....	11
2.2 Relevant Survey Guidelines .....	14
2.3 Field Survey.....	15
2.3.1 Weather, Climate and Astronomical Conditions.....	15
2.3.2 Techniques .....	17
2.3.3 Survey Effort.....	44
2.3.4 Limitations .....	44
2.3.5 Nomenclature .....	44
2.3.6 Targeted Searches for Threatened Fauna .....	46
2.3.7 Species Credit Species Habitat Polygon Mapping .....	46
3 Survey Results .....	56
3.1 Fauna Habitat Types .....	56
3.2 Fauna Species.....	62
3.3 Threatened Fauna Species Listed under the BC Act.....	62
3.3.1 Threatened Fauna Species Recorded During the Surveys.....	62
3.3.2 Other Threatened Fauna Species Previously Recorded within the Study Area .....	94
3.3.3 Other Threatened Fauna Species Not Recorded in the Study Area .....	101
3.4 Potential Koala Habitat – SEPP 44 .....	104
3.5 Threatened Fauna Species Listed under the EPBC Act.....	107
4 Conclusion.....	109
5 References .....	110
Appendix A Fauna Species Detected .....	115
Appendix B Fauna Survey Site Descriptions .....	131
Appendix C Threatened Biodiversity Data Collection Species PCT Associations Sydney Basin – Hunter IBRA Sub-region.....	149

# Figures

Figure 1	Regional Location
Figure 2	Project General Arrangement
Figure 3a	Vegetation Mapping – Maxwell Underground
Figure 3b	Vegetation Mapping – Maxwell Infrastructure
Figure 4	Fauna Survey Sites
Figure 5	Bat Survey Sites
Figure 6	Amphibian Survey Sites
Figure 7a	Broad Fauna Habitat Types – Maxwell Underground
Figure 7b	Broad Fauna Habitat Types – Maxwell Infrastructure
Figure 8	Threatened Reptiles and Amphibians
Figure 9	Threatened Birds
Figure 10	Threatened Mammals
Figure 11	Threatened Bats
Figure 12	Species Credit Species Relevant to the Study Area
Figure 13	Threatened Species Listed under the EPBC Act
Figure 14	Pink-tailed Legless Lizard Species Polygon
Figure 15	Striped Legless Lizard Species Polygon
Figure 16	Painted Honeyeater Potential Habitat
Figure 17	Squirrel Glider Species Polygon
Figure 18	Grey-headed Flying-fox Potential Habitat
Figure 19	Southern Myotis Species Polygon
Figure 20	Swift Parrot Potential Habitat
Figure 21	Spotted-tailed Quoll Potential Habitat
Figure 22	Koala Potential Habitat

# Tables

Table 1: Threatened Fauna Species Known or Predicted to occur in the Locality .....	8
Table 2: Weather and Astronomical Records during Survey Periods .....	16
Table 3: Fauna Survey Sites for the Study Area .....	21
Table 4: Habitat Constraints Identified in the Threatened Biodiversity Data Collection .....	24
Table 5: Bird Survey Effort for January 2018 Survey Period .....	25
Table 6: Bird Survey Effort for June 2018 Survey Period .....	26
Table 7: Bird Survey Effort for August 2018 Survey Period .....	26
Table 8: Bird Survey Effort for September 2018 Survey Period .....	27
Table 9: Bird Survey Effort for November 2018 Survey Period .....	28
Table 10: Bird Survey Effort for December 2018 Survey Period .....	29



Table 11: ' <i>Species Credit</i> ' threatened bats and their habitats: NSW survey guide for the Biodiversity Assessment Method (OEH, 2018).....	33
Table 12: Herpetofauna Survey Effort for January 2018 Survey Period.....	42
Table 13: Herpetofauna Survey Effort for November 2018 Survey Period.....	42
Table 14: Herpetofauna Survey Effort for December 2018 Survey Period.....	43
Table 15: Summary of Survey Techniques and Effort Used at Each Site within the Study Area.....	45
Table 16: Targeted Searches for Conservation Significant Fauna Species.....	47
Table 17: Threatened Fauna Species Recorded During this Study.....	62
Table 18: Threatened Species Recorded by Others in the Study Area but not Future Ecology.....	94
Table 19: Other Threatened Fauna Species Not Recorded in the Study Area.....	102
Table 20: Threatened Fauna Species Listed under the EPBC Act.....	107

## Plates

- Plate 1 Pink-tailed Legless Lizard at Site 5, November 2018 (Henry Cook)
- Plate 2 Striped Legless Lizard, November-December 2018 (Alex Dudley)
- Plate 3 Spotted Harrier, November 2018 (Alex Dudley)

## Executive Summary

Maxwell Ventures (Management) Pty Ltd, a wholly owned subsidiary of Malabar Coal Limited (Malabar), is seeking consent to develop an underground coal mining operation, referred to as the Maxwell Project (the Project). The Project is in the Upper Hunter Valley of New South Wales (NSW), east-southeast of Denman and south-southwest of Muswellbrook.

This baseline fauna survey report has been prepared by Future Ecology for the Project. This report provides a summary of previous fauna surveys as well as the methods and results of additional fauna surveys undertaken for the Project.

There have been a number of fauna surveys previously undertaken partly within and/or adjacent to the study area since the year 2000. These previous reports provide a good background on the fauna likely to be present in the study area. Additional fauna surveys were completed by Future Ecology in January 2018, June 2018, August 2018, September 2018, November 2018 and December 2018 using a team of up to five ecologists including specialists in birds, reptiles, amphibians and mammals.

A number of sites were surveyed within the study area using a variety of techniques in accordance with relevant NSW and national guidelines. Threatened fauna species listed under the NSW *Biodiversity Conservation Act, 2016* (BC Act) and/or Commonwealth *Environment Protection and Biodiversity Conservation Act, 1999* (EPBC Act) which are known or likely to occur in the study area were specifically targeted during the surveys.

Ten broad fauna habitat types were observed within the study area, comprising three natural habitats (Dry Sclerophyll Forest, Grass Woodlands, Forested Wetlands) and seven secondary habitats (Derived Native Grassland, Planted Trees, Cultivation, Waterbody/Dam, Woodland Rehabilitation, Pasture Rehabilitation and Infrastructure/Cleared Land). Most woodland/forest patches showed evidence of historic and ongoing disturbance from grazing. Most woodland/forest patches were small to medium size (< 150 ha), fragmented and lacked structural diversity in terms of subcanopy and understorey layers due to grazing pressure. Connectivity between woodland/forest patches was generally poor across the study area. However, some fauna habitat features such as hollow bearing trees, hollow logs, fallen timber, were present at most survey sites.

A total of 227 fauna species were recorded in the study area during the surveys namely 8 amphibian, 22 reptile, 148 bird, and, 49 mammal species. A total of 25 threatened fauna species listed under the BC Act (all listed as vulnerable) were recorded by Future Ecology in the study area during the surveys completed in 2018.

Four of the threatened fauna species recorded are considered relevant 'species credit species' under the *Threatened Biodiversity Data Collection* in the study area, namely, Pink-tailed Legless Lizard (also known as the Pink-tailed Worm-lizard) (*Aprasia parapulchella*), Striped Legless Lizard (*Delma impar*), Squirrel Glider (*Petaurus norfolcensis*) and Southern Myotis (*Myotis macropus*).

Five of the threatened fauna species recorded are listed under the EPBC Act, namely, the Pink-tailed Legless Lizard, Striped Legless Lizard, Painted Honeyeater (*Grantiella picta*), Grey-headed Flying-fox (*Pteropus poliocephalus*) and Large-eared Pied Bat (*Chalinolobus dwyeri*). Two additional threatened fauna species listed under the EPBC Act were recorded during surveys undertaken prior to 2018 in the study area, namely, the Swift Parrot (*Lathamus discolor*) and Spotted-tailed Quoll (*Dasyurus maculatus maculatus*) (south-eastern mainland population). The Corben's Long-eared Bat (*Nyctophilus corbeni*) may also have been recorded in the study area nearly 20 years ago but the record is uncertain as the detection method is not known.

# 1 Introduction and Project Description

Maxwell Ventures (Management) Pty Ltd, a wholly owned subsidiary of Malabar Coal Limited (Malabar), is seeking consent to develop an underground coal mining operation, referred to as the Maxwell Project (the Project). The Project is in the Upper Hunter Valley of New South Wales (NSW), east-southeast of Denman and south-southwest of Muswellbrook (Figure 1).

Underground mining is proposed within Exploration Licence (EL) 5460, which was acquired by Malabar in February 2018. Malabar also acquired existing infrastructure within Coal Lease (CL) 229, Mining Lease (ML) 1531 and CL 395, known as the “Maxwell Infrastructure”. The Project would include the use of the substantial existing Maxwell Infrastructure, along with the development of some new infrastructure (Figure 2).

This assessment forms part of an Environmental Impact Statement (EIS) which has been prepared to accompany a Development Application for the Project in accordance with Part 4 of the NSW *Environmental Planning and Assessment Act, 1979* (EP&A Act). Malabar also owns EL 7429, an undeveloped EL called the Spur Hill Underground Coking Coal Project in the Upper Hunter.

The Project would involve an underground mining operation that would produce high quality coals over a period of approximately 26 years. At least 75% of coal produced by the Project would be capable of being used in the making of steel (coking coals). The balance would be export thermal coals suitable for the new generation High Efficiency, Low Emissions power generators.

The Project would involve extraction of run-of-mine (ROM) coal from four seams within the Wittingham Coal Measures using the following underground mining methods:

- underground bord and pillar mining with partial pillar extraction in the Whynot Seam; and
- underground longwall extraction in the Woodlands Hill Seam, Arrowfield Seam and Bowfield Seam.


The substantial existing Maxwell Infrastructure would be used for handling, processing and transportation of coal for the life of the Project. The Maxwell Infrastructure includes an existing coal handling and preparation plant (CHPP), train load-out facilities and other infrastructure and services (including water management infrastructure, administration buildings, workshops and services).

A mine entry area (MEA) would be developed for the Project in a natural valley in the north of EL 5460 to support underground mining and coal handling activities and provide for personnel and materials access.


ROM coal brought to the surface at the MEA would be transported to the Maxwell Infrastructure area. Early ROM coal would be transported via internal roads during the construction and commissioning of a covered overland conveyor system. Subsequently, ROM coal would be transported to the Maxwell Infrastructure area via the covered overland conveyor system.

The existing product coal stockpile area at the Maxwell Infrastructure would be extended to allow for better management of different product coal blends. An additional ROM stockpile would also be developed adjacent to the CHPP to cater for delivery of ROM coal via the covered overland conveyor.

The Project would support continued rehabilitation of previously mined areas and overburden emplacements areas within CL 229, ML 1531 and CL 395. The volume of the East Void would be reduced through the emplacement of reject material generated by Project coal processing activities and would be capped and rehabilitated at the completion of mining.



SHK: 18-03 Maxwell\_ES\_App\_Figure 1\_Rev 0




- LEGEND**
- Mining Operation
  - Proposed Mining Operation
  - Railway
  - Local Government Boundary
  - State Forest
  - National Parks and Wildlife Service Estate
  - Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services and Innovation (2019);  
 Office of Environment and Heritage NSW (2019)

**MAXWELL PROJECT**
  
 Regional Location

**Figure 1**



- LEGEND**
- Railway
  - Exploration Licence Boundary
  - Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Indicative Surface Development Area
  - CHPP Reject Emplacement Area
  - Proposed 66 kV Power Supply
  - Proposed Ausgrid 66 kV Power Supply Extension #
  - Ecology Study Area

# Subject to separate assessment and approval.

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011

MALABAR COAL  
 MAXWELL PROJECT  
 Project General Arrangement



Figure 2

An indicative Project general arrangement is shown on Figure 2. The Project area comprises the following main domains:

- Maxwell Underground – comprising the proposed area of underground mining operations and the MEA within EL 5460.
- Maxwell Infrastructure – the area within existing mining leases comprising the substantial existing infrastructure (including the CHPP) and previous mining areas.
- The transport and services corridor between the Maxwell Underground and Maxwell Infrastructure – this would comprise a site access road, a covered overland conveyor, power supply and other ancillary infrastructure and services.
- A potential realignment of Edderton Road.

A detailed description of the Project is provided in the main document of the EIS.

### 1.1 Purpose of Report

The purpose of the fauna survey and report is to, within the study area:

- survey and document potentially occurring threatened fauna species listed under the NSW *Biodiversity Conservation Act, 2016* (BC Act) in accordance with the relevant survey guidelines;
- survey and document potentially occurring threatened and protected migratory fauna species listed under the Commonwealth *Environment Protection and Biodiversity Conservation Act, 1999* (EPBC Act) in accordance with the relevant survey guidelines;
- survey and document threatened fauna species according to the *Biodiversity Assessment Methodology* (BAM) (Office of Environment and Heritage [OEH], 2017); and
- document the broad fauna habitats and the habitat for relevant 'species credit species'.

### 1.2 Site Description

#### 1.2.1 Study Area

The study area is shown on Figure 2 and is approximately 5,000 hectares (ha) in size. The northern area includes the Maxwell Infrastructure and consists primarily of old open cut workings and infrastructure, with some woodland areas. The southern area consists of a mosaic of cleared grazing land and woodland.

#### 1.2.2 Land Use

Agricultural industries in the surrounding area include cattle grazing, cropping, horse breeding and viticulture. Freehold land in the Project area is owned by Malabar, except for a small area in the northern part of the transport corridor and services corridor and a portion of the Maxwell Infrastructure, which are owned by AGL Energy Limited (AGL).

Land within the Project area is primarily cleared, open paddock grazing land, with some areas of remnant forest and open woodland and mainly used for cattle grazing along with minor cropping.

These agricultural activities are supported by farm dams, unsealed tracks, land contouring, cattle yards and fencing. Land to the north of the Maxwell Underground area is associated with active or previous open cut coal mining activities (i.e. the Mt Arthur Mine).

AGL-owned land associated with the Bayswater and Liddell Power Stations is located to the east of the Project. The Plashett Reservoir serves as an off-river water storage for the Bayswater Power Station along with water supply to the Jerrys Plains township.

The Golden Highway is located to the south and Thomas Mitchell Drive is located to the north of the study area. Edderton Road crosses through the western section of the study area.

### 1.2.3 Regional Setting

The following encompass the study area:

- Hunter Local Land Services Region;
- Sydney Basin Interim Biogeographic Regionalisation for Australia (IBRA) Bioregion and Hunter IBRA sub-region; and
- the Muswellbrook Local Government Area (LGA).

### 1.2.4 Landform and Hydrology

The landform above the underground mining area consists of undulating foothills to moderately-sloping hills over open paddock grazing land, with some areas of remnant forest and open woodland. Surface elevations vary from a low point of approximately 110 metres above Australian Height Datum (mAHD) to a high point of approximately 240 mAHD along a north-east to south-west trending ridgeline.

The Project is located in the Hunter River catchment, with the thalweg of the Hunter River approximately 525 metres (m) south of the underground mining area, at its closest point.

Saddlers Creek, an intermittent stream, is located north of the Maxwell Underground area. Saddlers Creek is a 4<sup>th</sup> order stream to the north of the underground mining area, and a 5<sup>th</sup> order stream downstream of Edderton Road. Saddlers Creek is fed by several small ephemeral creeks and drainage lines that traverse the central and northern portions of the Maxwell Underground area. These creeks and drainage lines form complex drainage networks that comprise the central reaches of the Saddlers Creek catchment area. Dry for much of the year, these watercourses commonly only flow after large rain events.


In the eastern portion of the Maxwell Underground area, another series of ephemeral creeks and drainage lines drain moderate to steeply sloping hills before feeding into Saltwater Creek, a 5<sup>th</sup> order stream immediately upstream of the Hunter River.

### 1.2.5 Vegetation

Hunter Eco (2019) has undertaken flora surveys across the study area. Eleven native vegetation communities were identified and several of these had corresponding 'Derived Natural Grasslands' (DNG) associated with them (Figures 3a and 3b). The majority of the study area comprises White Box – Ironbark – Red Gum shrubby forest (DNG) (approximately 2,200 ha).

### 1.2.6 Summary of Previous Threatened Species Recorded in the Study Area

As detailed in Section 2.1, a literature and database review was undertaken to identify threatened fauna species which are known or likely to occur in the study area. Table 1 lists the threatened fauna species that have previous survey or database records in or close to the study area and/or are predicted to occur in the study area.



- LEGEND**
- Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Dry Sclerophyll Forests (Shrub/grass sub-formation)
  - 1 Red Gum - Ironbark - Apple Shrubby Woodland (PCT1607)
  - 1a Red Gum - Ironbark - Apple Shrubby Woodland - Derived Native Grassland (PCT1607)
  - 2 White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)<sup>1</sup>
  - 2a White Box - Ironbark - Red Gum Shrubby Forest - Derived Native Grassland (PCT1606)<sup>1</sup>
  - Dry Sclerophyll Forests (Shrubby sub-formation)
  - 3 Slaty Box Shrubby Woodland (PCT1655)<sup>2</sup>
  - 3a Slaty Box Shrubby Woodland - Derived Native Grassland (PCT1655)
  - Forested Wetlands
  - 4 Swamp Oak Forest (PCT1731)
  - Grassy Woodlands
  - 6 Bull Oak Grassy Woodland (PCT1692)<sup>3</sup>
  - 7 Yellow Box - Apple Grassy Woodland (PCT1693)<sup>1</sup>
  - 7a Yellow Box - Apple Grassy Woodland - Derived Native Grassland (PCT1693)<sup>1</sup>
  - 8 Fuzzy Box Woodland (PCT201)
  - 8a Fuzzy Box Woodland - Derived Native Grassland (PCT201)
  - 9 Ironbark - Grey Box Grassy Woodland (PCT1691)<sup>4</sup>
  - 9a Ironbark - Grey Box Grassy Woodland - Derived Native Grassland (PCT1691)
  - 10 Weeping Myall Woodland (PCT116)<sup>5</sup>
  - 11 Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)<sup>6</sup>
  - Other**
  - A Planted Trees
  - B Cultivation
  - C Waterbody/Dam
  - D Woodland Rehabilitation
  - E Pasture Rehabilitation
  - F Infrastructure/Cleared Land


<sup>1</sup> Listed BC Act, White Box Yellow Box Blakely's Red Gum Woodland; Listed EPBC Act, White Box - Yellow Box - Blakely's Red Gum Woodland  
<sup>2</sup> Listed BC Act, Hunter Valley Footslopes Slaty Gum Woodland; Listed EPBC Act, Central Hunter Valley Eucalypt Forest and Woodland  
<sup>3</sup> The hatched portions of this PCT is Listed EPBC Act, Central Hunter Valley Eucalypt Forest and Woodland  
<sup>4</sup> Listed BC Act, Central Hunter Grey Box - Ironbark Woodland; Listed EPBC Act, Central Hunter Valley Eucalypt Forest and Woodland  
<sup>5</sup> Listed BC Act, Hunter Valley Weeping Myall Woodland; Listed EPBC Act, Hunter Valley Weeping Myall (Acacia pendula) Woodland  
<sup>6</sup> Listed BC Act, Central Hunter Ironbark - Spotted Gum - Grey Box Forest; Listed EPBC Act, Central Hunter Valley Eucalypt Forest and Woodland

\* Potential ponding impacts associated with subsidence have also been included as part of the surface disturbance area consistent with the predicted impacts in the Geomorphology Assessment prepared by Fluvial Systems (2019).

Source: © NSW Department of Planning and Environment (2018); NSW Department of Finance, Services & Innovation (2018); Orthophoto Mosaic: 2018, 2016, 2011

**Figure 3a**





- LEGEND**
- Biodiversity Assessment Development Footprint
  - Dry Sclerophyll Forests (Shrub/grass sub-formation)
  - Red Gum - Ironbark - Apple Shrubby Woodland (PCT1607)
  - Red Gum - Ironbark - Apple Shrubby Woodland - Derived Native Grassland (PCT1607)
  - White Box - Ironbark - Red Gum Shrubby Forest - Derived Native Grassland (PCT1606)<sup>1</sup>
  - Forested Wetlands
  - 5 Hunter Lowland Red Gum Forest (PCT1598)<sup>7</sup>
  - Grassy Woodlands
  - 11 Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)<sup>6</sup>
  - 11a Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland - Derived Native Grassland (PCT1604)
  - Other
  - C Waterbody/Dam
  - D Woodland Rehabilitation
  - E Pasture Rehabilitation
  - F Infrastructure/Cleared Land

<sup>1</sup> Listed BC Act, White Box Yellow Box Blakely's Red Gum Woodland;  
 Listed EPBC Act, White Box - Yellow Box - Blakely's Red Gum Woodland  
<sup>6</sup> Listed BC Act, Central Hunter Ironbark - Spotted Gum - Grey Box Forest;  
 Listed EPBC Act, Central Hunter Valley Eucalypt Forest and Woodland  
<sup>7</sup> Listed BC Act, Hunter Lowland Redgum Forest

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011

  
**MAXWELL PROJECT**  
**Vegetation Mapping**  
**- Maxwell Infrastructure**



**Figure 3b**

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 1: Threatened Fauna Species Known or Predicted to occur in the Locality

Common Name	Scientific Name	Conservation Status			Database Records				Recorded in Previous Studies <sup>8</sup>
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>	Potentially Associated with PCTs in the Study Area <sup>4</sup>	EPBC Act Protected Matters Search <sup>5</sup>	BioNet Atlas <sup>6</sup>	ALA <sup>7</sup>	
<b>Amphibians</b>									
Green and Golden Bell Frog	<i>Litoria aurea</i>	V	E	S	Yes	Predicted	Yes	Yes	-
Booroolong Frog	<i>Litoria booroolongensis</i>	E	E	S	-	Predicted	-	-	-
Green-thighed Frog	<i>Litoria brevipalmata</i>	-	V	S	Yes	-	-	-	-
<b>Reptiles</b>									
Pink-tailed Legless Lizard	<i>Aprasia parapulchella</i>	V	V	S	-	-	-	-	-
Striped Legless Lizard	<i>Delma impar</i>	V	V	S	Yes	-	-	-	-
Pale-headed Snake	<i>Hoplocephalus bitorquatus</i>	-	V	S	Yes	-	-	-	-
<b>Birds</b>									
Freckled Duck	<i>Stictonetta naevosa</i>	-	V	E	-	-	-	Yes	-
Australasian Bittern	<i>Botaurus poiciloptilus</i>	E	E	E	-	Predicted	-	-	-
Black Falcon	<i>Falco subniger</i>	-	V	E	-	-	-	Yes	-
Square-tailed Kite	<i>Lophoictinia isura</i>	-	V	S/E	Yes	-	Yes	Yes	-
White-bellied Sea-eagle	<i>Haliaeetus leucogaster</i>	Ma	V	S/E	Yes	-	Yes	Yes	-
Spotted Harrier	<i>Circus assimilis</i>	-	V	E	Yes	-	Yes	Yes	A
Red Goshawk	<i>Erythrorhynchus radiatus</i>	V	CE	S	-	Predicted	-	-	-
Little Eagle	<i>Hieraaetus morphnoides</i>	-	V	S/E	Yes	-	Yes	Yes	A, I
Bush Stone-curlew	<i>Burhinus grallarius</i>	-	E	S	Yes	-	-	Yes	-
Australian Painted Snipe	<i>Rostratula australis</i>	E	E	E	-	Predicted	-	-	-
Eastern Curlew	<i>Numenius madagascariensis</i>	CE	-	S/E	-	Predicted	-	-	-
Curlew Sandpiper	<i>Calidris ferruginea</i>	CE	E	S/E	-	Predicted	-	-	-
Glossy Black-Cockatoo	<i>Calyptorhynchus lathami</i>	-	V	S/E	Yes	-	-	-	-
Gang-gang Cockatoo	<i>Callocephalon fimbriatum</i>	-	V	S/E	Yes	-	-	Yes	-
Little Lorikeet	<i>Glossopsitta pusilla</i>	-	V	E	Yes	-	Yes	Yes	J
Turquoise Parrot	<i>Neophema pulchella</i>	-	V	E	Yes	-	-	Yes	-
Swift Parrot	<i>Lathamus discolor</i>	CE	E	S/E	Yes	Predicted	-	-	A
Eastern Grass Owl	<i>Tyto longimembris</i>	-	V	E	Yes	-	-	-	-
Masked Owl	<i>Tyto novaehollandiae</i>	-	V	S/E	Yes	-	-	-	-
Powerful Owl	<i>Ninox strenua</i>	-	V	S/E	Yes	-	Yes	Yes	-

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 1 (Continued): Threatened Fauna Species Known or Predicted to occur in the Locality

Common Name	Scientific Name	Conservation Status			Database Records				Recorded in Previous Studies <sup>8</sup>
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>	Potentially Associated with PCTs in the Study Area <sup>4</sup>	EPBC Act Protected Matters Search <sup>5</sup>	BioNet Atlas <sup>6</sup>	ALA <sup>7</sup>	
Barking Owl	<i>Ninox connivens</i>	-	V	S/E	Yes	-	Yes	Yes	B
Brown Treecreeper (eastern subspecies)	<i>Climacteris picumnus victoriae</i>	-	V	E	Yes	-	Yes	Yes	A
Speckled Warbler	<i>Chthonicola sagittata</i>	-	V	E	Yes	-	Yes	Yes	A
Black-chinned Honeyeater (eastern subspecies)	<i>Meliphreptus gularis gularis</i>	-	V	E	Yes	-	Yes	-	
Regent Honeyeater	<i>Anthochaera phrygia</i>	CE	CE	S/E	Yes	Predicted	-	-	-
Painted Honeyeater	<i>Grantiella picta</i>	V	V	E	Yes	Predicted	-	-	-
Hooded Robin (south-eastern form)	<i>Melanodryas cucullata cucullata</i>	-	V	E	Yes	-	Yes	-	
Flame Robin	<i>Petroica phoenicea</i>	-	V	E	Yes	-	-	Yes	-
Scarlet Robin	<i>Petroica boodang</i>	-	V	E	Yes	-	Yes	-	A
Grey-crowned Babbler (eastern subspecies)	<i>Pomatostomus temporalis temporalis</i>	-	V	E	Yes	-	Yes	-	A
Varied Sittella	<i>Daphoenositta chrysoptera</i>	-	V	E	Yes	-	Yes	Yes	C
Dusky Woodswallow	<i>Artamus cyanopterus cyanopterus</i>	-	V	E	Yes	-	Yes	Yes	-
Diamond Firetail	<i>Stagonopleura guttata</i>	-	V	E	Yes	-	Yes	Yes	A, B, J
<b>Mammals</b>									
Spotted-tailed Quoll	<i>Dasyurus maculatus maculatus</i> (south-eastern mainland population)	E	V	E	Yes	Predicted	Yes	Yes	D, E
Brush-tailed Phascogale	<i>Phascogale tapoatafa</i>	-	V	S	Yes	-	Yes	-	-
Common Planigale	<i>Planigale maculata</i>	-	V	S	Yes	-	-	-	-
Koala	<i>Phascolarctos cinereus</i>	V	V	S/E	Yes	Predicted	Yes	-	-
Eastern Pygmy-possum	<i>Cercartetus nanus</i>	-	V	S	Yes	-	-	-	-
Yellow-bellied Glider	<i>Petaurus australis</i>	-	V	E	Yes	-	-	-	-
Squirrel Glider	<i>Petaurus norfolcensis</i>	-	V	S	Yes	-	Yes	-	A, D, E, F, J
Greater Glider	<i>Petauroides volans</i>	V	-	S	-	Predicted	-	-	-
Brush-tailed Rock-wallaby	<i>Petrogale penicillata</i>	V	E	S	Yes	Predicted	Yes	-	-
Grey-headed Flying-fox	<i>Pteropus poliocephalus</i>	V	V	S/E	Yes	Predicted	Yes	-	J

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 1 (Continued): Threatened Fauna Species Known or Predicted to occur in the Locality

Common Name	Scientific Name	Conservation Status			Database Records				Recorded in Previous Studies <sup>8</sup>
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>	Potentially Associated with PCTs in the Study Area <sup>4</sup>	EPBC Act Protected Matters Search <sup>5</sup>	BioNet Atlas <sup>6</sup>	ALA <sup>7</sup>	
Yellow-bellied Sheath-tail-bat	<i>Saccolaimus flaviventris</i>	-	V	E	Yes	-	Yes	-	A, J
Eastern Freetail-bat	<i>Mormopterus norfolkensis</i>	-	V	E	Yes	-	Yes	-	A, B, C, E, G, J
Northern Freetail-bat	<i>Mormopterus lumsdenae</i>	-	V	E	No	-	-	-	G
Little Bentwing-bat	<i>Miniopterus australis</i>	-	V	S/E	Yes	-	Yes	-	G
Eastern Bentwing-bat	<i>Miniopterus schreibersii oceanensis</i>	-	V	S/E	Yes	-	Yes	-	A, C, D, E, F, G, H, J
Corben's Long-eared Bat	<i>Nyctophilus corbeni</i>	V	V	E	Yes	Predicted	Yes	-	B
Large-eared Pied Bat	<i>Chalinolobus dwyeri</i>	V	V	S	Yes	Predicted	Yes	-	A, C, G
Eastern False Pipistrelle	<i>Falsistrellus tasmaniensis</i>	-	V	E	Yes	-	Yes	-	E, F
Southern Myotis	<i>Myotis macropus</i>	-	V	S	Yes	-	Yes	-	A, B, G
Greater Broad-nosed Bat	<i>Scoteanax rueppellii</i>	-	V	E	Yes	-	Yes	-	B, D, E, J
Eastern Cave Bat	<i>Vespadelus troungtoni</i>	-	V	S	Yes	-	Yes	-	A, G, J
New Holland Mouse	<i>Pseudomys novaehollandiae</i>	V	-	E	-	Predicted	-	-	-

Highlighted species are species recorded in the study area.

<sup>1</sup> Conservation status under the EPBC Act (current as at March 2019). V = Vulnerable, E = Endangered, CE = Critically Endangered, Ma = Marine.

<sup>2</sup> Conservation status under the BC Act (current as at March 2019). V = Vulnerable, E = Endangered, CE = Critically Endangered.

<sup>3</sup> Biodiversity credit class under the *Threatened Biodiversity Data Collection* (OEH, 2019a) (current as at March 2019), S = Species, E = Ecosystem.

<sup>4</sup> OEH (2019a).

<sup>5</sup> Department of the Environment and Energy (2018a).

<sup>6</sup> OEH (2019b).

<sup>7</sup> Atlas of Living Australia (2018).

<sup>8</sup> A – Cumberland Ecology (2009a) and/or Cumberland Ecology (2012)

B – Ecotone (2000).

C – Eco Logical Australia (2015).

D – Eco Logical Australia (2016a).

E – Eco Logical Australia (2016b).

F – Eco Logical Australia (2014).

G – Eco Logical Australia (2017).

H – Umwelt Environmental Consultants (Umwelt) (2006b).

I – Umwelt (2007b).

J – Hansen Bailey (2007).

PCT = Plant Community Type.

## 2 Methods

### 2.1 Literature and Database Review

A literature and database review was undertaken prior to undertaking the field surveys (Section 2.3) to identify known or potentially occurring threatened fauna species or their habitats.

The following databases were reviewed:

- *Birdlife Australia Atlas Database* (Birdlife Australia, 2018);
- *BioNet Atlas* (OEH, 2019b);
- *Protected Matters Search Tool* (Department of the Environment and Energy [DEE], 2018a); and
- *Atlas of Living Australia* (Atlas of Living Australia [ALA], 2018).

The following mapping sources were reviewed:

- *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019);
- *SIX Maps* (NSW Spatial Services, 2018); and
- *Google Earth Pro* (Google, 2018).

The following local survey reports were also reviewed:

- *Ecological Assessment – Proposed South Pit Extension Project* (Umwelt, 2006b).
- *Ecological Assessment – Proposed Mt Arthur Underground Project* (Umwelt, 2007b).
- *Drayton Mine Extension Flora and Fauna Impact Assessment* (Hansen Bailey, 2007).
- *Ecological Assessment of Section 75W Modification for Drayton Mine* (Cumberland Ecology, 2009a).
- *Mt Arthur Coal Consolidation Project Ecological Assessment* (Cumberland Ecology, 2009b).
- *Mt Arthur Coal Open Cut Modification – Ecological Assessment* (Hunter Eco, 2013).
- *Mt Arthur Coal – Fauna Survey Report* (Niche Environment and Heritage, 2012).
- *Drayton South Coal Project Ecology Impact Assessment* (Cumberland Ecology, 2012).
- *Drayton South Coal Project Biodiversity Assessment Report* (Cumberland Ecology, 2015a).
- *Drayton South Coal Project Biodiversity Offset Strategy* (Cumberland Ecology, 2015b).
- 2013-2017 Spring Biodiversity Monitoring Reports of the former Drayton Mine (Eco Logical Australia, 2014-2017).

Since 2000, several surveys have been undertaken for surrounding coal projects. Results for each of the surveys are summarised below.

### **Saddlers Creek Survey**

Ecotone (2000 in Cumberland Ecology, 2012) undertook flora and fauna surveys of Saddlers Creek in February 2000. The fauna survey included harp trapping, spotlighting, call playback, hair tube, Anabat and tripline bat surveys and a bird census. During the survey the following threatened species were recorded: the Barking Owl (*Ninox connivens*), Speckled Warbler (*Chthonicola sagittata*), Brown Treecreeper (eastern subspecies) (*Climacteris picumnus victoriae*), Black-chinned Honeyeater (eastern subspecies) (*Melithreptus gularis gularis*), Hooded Robin (south-eastern form) (*Melanodryas cucullata cucullata*), Grey-crowned Babbler (eastern subspecies) (*Pomatostomus temporalis temporalis*), Diamond Firetail (*Stagonopleura guttata*), Koala (*Phascolarctos cinereus*), Eastern Freetail-bat (*Mormopterus norfolkensis*), Eastern Bentwing-bat (*Miniopterus schreibersii oceanensis*), Corben's Long-eared Bat (*Nyctophilus corbeni*), Southern Myotis (*Myotis macropus*) and Greater Broad-nosed Bat (*Scoteanax rueppellii*) (Table 1). Two additional species which are not listed as threatened in NSW but are listed as nationally protected migratory species under the EPBC Act were also detected, namely the White-throated Needletail (*Hirundapus caudacutus*) and Rainbow Bee-eater (*Merops ornatus*).

### **Mt Arthur Mine**

Umwelt (2006b and 2007b) conducted various fauna monitoring surveys of the Mt Arthur Mine and surrounds between 2004 and 2006. Survey techniques involved hair traps, Elliott traps, cage traps, spotlight and diurnal surveys, Anabat surveys and call playback.

Between 21-25 February 2005, a fauna monitoring survey of the Mt Arthur Mine and the area located to the south-east; near Saddlers Creek and adjacent to the Maxwell Infrastructure, was undertaken. Threatened species recorded during this survey include the Squirrel Glider (*Petaurus norfolkensis*), Eastern Bentwing-bat, Eastern Cave Bat (*Vespadelus troughtoni*) and Southern Myotis (Umwelt, 2006b).

In December 2005, Umwelt (2006a in Hunter Eco, 2013) undertook a monitoring fauna survey of McLeans Hill, Saddlers Creek and Mt Arthur Mine and surrounds. Threatened species recorded include the Speckled Warbler, Grey-crowned Babbler (eastern subspecies), Varied Sittella (*Daphoenositta chrysoptera*), Squirrel Glider, Eastern Freetail-bat, Eastern Bentwing-bat and Southern Myotis.

On 7-11 March and 5-7 December 2005, Umwelt (2007b) conducted surveys for the Mt Arthur Underground Project, in areas located to the south and south-west of the Mt Arthur open cut mining areas, including near Saddlers Creek. Survey techniques included trapping (Elliott traps, cage traps, hair funnels and tubes and harp traps), spotlight surveys, diurnal surveys and Anabat surveys. Threatened species recorded included the Little Eagle (*Hieraaetus morphnoides*), Grey-crowned Babbler (eastern subspecies), Spotted Harrier (*Circus assimilis*), Speckled Warbler, Squirrel Glider, Eastern Bentwing-bat, Eastern Cave Bat, Greater Broad-nosed Bat, Eastern Freetail-bat and Southern Myotis (Umwelt, 2007b). Commonwealth listed migratory species recorded during the survey include the Black-shouldered Kite (*Elanus axillaris*), Wedge-tailed Eagle (*Aquila audax*), Nankeen Kestrel (*Falco cenchroides*), Masked Lapwing (*Vanellus miles*), White-throated Needletail and Rainbow Bee-eater (Umwelt, 2007b).

Umwelt (2007a in Hunter Eco, 2013) undertook a survey in December 2006 of the Mt Arthur Mine and McLeans Hill. Umwelt (2007a in Hunter Eco, 2013) recorded the Speckled Warbler, Eastern Freetail-bat, Large-eared Pied Bat (*Chalinolobus dwyeri*) and Greater Broad-nosed Bat.

The Spotted-tailed Quoll (south-eastern mainland population) (*Dasyurus maculatus maculatus*) was tentatively recorded during the first half of 2006 by a Hunter Valley Energy Coal Pty Ltd (HVEC) staff member on the main access road to the Mt Arthur Mine offices, near the intersection with Thomas Mitchell Drive (Hunter Eco, 2013).

Cumberland Ecology (2010 in Hunter Eco, 2013) conducted flora and fauna surveys of the Mt Arthur Mine surrounds between 20-23 September 2010. During the monitoring survey the Squirrel Glider and Eastern Bentwing-bat were recorded.

In 2012, a Koala was recorded within the Mt Arthur Mine, to the south-west of the Thomas Mitchell Drive Offset Area. The Koala was taken by wildlife carers who relocated him into a rehabilitated area, near where he was originally found (HVEC pers. comm., 2012 in Hunter Eco, 2013).

Niche Environment and Heritage (2012) conducted fauna surveys on 1 May and 7-11 May 2012, for areas associated with expansion of open cut coal mining activities at the Mt Arthur Mine site and adjacent to the Maxwell Infrastructure, but outside the study area.

Survey methods included arboreal Elliott traps, infra-red camera traps, hair tubes, ultrasonic call recording, diurnal bird surveys, spotlight surveys, call playback, stag watching, koala scat searches, herpetological surveys and frog chorus surveys. Two threatened species were recorded during the survey; the Grey-headed Flying-fox and Eastern Freetail-bat (Table 1) (Niche, 2012). The Eastern Bentwing-bat, Eastern Cave Bat and Eastern False Pipistrelle may have been recorded, however call recordings from these species were not of sufficient quality to be certain. The White-bellied Sea-eagle (*Haliaeetus leucogaster*), listed as migratory under the EPBC Act, was recorded near the Mt Arthur Mine during the flora surveys (Hunter Eco, 2013).

#### **Former Drayton South Coal Project**

Cumberland Ecology (2012 and 2015a) conducted several fauna surveys within the current study area for the former Drayton South Coal Project. Surveys were undertaken from 30 September-2 October, 26-28 October and 2-3 November in 2009 and from 14-18 March, 2-3 May and 16-24 June in 2011. Survey techniques included hair tubes, Anabat surveys, bird census, spotlight surveys, call playback, Elliott traps, cage traps, infra-red cameras, harp traps, targeted surveys and diurnal surveys.

The following threatened species were recorded during the 2009 and 2011 surveys: Spotted Harrier, Little Eagle, Swift Parrot (*Lathamus discolor*), Brown Treecreeper (eastern subspecies), Speckled Warbler, Grey-crowned Babbler (eastern subspecies), Scarlet Robin (*Petroica boodang*), Diamond Firetail, Yellow-bellied Sheath-tail-bat (*Saccolaimus flaviventris*), Eastern Freetail-bat, Large-eared Pied Bat, Eastern Bentwing-bat, Southern Myotis (non-definite call identification), and Eastern Cave Bat (non-definite call identification) (Table 1), The Commonwealth listed migratory species, the Rainbow Bee-eater, was also recorded.

#### **Maxwell Infrastructure**

Hansen Bailey (2007) conducted fauna surveys of the Maxwell Infrastructure and surrounds for the periods over 14-17 February 2006, 6 September 2006 and 12-16 February 2007. Survey methods included arboreal mammal trapping, arboreal hair tube sampling, spotlighting, call playback, Anabat surveys, avian fauna surveys and opportunistic sightings

Hansen Bailey (2007) recorded the Little Lorikeet (*Glossopsitta pusilla*) (no location given), Diamond Firetail (north of rail loop, north-west of Site 17), Squirrel Glider (between the rail loop and coal stockpiles [equivalent to Site 17]), Grey-headed Flying-fox (flying near the Access Road Dam), Yellow-bellied Sheath-tail-bat (non-definite call identification south-west to Site 17), Eastern Freetail-bat (equivalent to Site 17), Eastern Bentwing-bat (equivalent to Site 17), Greater Broad-nosed Bat (non-definite call identification at what is equivalent to Site 16b), and the Eastern Cave Bat (non-definite call identification at equivalent to Sites 16 and 17). The Commonwealth listed migratory species White-throated Needletail and Rainbow Bee-eater were also recorded near the rail loop and south-west of the Access Road Dam, respectively.

Cumberland Ecology (2009a) conducted a site inspection and fauna habitat assessment of the Maxwell Infrastructure in May 2009, During the field surveys call playback surveys were undertaken but no targeted threatened fauna surveys. One threatened species, the Speckled Warbler, was recorded outside the Maxwell Infrastructure, in the Northern Offset.

Eco Logical Australia (2014 to 2017) undertook annual flora and fauna monitoring surveys between 2013 and 2017 of the rehabilitation areas within the Maxwell Infrastructure and surrounds. All surveys were conducted during the spring season, with eight permanent fauna survey plots established and monitored annually for signs of fauna activity. Survey techniques consisted hair tube traps (arboreal and terrestrial) and remote cameras, spotlight surveys, bird census, call playback, herpetological and anabat surveys. Threatened species recorded during the monitoring period included the Squirrel Glider, Varied Sittella, Speckled Warbler, Little Lorikeet, Little Eagle, Grey-crowned Babbler (eastern subspecies), Spotted-tail Quoll (south-eastern mainland population), Brush-tailed Phascogale (*Phascogale tapoatafa*), Eastern Bentwing-bat, Little Bentwing-bat (*Miniopterus australis*), Eastern False Pipistrelle (*Falsistrellus tasmaniensis*), Large-eared Pied Bat, Eastern Freetail Bat, Greater Broad-nosed Bat, Yellow-bellied Sheathtail-bat, Eastern Cave Bat, Southern Myotis, and Northern Freetail-bat (*Mormopterus lumsdenae*) (Table 1). The following Commonwealth listed migratory species were also recorded in during the monitoring period: Satin Flycatcher (*Myiagra cyanoleuca*), White-throated Needle-tail and Rainbow Bee-eater.

## 2.2 Relevant Survey Guidelines

Relevant guidelines that were followed during fauna surveys are as follows:

- *Threatened Biodiversity Survey and Assessment: Guidelines for Developments and Activities (Working Draft)* (Department of Environment and Conservation [DEC], 2004a).
- *Hygiene Protocol for The Control of Disease in Frogs*. (Department of Environment and Climate Change [DECC], 2008a).
- *Threatened Species Survey and Assessment Guidelines: Field Survey Methods for Fauna – Amphibians* (DECC, 2009).
- *'Species Credit' Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018).
- *Survey Guidelines for Australia's Threatened Frogs* (Department of Environment, Water, Heritage and Arts [DEWHA], 2010a).
- *Survey Guidelines for Australia's Threatened Bats* (DEWHA, 2010b).
- *Survey Guidelines for Australia's Threatened Birds* (DEWHA, 2010c).
- *Survey Guidelines for Australia's Threatened Mammals* (Department of Sustainability, Environment, Water, Population and Communities [SEWPaC], 2011a).
- *Survey Guidelines for Australia's Threatened Reptiles* (SEWPaC, 2011b).
- *EPBC Act Referral Guidelines for the Vulnerable Striped Legless Lizard, Delma impar* (SEWPaC, 2011c).
- *EPBC Act Referral Guidelines for the Vulnerable Koala* (Department of the Environment, 2014).
- SPRAT profiles of relevant Commonwealth listed threatened and/or migratory fauna species (DEE, 2018b).



## 2.3 Field Survey

### 2.3.1 Weather, Climate and Astronomical Conditions

Fauna surveys took place over several separate periods:

1. 22 to 28 January 2018;
2. 4 to 7 June 2018;
3. 28 to 30 August 2018;
4. 17 to 20 September 2018;
5. 12 to 16 November 2018;
6. 19 to 23 November 2018;
7. 3 to 7 December 2018; and
8. 17 to 21 December 2018.

Weather records during the surveys were taken from the Maxwell Infrastructure CHPP Automatic Weather Station (AWS), and closest operating Bureau of Meteorology (BoM) AWS at Singleton Sewage Treatment Plant (STP) (station 061397) (BoM, 2018a), approximately 28 kilometres (km) south-east of the study area. Astronomical records were taken from the Geoscience Australia website (2018a, 2018b), the Museum of Applied Arts and Sciences – Sydney Observatory website (2018) and the United States Naval Observatory website (2019).

Since April 2017, there has been serious to severe rainfall deficiencies across large areas of NSW including the study area (BoM, 2018b). Below average rainfall conditions continued into Spring 2018 with only 81% of the average spring rainfall recorded at Singleton STP (BoM, 2018c). Rainfall for 2018 as a whole was exceptionally low over the south eastern quarter of the mainland, with much of the region experiencing totals in the lowest 10% of records. As of January 2019, significant rainfall deficiencies continued to affect large areas of eastern Australia at timescales out to around two years' duration (BoM, 2019a).

In terms of temperature, 2018 was Australia's third-warmest year on record (BoM, 2019b). At Singleton STP the mean maximum temperature for Spring 2018 was 0.3°C below the average and the mean minimum temperature was 2.2°C above the average (BoM, 2018c).

Weather conditions during the January survey period were very hot, with a maximum temperature of 40.8°C recorded and each survey date reaching over 32.5°C. Minimum nightly temperatures were also warm (the minimum recorded temperature being 19.5°C). There was 31.2 millimetres (mm) of rainfall recorded during the survey period from 25 to 28 January 2018.

Weather conditions during the June survey period were cool to mild with temperatures ranging from 9.3°C to 18.2°C. Some very minor rainfall (2 mm) was recorded on each of the last three days of survey, and an additional 4.4 mm of rainfall recorded in the two days prior to the survey period from 2 to 3 June 2018.

The August survey period had very cold frosty mornings and mild days with temperatures ranging from 1.4°C to 18.4°C. Some very minor rainfall (0.2 mm) was recorded on the first day of the survey period, and an additional 11.2 mm of rainfall recorded in the two days prior to the survey period from 26 to 27 August 2018.

Weather conditions during the September survey period were cool mornings and mild to hot days with temperatures ranging from 3.7°C to 27.9°C. No rainfall was recorded during the survey period.

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Weather conditions during the November survey periods were warm to hot, with temperatures ranging from 12.5°C to 33.0°C. There was 9.6 mm of rainfall recorded during the last two days of the first survey period, and an additional 17 mm of rainfall recorded four to five days prior to the survey from 7 to 8 November 2018. During the second survey period 1.2 mm of rainfall was recorded on 21 November 2018.

Weather conditions during the December survey periods were warm to hot, with temperatures ranging from 13.6°C to 36.5°C. There was no rainfall recorded in the first survey period, however there was 36.4 mm of rainfall recorded four to six days prior to this survey from 27 to 29 November 2018. During the second survey period 5.2 mm of rainfall was from 17 to 21 December 2018, and an additional 77.2 mm of rainfall recorded in the week prior to the survey from 10 to 16 December 2018.

A summary of these weather records in addition to astronomical records relevant to the survey periods are presented in Table 2.

**Table 2: Weather and Astronomical Records during Survey Periods**

Date	Minimum temperature (°C)	Maximum temperature (°C)	Rainfall (mm)	Sunrise	Sunset	Moonrise	Moonset	Moonphase
<u>January 2018</u>								
22/01/2018	19.5	40.8	0	0511	1905	0932	2212	Waxing Crescent
23/01/2018	22.5	36.8	0	-	-	-	-	-
24/01/2018	21.2	37.3	0	-	-	-	-	-
25/01/2018	21.0	33.5	0.2	-	-	-	-	First Quarter
26/01/2018	22.2	34.0	0.2	-	-	-	-	-
27/01/2018	22.5	33.7	18.4	-	-	-	-	-
28/01/2018	22.6	32.5	12.4	0517	1902	1543	0130	Waxing Gibbous
<u>June 2018</u>								
4/06/2018	9.3	17.1	0	0651	1658	2148	1056	Waning Gibbous
5/06/2018	10.9	15.4	0.4	-	-	-	-	-
6/06/2018	10.7	12.9	1.4	-	-	-	-	-
7/06/2018	10.2	18.2	0.2	0653	1658	-	1242	Third Quarter
<u>August 2018</u>								
28/08/2018	6.7	16.3	0.2	0619	1737	1910	0720	Full Moon
29/08/2018	2.4	18.4	0	-	-	-	-	-
30/08/2018	1.4	18.1	0	0616	1738	2103	0824	Waning Gibbous
<u>September 2018</u>								
17/09/2018	3.7	19.8	0	0553	1749	1058	0021	First Quarter
18/09/2018	4.5	25.9	0	-	-	-	-	-
19/09/2018	14.4	27.9	0	-	-	-	-	-
20/09/2018	8.6	18.3	0	0549	1751	1325	0249	Waxing Gibbous
<u>November 2018</u>								
12/11/2018	12.5	28.1	0	0549	1931	0915	2338	Waxing Crescent
13/11/2018	14.1	29.8	0	-	-	-	-	-
14/11/2018	16.4	26.3	0	-	-	-	-	-

**Table 2 (Continued): Weather and Astronomical Records during Survey Periods**

Date	Minimum temperature (°C)	Maximum temperature (°C)	Rainfall (mm)	Sunrise	Sunset	Moonrise	Moonset	Moonphase
15/11/2018	16.1	31.3	8.6	-	-	1151	0105	First Quarter
16/11/2018	14.0	20.3	1	-	-	-	-	-
19/11/2018	13.5	26.8	0	-	-	-	-	-
20/11/2018	13.4	33.0	0	-	-	1630	0356	Waxing Gibbous
21/11/2018	19.3	28.3	1.2	-	-	-	-	-
22/11/2018	18.7	25.6	0	-	-	-	-	-
23/11/2018	14.5	23.3	0	0544	1941	1937	0544	Full Moon
<b>December 2018</b>								
3/12/2018	15.0	31.9	0	0542	1950	0310	1545	Waning Crescent
4/12/2018	16.7	30.4	0	-	-	-	-	-
5/12/2018	17.2	24.5	0	-	-	-	-	-
6/12/2018	15.2	28.5	0	-	-	-	-	-
7/12/2018	13.6	29.9	0	-	-	0537	1943	New Moon
17/12/2018	20.8	33.7	0.2	-	-	1414	0154	Waxing Gibbous
18/12/2018	20.8	30.1	0	-	-	-	-	-
19/12/2018	21.6	31.2	4	-	-	-	-	-
20/12/2018	20.4	36.5	1	-	-	-	-	-
21/12/2018	19.3	26.3	0.2	0546	2003	1821	0417	Waxing Gibbous
(22/12/2018)	-	-	-	-	-	-	-	(Full Moon)


Sources: BOM (2018a), Geoscience Australia (2018a, 2018b), Sydney Observatory (2018), Maxwell Infrastructure CHPP AWS.

### 2.3.2 Techniques

#### **Stratification of the study area and site selection**

The study area was initially assessed through interpretation of digital aerial imagery and from literature generated from previous studies. The landscape is mostly cleared agricultural lands and therefore remnant patches of treed vegetation within the study area were used as a basis for the initial stratification. Further stratification considered previous threatened and/or protected migratory fauna records within the study area and the spacing of survey sites.

General fauna survey sites are listed in Table 3 and shown on Figure 4. Bat survey sites are shown on Figure 5 and amphibian survey sites are shown on Figure 6. The previous survey sites by Ecotone (2000), Cumberland Ecology in 2011 (2012) and Eco Logical Australia (2017) are also shown on Figures 4 and 5.




- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
  - Future Ecology Sites**
  - + General Fauna Survey Site
  - Mammal Survey Site
  - ▼ Pitfall Trap
  - Artificial Habitat (Tiles)
  - + Eco Logical Australia (2017) Sites
  - + Monitoring Location
  - + Cumberland Ecology (2012) Sites
  - ◇ Bird Census
  - ☆ Cage Trap
  - ▽ Infrared Camera
  - + Ecotone (2000) Sites
  - Call Playback

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011




**Figure 4**



- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
  - Waterbody/Dam
  - Future Ecology Sites
  - ▲ Harp Trap
  - Acoustic Detector
  - Habitat Inspection
  - ▲ Cumberland Ecology (2012) Sites
  - Acoustic Detector
  - ▲ Ecotone (2000) Sites
  - Harp Trap
  - ▲ Acoustic Detector
  - Harp Trap
  - Acoustic Detector

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011

**Figure 5**



- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
  - Waterbody/Dam
  - ◆ Amphibian Survey Site

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011

  
**MAXWELL PROJECT**  
 Amphibian Survey Sites

**Figure 6**

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 3: Fauna Survey Sites for the Study Area

Site	Location (Lat/Long GDA)		Site Type
1	-32.386	150.891	General Fauna Survey Site
1a	-32.386	150.888	Mammal Survey Site
1b	-32.388	150.891	Mammal Survey Site
2	-32.396	150.892	General Fauna Survey Site
2a	-32.396	150.890	Mammal Survey Site
2b/3b^	-32.399	150.895	Mammal Survey Site
2c	-32.396	150.894	Mammal Survey Site
2d	-32.396	150.897	Pitfall Trap
3	-32.404	150.890	General Fauna Survey Site
3a	-32.401	150.889	Mammal Survey Site and Pitfall Trap
3c	-32.403	150.896	Mammal Survey Site
4	-32.402	150.883	General Fauna Survey Site
4a	-32.402	150.883	Mammal Survey Site and Artificial Habitat (Tiles)
5	-32.414	150.867	General Fauna Survey Site
5a	-32.419	150.873	Mammal Survey Site
5b	-32.415	150.870	Mammal Survey Site and Pitfall Trap
6	-32.413	150.852	General Fauna Survey Site
6a	-32.411	150.851	Mammal Survey Site
6b	-32.413	150.851	Mammal Survey Site and Artificial Habitat (Tiles)
6c	-32.410	150.856	Mammal Survey Site
7	-32.417	150.839	General Fauna Survey Site
7a	-32.415	150.843	Mammal Survey Site and Artificial Habitat (Tiles)
7b	-32.420	150.840	Mammal Survey Site
7c	-32.417	150.843	Mammal Survey Site
8	-32.410	150.826	General Fauna Survey Site
8a	-32.410	150.826	Mammal Survey Site
9	-32.440	150.829	General Fauna Survey Site
10	-32.438	150.852	General Fauna Survey Site
10a	-32.444	150.857	Mammal Survey Site
10b	-32.445	150.854	Mammal Survey Site
11	-32.423	150.879	General Fauna Survey Site
11a	-32.425	150.878	Mammal Survey Site
12	-32.430	150.871	General Fauna Survey Site
13	-32.414	150.821	General Fauna Survey Site
14	-32.429	150.824	General Fauna Survey Site
14a	-32.425	150.822	Mammal Survey Site
14b	-32.419	150.823	Mammal Survey Site
15	-32.380	150.931	General Fauna Survey Site
15a	-32.378	150.926	Mammal Survey Site
16	-32.335	150.935	General Fauna Survey Site
16a	-32.337	150.936	Mammal Survey Site
16b	-32.335	150.933	Mammal Survey Site
16c	-32.337	150.937	Mammal Survey Site
17	-32.336	150.924	General Fauna Survey Site, Mammal Survey Site and Artificial Habitat (Tiles)
18	-32.414	150.843	General Fauna Survey Site
18a	-32.406	150.850	Mammal Survey Site

**Table 3 (Continued): Fauna Survey Sites for the Study Area**

Site	Location (Lat/Long GDA)		Site Type
IncA*	-32.417	150.857	Mammal Survey Site
IncB*	-32.442	150.835	Mammal Survey Site
Savoy Dam	-32.380	150.901	Mammal Survey Site

<sup>^</sup> Site 2b/3b was a call-playback survey site located close to the boundary of Sites 2 and 3 and therefore covered both sites.

\* Sites IncA and IncB were incidental mammal survey sites.

Eleven survey sites were initially selected for the January 2018 survey period with a further three sites added to the June 2018 survey period to cover additional areas including west of Edderton Road. Some of the survey sites (Sites 2, 3, 4, 5, 6, 7, 9, 10, 11, 12 and 14) had been previously surveyed (or partly surveyed) by Ecotone (2000) and/or Cumberland Ecology (2012). The study area was extended to the north to cover the Maxwell Infrastructure and an additional two sites (Sites 15 and 16) added to this area and initially surveyed in August 2018. The two site locations were selected as they were the largest areas of remaining treed habitat within CL 229. Site 17 was added in October 2018 adjacent to the existing coal stockpile area within CL 229, to cover some additional proposed surface development.

A mine dam (known as Savoy Dam) in the southern part of CL 229 was also added in October 2018. Site 18 was also added at the same time to cover a plant community type (PCT) identified above the Maxwell Underground.

The following survey techniques were undertaken at “General Fauna Survey Sites” listed in Table 3: general diurnal and nocturnal bird, reptile, amphibian and reptile surveys using a variety of standard techniques including observation, listening, spotlighting, call-playback and habitat searches,

The following survey techniques were undertaken at “Mammal Survey Sites” listed in Table 3: live trapping for arboreal and terrestrial mammals using Elliott traps and cages together with the use of hair tubes/funnels, nest boxes and wildlife cameras for longer term monitoring. Insectivorous bats were sampled using harp traps and acoustic devices.

Artificial habitat (second-hand terracotta roofing tiles) were placed at “Artificial Habitat (Tiles) Sites” listed in Table 3. Pitfall traps were placed at the “Pitfall Trap Sites” listed in Table 3.

### **Field Surveys**

It should be noted that some surveys were often done concurrently e.g. spotlighting surveys were carried out at the same time as nocturnal bird and herpetofauna surveys. Given that there was a team of five ecologists in January 2018, two ecologists in June, August and September 2018, and five ecologists in November and December 2018, total survey effort would be in the order of at least two times what is described below for the specialist avifauna and herpetofauna surveys.



The below sub-sections describe the following survey techniques:

- habitat surveys;
- diurnal and nocturnal bird surveys;
- ground Elliott trapping;
- arboreal Elliott trapping;
- cage trapping;
- hair tubes;
- camera trapping;
- nest boxes;
- bat surveys;
- harp trapping;
- ultrasonic bat detection;
- microbat habitat searches;
- nocturnal call playback;
- spotlighting;
- Koala scat searches;
- searches for reptiles and amphibians (active searches, pitfall traps and artificial shelter habitat);
- tadpole surveys; and
- opportunistic observations.

### **Habitat Surveys**

Fauna habitat searches were conducted for potential foraging, roosting, breeding or nesting habitat of nocturnal and diurnal species. This includes inspection for the presence of tree hollows, stags, bird nests, possum dreys, decorticating bark, rock shelters, rock outcrops/crevices, mature/old growth trees, food trees (*Banksia spp.*, *Allocasuarina spp.*, and winter-flowering eucalypts), culverts, dens, dams, riparian areas and refuge habitats within man-made structures.

The quality of the fauna habitat was assessed and categorised (low, medium or high) by the presence or absence of components of the ecosystems used by different fauna groups, e.g. large hollow-bearing trees for hollow dependent species, presence of understorey and composition of understorey for reptile, mammals and woodland birds.

One or more photos representing the habitat types on each site were taken at the beginning of the first survey of each of the sites. The structure of the canopy, shrub cover and ground cover was recorded for each site along with up to five of the most abundant plant species for each vegetation layer. Fauna habitat types were characterised in the study area in consideration of the vegetation mapping undertaken by Hunter Eco (2019).

Consideration was also given to the occurrence of habitat constraints in the *Threatened Biodiversity Data Collection* (OEH, 2019a) (Table 4).

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

**Table 4: Habitat Constraints Identified in the Threatened Biodiversity Data Collection**

Common Name	Credit Class	Habitat Constraints identified in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a)
<b>Amphibians</b>		
Green and Golden Bell Frog	Species	<i>Semi-permanent/ephemeral wet areas (within 1km of wet areas). Swamps (within 1km of swamp). Waterbodies (within 1km of waterbody).</i>
Green-thighed Frog	Species	<i>None.</i>
<b>Reptiles</b>		
Pink-tailed Legless Lizard	Species	<i>Rocky areas or within 50 m of rocky areas.</i>
Striped Legless Lizard	Species	<i>None.</i>
Pale-headed Snake	Species	<i>None.</i>
<b>Birds</b>		
Square-tailed Kite	Species/Ecosystem	<i>Breeding constraint: Other (Nest trees). Foraging constraint: none.</i>
White-bellied Sea-eagle	Species/Ecosystem	<i>Breeding constraint: Other (Living or dead mature trees within suitable vegetation within 1km of a rivers, lakes, large dams or creeks, wetlands and coastlines). Foraging constraint: Waterbodies (Within 1km of a rivers, lakes, large dams or creeks, wetlands and coastlines)</i>
Little Eagle	Species/Ecosystem	<i>Breeding constraint: Other (Nest trees - live (occasionally dead) large old trees within vegetation.). Foraging constraint: none</i>
Bush Stone-curlew	Species	<i>Fallen/standing dead timber including logs.</i>
Glossy Black-cockatoo	Species/Ecosystem	<i>Breeding constraint: Hollow-bearing trees (Living or dead tree with hollows greater than 15cm diameter and greater than 5m above ground). Foraging constraint: Other (Presence of Allocasuarina and Casuarina species)</i>
Gang-gang Cockatoo	Species/Ecosystem	<i>Breeding constraint: Hollow-bearing trees (Eucalypt tree species with hollows greater than 9 cm diameter) Foraging constraint: none</i>
Swift Parrot	Species/Ecosystem	<i>Breeding constraint: Other (As per mapped important areas – contact OEH for information). Foraging constraint: none.</i>
Masked Owl	Species/Ecosystem	<i>Breeding constraint: Hollow-bearing tree (Living or dead trees with hollows greater than 20cm diameter). Foraging constraint: none.</i>
Powerful Owl	Species/Ecosystem	<i>Breeding constraint: Hollow-bearing tree (Living or dead trees with hollows greater than 20cm diameter). Foraging constraint: none.</i>
Barking Owl	Species/Ecosystem	<i>Breeding constraint: Hollow-bearing tree (Living or dead trees with hollows greater than 20 cm diameter and greater than 4m above the ground). Foraging constraint: none.</i>
Regent Honeyeater	Species/Ecosystem	<i>Breeding constraint: Other (As per mapped areas; contact OEH) Foraging constraint: none</i>
<b>Mammals</b>		
Brush-tailed Phascogale	Species	<i>Hollow-bearing trees</i>
Common Planigale	Species	<i>None.</i>
Koala	Species/Ecosystem	<i>Breeding constraint: Other (Areas identified via survey as important habitat (see comments) Foraging constraint: none.</i>
Eastern Pygmy-possum	Species	<i>None</i>
Squirrel Glider	Species	<i>None</i>
Brush-tailed Rock-wallaby	Species	<i>Other (Land within 1 km of rocky escarpments, gorges, steep slopes, boulder piles, rock outcrops or cliff lines).</i>
Grey-headed Flying-fox	Species/Ecosystem	<i>Breeding constraint: Other (Breeding camps) Foraging constraint: none.</i>

**Table 4 (Continued): Habitat Constraints Identified in the Threatened Biodiversity Data Collection**

Common Name	Credit Class	Habitat Constraints identified in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a)
Little Bentwing-bat	Species/Ecosystem	<i>Breeding constraint: Caves (Cave, tunnel, mine, culvert or other structure known or suspected to be used for breeding).</i> <i>Foraging constraint: none.</i>
Eastern Bentwing-bat	Species/Ecosystem	<i>Breeding constraint: Caves (Cave, tunnel, mine, culvert or other structure known or suspected to be used for breeding).</i> <i>Foraging constraint: none.</i>
Large-eared Pied Bat	Species	<i>Cliffs (Within two kilometres of rocky areas containing caves, overhangs, escarpments, outcrops, or crevices, or within two kilometres of old mines or tunnels).</i>
Southern Myotis	Species	<i>Hollow-bearing trees (Within 200 m of riparian zone).</i> <i>Other (Bridges, caves or artificial structures within 200 m of riparian zone).</i>
Eastern Cave Bat	Species	<i>Caves (Within two kilometres of rocky areas containing caves, overhangs, escarpments, outcrops, crevices or boulder piles, or within two kilometres of old mines, tunnels, old buildings or sheds).</i>

### **Diurnal and Nocturnal Bird Surveys**

All ecologists recorded birds as they were encountered during the survey periods however the following information discussed in this section and within Tables 5 to 8 is provided by the dedicated avifauna specialist (Tony Saunders) on the ecological team.

All species that were encountered and identified by sight or call were recorded using the 'Sightings' App. Each significant patch of woodland habitat (a site) was searched using a 500 m radius survey. The time spent on a site was determined by the habitat quality, with the survey effort increased for higher quality sites when compared with lower quality sites. Incidental records of additional or locally significant species were also recorded while travelling around the site and between survey sites.

Specialist bird surveys were conducted in January 2018, June 2018, August 2018, September 2018, November 2018 and December 2018 to cover seasonality and detection requirements of several species.

In January 2018, early morning surveys were conducted between 05:30 and 11:30 hours. Evening surveys were run between 17:30 and 23:30 hours and included listening for calls and spotlighting. No surveys were conducted during the middle of the day in January 2018, as conditions were too hot for bird activity. A total of 42.25 hours was spent surveying 9 sites and an additional 11.25 hours was spent surveying while travelling between sites. The total survey effort over the 6 days from the 22-27 January 2018 was 53.5 hours. Diurnal survey effort was 20.65 hours and nocturnal survey effort was 21.6 hours. The survey effort for each site is summarised below with a break down showing time spent between diurnal and nocturnal surveys.

**Table 5: Bird Survey Effort for January 2018 Survey Period**

Site	1, 1a, 1b	2, 2a, 2b/3b, 2c, 2d	3, 3a, 3c	4, 4a	5, 5a, 5b	6, 6a, 6b, 6c	7, 7a, 7b, 7c	10, 10a, 10b	11, 11a	Totals
Diurnal (hrs)	3.45	1.70	2.25	1.00	3.00	4.75	1.50	1.00	2.00	<b>20.65</b>
Nocturnal (hrs)	5.00	3.30	1.80	3.00	4.25	4.25	Nil	Nil	Nil	<b>21.6</b>
<b>Totals (hrs)</b>	<b>8.45</b>	<b>5.00</b>	<b>4.05</b>	<b>4.00</b>	<b>7.25</b>	<b>9.00</b>	<b>1.50</b>	<b>1.00</b>	<b>2.00</b>	<b>42.25</b>

Sites 8 and 9 were not surveyed in January 2018 due to poor habitat.

In June 2018 diurnal surveys were conducted between 06:30 and 16:30 hours. Conditions were cool and birds were active throughout the day. Evening surveys were run between 18:30 and 20:30 hours and included listening for calls and spotlighting. Between 20 and 30 live trees and approximately 10 standing dead trees were spotlighted during each nocturnal survey in June 2018. A total of 29.00 hours was spent surveying 13 sites and an additional 6.00 hours was spent surveying while travelling between sites. The total survey effort over the 4 days from the 4-7 June 2018 was 35.00 hours. Diurnal survey effort was 23.75 hours and nocturnal survey effort was 5.25 hours. The survey effort for each site is summarised in the table below with a break down showing time spent between diurnal and nocturnal surveys.

**Table 6: Bird Survey Effort for June 2018 Survey Period**

Site	1, 1a, 1b	2, 2a, 2b/3b, 2c, 2d	3, 3a, 3c	4, 4a	5, 5a, 5b	6, 6a, 6b, 6c	7, 7a, 7b, 7c	10, 10a, 10b
Diurnal (hrs)	2.00	2.00	2.00	1.00	2.00	2.00	2.25	1.00
Nocturnal (hrs)	-	-	-	-	2.25	-	-	-
<b>Totals (hrs)</b>	<b>2.00</b>	<b>2.00</b>	<b>2.00</b>	<b>1.00</b>	<b>4.25</b>	<b>2.00</b>	<b>2.25</b>	<b>1.00</b>

Site	11, 11a	12	13	14a, 14b	Totals
Diurnal (hrs)	2.25	1.5	1.75	4.00	<b>23.75</b>
Nocturnal (hrs)	1.5	-	-	1.5	<b>5.25</b>
<b>Totals (hrs)</b>	<b>3.75</b>	<b>1.5</b>	<b>1.75</b>	<b>5.5</b>	<b>29</b>

In August 2018, diurnal surveys were conducted between 7:00 and 18:00 hours at Sites 15 and 16 located within the Maxwell Infrastructure area. Conditions were cool and birds were active throughout the day. Evening surveys were run between 18:00 and 19:30 hours and included listening for calls and spotlighting. Two hollow-bearing trees were stag watched on dusk at Site 15 on 28 August 2018. A total of 17.25 hours was spent surveying these two sites. Diurnal survey effort was 14.00 hours and nocturnal survey effort was 3.25 hours. The survey effort for each site is summarised in the table below with a break down showing time spent between diurnal and nocturnal surveys.

**Table 7: Bird Survey Effort for August 2018 Survey Period**

Site	15, 15a	16, 16a, 16b, 16c	Totals
Diurnal (hrs)	7.5	6.5	<b>14</b>
Nocturnal (hrs)	1.5	1.75	<b>3.25</b>
<b>Totals (hrs)</b>	<b>9.0</b>	<b>8.25</b>	<b>17.25</b>

In September 2018, diurnal surveys were conducted between 6:30 and 18:00 hours. Evening surveys were run between 18:30 and 20:30 hours and included listening for calls and spotlighting. Between 20 and 30 live trees and approximately 10 standing dead trees were spotlighted during each nocturnal survey. Incidental records of additional or locally significant species were also recorded while travelling around the site and between survey sites. Conditions were cool to warm, sunny and with only very light rain recorded 3 weeks previous to surveys. Heavy grazing pressure combined with the dry conditions had negatively impacted on bird diversity onsite during the survey period. Conditions were cool and calm in the mornings but became windy in the middle of the day and in the early afternoon.

Each significant patch of woodland habitat (a site) was searched using a 500 m radius survey. The time spent on a site was determined by the habitat quality, so that survey effort was increased for higher quality sites compared with lower quality sites.

In addition, several surveys were conducted targeting raptor species such as the Little Eagle and the Square-tailed Kite (*Lophoictinia isura*). For each of these surveys a good vantage point was selected to cover as much of the area under the development footprint as possible. Several vantage points were selected to ensure that the whole site was covered. From each vantage point the horizon was scanned for raptors hunting over the canopy and near the edges of remnant woodland, moving between remnants or circling in thermals.

In September 2018 a total of 21.0 hours was spent surveying eight sites, 7.5 hours was spent on aerial raptor searches and an additional 4.00 hours was spent surveying while travelling between sites. The total survey effort over the 4 days of survey was 32.5 hours. Diurnal survey effort was 16.0 hours and nocturnal survey effort was 5.0 hours. The survey effort for each site is summarised in the table below with a break down showing time spent between diurnal and nocturnal surveys.

**Table 8: Bird Survey Effort for September 2018 Survey Period**

Site	1, 1a, 1b	2, 2a, 2b/3b, 2c, 2d	3, 3a, 3c	5, 5a, 5b	6, 6a, 6b, 6c	7, 7a, 7b, 7c	8, 8a	11, 11a	Totals
Diurnal (hrs)	4.5	2.5	1.0	2.5	1.5	1.5	0.5	2.0	<b>16</b>
Nocturnal (hrs)	1.5	1.5	-	2.0	-	-	-	-	<b>5</b>
<b>Totals (hrs)</b>	<b>6.0</b>	<b>4.0</b>	<b>1.0</b>	<b>4.5</b>	<b>1.5</b>	<b>1.5</b>	<b>0.5</b>	<b>2.0</b>	<b>21</b>

In November 2018 surveys were conducted over an eight-day period between 12-23 November 2018 inclusive. Conditions were cool to warm, sunny, dry and sometimes windy. There was evidence of some heavy rain events since the previous surveys and most dams had good levels of water in them. Heavy grazing pressure combined with very dry conditions had negatively impacted on bird diversity onsite during the survey period. There were some signs of recovery, but this was not very advanced at the time of the surveys. Conditions were cool and calm in the mornings and observing conditions were ideal. When conditions became windy surveys were terminated.

Each significant patch of woodland habitat (a site) was searched using a 500 m radius survey. The time spent on a site was determined by the habitat quality, so that survey effort was increased for higher quality sites compared with lower quality sites. Based on the PCTs present, there was an emphasis on sites that contained potential habitat for Regent Honeyeaters (*Anthochaera phrygia*).

In addition, several surveys were conducted targeting raptor species such as the Little Eagle, Square-tailed Kite and Spotted Harrier. For each of these surveys a good vantage point was selected to cover as much of the area under the development footprint as possible. Several vantage points were selected to ensure that the whole site was covered. From each vantage point the horizon was scanned for raptors hunting over the canopy and near the edges of remnant woodland, moving between remnants or circling in thermals. Several potential raptor nests were also recorded within the 'Sightings' App and whether birds were present or not.

Most sites were surveyed by two observers who covered different parts of a site during each visit. Diurnal surveys were conducted between 05:00 and 19:30 hours. Evening surveys were run between 19:30 and 22:30 hours and included listening for calls and spotlighting. A minimum of 30 live trees and 10 standing dead trees (if present) were spotlighted during each nocturnal survey. Incidental records of additional or locally significant species were also recorded while travelling around the site and between survey sites.

In November 2018 a total of 124.25 hours was spent surveying 11 sites. An additional 24 hours were spent on aerial raptor searches and incidental surveying while travelling between sites. Diurnal survey effort was 136.25 hours and nocturnal survey effort was 12.0 hours. Total survey effort was 148.25 hours. The survey effort for each site is summarised in the table below with a break down showing time spent between diurnal and nocturnal surveys.

**Table 9: Bird Survey Effort for November 2018 Survey Period**

Site	1, 1a, 1b	2, 2a, 2b/3 b, 2c, 2d	3, 3a, 3c	4, 4a	5, 5a, 5b	6, 6a, 6b, 6c	7, 7a, 7b, 7c	9	10, 10a, 10b	17	18, 18a	Raptors / Incidental	Totals
Diurnal (hrs)	4.5	23	8	6	36	8	7.5	1	1.75	12	4.5	24	<b>136.25</b>
Nocturnal (hrs)	-	5	-	-	4	-	-	-	-	-	3	-	<b>12</b>
<b>Totals (hrs)</b>	<b>4.5</b>	<b>28</b>	<b>8</b>	<b>6</b>	<b>40</b>	<b>8</b>	<b>7.5</b>	<b>1</b>	<b>1.75</b>	<b>12</b>	<b>7.5</b>	<b>24</b>	<b>148.25</b>

In December 2018 surveys were conducted over a six and a half-day period between 3-19 December 2018 inclusive. Conditions were warm, sunny, dry and sometimes windy to hot and humid with storms building. There was evidence of some heavy rain events since the previous surveys and most dams had good levels of water in them. Heavy grazing pressure combined with very dry conditions negatively impacted on bird diversity onsite during the surveys. The recovery was more advanced at the time of the final surveys undertaken in December 2018 as the effects of the rain storms and the removal of cattle from the property was evident. Conditions were cool and calm in the mornings and observing conditions were ideal.

Each significant patch of woodland habitat (a site) was searched using a 500 m radius survey. The time spent on a site was determined by the habitat quality, so that survey effort was increased for higher quality sites compared with lower quality sites. There was an emphasis on sites that contained potential habitat for Regent Honeyeaters.

All sites containing PCTs that were possible habitat for the Regent Honeyeater were surveyed for a minimum of 20 hours each. These include sites 2, 3, 4, 5, 6, 7 and 17.

Additional potential raptor nests were also recorded including whether birds were present or not in the 'Sightings' App.

Most sites were surveyed by two observers who covered different parts of a site during each visit. Diurnal surveys were conducted between 5:00 and 20:30 hours. Evening surveys were run between 20:30 and 22:30 hours and included listening for calls and spotlighting. A minimum of 30 live trees and 10 standing dead trees (if present) were spot-lighted during each nocturnal survey. Incidental records of additional or locally significant species were also recorded while travelling around the site and between survey sites.

In December 2018 a total of 67.75 hours was spent surveying 10 sites. An additional 12.5 hours were spent on aerial raptor searches and incidental surveying while travelling between sites. Diurnal survey effort was 80.25 hours and nocturnal survey effort was 11.75 hours. The total survey effort was 91 hours. The survey effort for each site is summarised in the table below with a break down showing time spent between diurnal and nocturnal surveys.

**Table 10: Bird Survey Effort for December 2018 Survey Period**

Site	1, 1a, 1b	3, 3a, 3c	4, 4a	6, 6a, 6b, 6c	7, 7a, 7b, 7c	11, 11a	14a, 14b	17	18, 18a	Savoy Dam	Raptors / Incidental	Totals
Diurnal (hrs)	11.0	12.75	14.25	0.75	1.75	10.0	-	8.0	3.25	6.0	12.5	<b>80.25</b>
Nocturnal (hrs)	4.0	-	-	-	-	-	3.0	3.75	-	-1.0	-	<b>11.75</b>
<b>Totals (hrs)</b>	<b>15.0</b>	<b>12.75</b>	<b>14.25</b>	<b>0.75</b>	<b>1.75</b>	<b>10.0</b>	<b>3.0</b>	<b>11.75</b>	<b>3.25</b>	<b>7.0</b>	<b>12.5</b>	<b>91</b>

During all survey periods signs of owl use were searched for under and on some hollow-bearing trees with hollows large enough to accommodate large forest owl species (i.e. owl pellets, remains of meals, faecal whitewash, feathers).

### **Ground Elliott Trapping**

Elliott traps targeting small to medium sized ground-dwelling mammals were set out for four consecutive nights from 23-27 January 2018 (Sites 1a, 2a and 3b), and 19-23 November 2018 (Site 7b). A total of 25 “A” Elliott traps (measuring 33 centimetres (cm) x 10 cm x 9 cm) were deployed at each of the relevant sites.

Figure 4 shows the location of the mammal trapping sites, which included ground Elliott traps.

Trap lines typically traversed areas of diverse vegetation or habitat features as identified from the habitat search as likely areas to support the target mammal. Each trap was baited with a bait mix of peanut butter, honey, molasses, rolled oats, vanilla essence, almond essence and fish sauce.

Dry bedding material (leaves or coconut husk) was placed in each Elliott trap and the traps were covered in plastic bags if wet weather threatened. Traps were checked early each morning for captures, with any captured animals identified and immediately released. Traps were left closed for the day and re-opened on dusk.

### **Arboreal Elliott Trapping**

Elliott traps targeting arboreal species identified from the literature review, namely the Squirrel Glider, were placed in habitat with large trees (some with hollows) and vegetated ground cover, for four consecutive nights from 23-27 January 2018 (Sites 1a, 2a and 3a) and 19-23 November 2018 (Sites 5d, 6b, 7b, 18a).

Figure 4 shows the location of the trapping sites.

Ten “B” Elliott traps (15 cm x 15 cm x 56 cm) were deployed at each of the relevant sites. The traps were placed greater than two meters off the ground on a platform fixed to the trunk of the tree at approximately 30-50 m spacing.

Each trap was baited with a bait mix of peanut butter, honey, molasses, rolled oats, vanilla essence, almond essence and fish sauce.

Dry bedding material (leaves or coconut husk) was placed in each Elliott trap and the traps were covered in plastic bags if wet weather threatened. Traps were checked early each morning for captures, with any captured animals identified and immediately released. Traps were left closed for the day and re-opened on dusk.

### **Cage Trapping**

Cage traps targeting medium mammals were deployed at Sites 1a, 2a and 3b, for four consecutive nights from 23-27 January 2018, as part of the terrestrial and arboreal Elliott trapping grids. An additional cage was deployed at Site 3. Figure 4 shows the location of the trapping sites.

Three cages were Mascot Wire Works steel traps measuring 20 cm x 20 cm x 56 cm with a 12.5 x 50 mm mesh (i.e. 'bandicoot' sized traps). The fourth cage was a larger 'dog' sized trap also from Mascot Wire Works and this was deployed at Site 2a.

Cages were baited with sardines and covered in hessian bags. Traps were checked early each morning for captures, with any captured animals identified and immediately released. Traps were left closed for the day and re-opened on dusk.

Between 27 January and 6 June 2018, a single bandicoot sized cage at Site 3 was closed, baited with fresh sardines and a jar of molasses and left onsite as part of bait stations for longer term camera trapping (130 nights).

### **Hair Tubes**

Hair tube surveys, targeting small to medium-sized arboreal and terrestrial mammals, were deployed for at least four-five consecutive evenings as follows:

- Site 1a – 21 tubes over five nights (22-27 January 2018).
- Site 2a – 22 tubes over four nights (23-27 January 2018).
- Site 3a – 20 tubes over four nights (23-27 January 2018).
- Site 5a – two tubes over four nights (23-27 January 2018).
- Site 6a – two tubes over four nights (23-27 January 2018).
- Site 16a – five tubes over 76 nights from (29 August 29 – 13 November 2018).
- Site 16b – five tubes over 76 nights from (29 August 29 – 13 November 2018).
- Site 18a – 20 tubes over 18 nights (16 November – 4 December 2018).

Figure 4 shows the location of the trapping sites.

All hair tubes were single sided and five sizes of were used, 90 mm diameter (large), 50 mm diameter (medium), 40 mm diameter (small), 30 mm diameter (extra small) and Faunatech funnels. Double-sided tape was placed at the entrance on the upper side of the tube to collect hairs of animals attracted to the bait. All tubes were baited with peanut butter, honey, molasses, rolled oats, vanilla essence, almond essence and fish sauce. Tubes at Sites 15 and 16 were baited with a mix of flour, sardines, and tuna oil.

Hair tubes were set on the ground, in shrub/tree bases and in trees at a height approximately 1-1.5 m above the ground.

Between 27 January and 6 June 2018, a number of fresh hair tubes were deployed as part of bait stations for longer term camera trapping as follows:

- Site 1a – two tubes over 130 nights.
- Site 2a – four tubes over 130 nights.
- Site 3a – four tubes over 130 nights.

Any hairs collected were sent to an expert in hair analysis (Barbara Triggs, Genoa Victoria) for analysis.



### **Camera Trapping**

Wildlife cameras were deployed as part of live trapping and/or hair tube transects at Sites 1, 2, 3, 5, 6, 7 and 17 in areas of suitable habitat. Several types of cameras were used including Scout Guard SG562C White Flash, Scout Guard SG550V, Reconyx PC900, Reconyx PC90 and Moultrie D80.

All cameras were set to take still photos only.

Cameras were deployed as follows:

- Site 1a – two cameras each placed for five nights (22-27 January 2018).
- Site 2b – two cameras each placed for four nights (23-27 January 2018).
- Site 3c – two cameras each placed for four nights (23-27 January 2018).
- Site 5a – one camera placed for four nights (23-27 January 2018).
- Site 6a – one camera placed for four nights (23-27 January 2018).
- Site 6b – one camera placed for 21 nights (13 November – 4 December 2018).
- Site 7c – one camera placed for 20 nights (14 November – 4 December 2018).
- Site 17a – two cameras each placed for 15 nights (4 December – 20 December 2018).
- Location IncA – two cameras each placed for 3 nights (26 December – 29 December 2018).

Figure 4 shows the location of the trapping sites.

Cameras were pointed at bait stations or closed cages baited with a lure of sardines, and/or hair tubes baited with peanut butter, honey, molasses, rolled oats, vanilla essence, almond essence and fish sauce.

Between 27 January and 6 June 2018, cameras were deployed as part of bait stations for longer term camera trapping as follows:

- Site 1a – one camera over 130 nights.
- Site 2a – one camera over 130 nights.
- Site 3a – one camera over 130 nights.

In addition, two cameras were deployed facing a fresh cattle carcass at a location between Sites 5 and 6 and also left for 130 nights as above.

The use of camera traps is an additional survey technique to those described in DEC (2004a) but is discussed in SEWPaC (2011a) in regard to threatened mammals.

### **Nest boxes**

15 nest boxes (five at each of Sites 1a, 2a and 3a – Figure 4) were deployed for 132-133 nights (24/25 January – June 6 2018). Nest boxes were made from natural recycled hollows and were generally of dimensions and entrance diameters suitable for use by small to medium sized arboreal mammals such as Squirrel Glider, Eastern Pygmy-possum (*Cercartetus nanus*) and microbats.

### **Bat Surveys**

Bats were surveyed in accordance with the 'Species Credit' *Threatened Bats and their Habitat: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018).

Surveys were undertaken by appropriately experienced bat surveyors, Adam Greenhalgh B.App.Sc., Garon Staines B.App.Sc., and Nick Everitt B.Env.Sc., and bat call identification was undertaken by Amanda Lo Cascio B.Sc.,M.Env (2018; 2019). These surveyors each have over 10 years of experience surveying and identifying bats in NSW.

All bat species in Table 1 were targeted during the surveys, however species in Table 11 were specifically targeted in accordance with the *'Species Credit' Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018). The survey requirements and survey details are provided in Table 11.

The bat surveys were undertaken in January 2018, November 2018 and December 2018 (Table 11) in suitable weather conditions (Section 2.3.1).

The bat surveys for the Grey-headed Flying-fox, Corben's Long eared Bat and Large-eared Pied Bat were also undertaken in consideration of the *Survey Guidelines for Australia's Threatened Bats* (DEWHA, 2010b).

The following sections provide the dates of each survey, details of the methods (harp trapping, ultrasonic bat detection and microbat habitat searches) and total survey effort for bats.

### **Harp Trapping**

Harp trapping for insectivorous bats was carried out at Sites 1, 2, 3, 5, 6 and a mine dam (known as Savoy Dam) as follows:

- Site 1a – one harp trap placed for two nights (22-23 January 2018).
- Site 1d – Dam 30 - one harp trap placed for one night (26 January 2018).
- Site 1b – Quarry Pond - two harp traps placed for four nights (3-7 December 2018).
- Site 1c – Power Easement - four harp traps placed for three nights (17, 18 and 20 December 2018).
- Site 2a – one harp trap placed for two nights (23-24 January 2018).
- Site 2b – Dam 29 - two harp traps place for three nights (12-15 November 2018).
- Site 3a – one harp trap placed for two nights (24-25 January 2018).
- Site 3b – Dam 28 - two harp traps placed for three nights (12-15 November 2018).
- Site 5b – Dam 22 - two harp traps placed for four nights (15, 19, 21 and 22 November 2018).
- Site 5c – Dam 23 - two harp traps placed for four nights (15, 19, 21 and 22 November 2018).
- Site 6b – one harp trap placed for two nights (25-26 January 2018).
- Savoy Dam – two harp traps placed for four nights (3-7 December 2018).

Harp traps were set at the above sites within potential flyways wherever possible and/or adjacent to waterbodies.

Harp traps were inspected for captures usually once at night and then again before dawn and then disarmed for the day. Any captures were identified to species level and then released prior to sunrise the same day or were held during the day in cotton bags back at the accommodation and then released at dusk.

Figure 5 shows the bat survey sites.

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 11: 'Species Credit' Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method (OEH, 2018)

Species	Credit Type <sup>1</sup>	Potential Habitat <sup>#</sup>	Survey Method	Survey Period	Required Survey Effort <sup>#</sup>	Required Minimum Number of Days <sup>#</sup>	Actual Survey Details
Grey-headed Flying-fox <i>Pteropus poliocephalus</i>	Species Credit for Breeding Habitat	<i>The initial search for camps should encompass any recorded camps and roosting habitat likely to occur on the subject land. If a camp is located the survey only needs to take place in the camp (that is the area occupied by the target species) to identify breeding females</i>	Daytime camp survey	Oct – Dec	6 hrs (two hours/day)	3 (one per month)	No camps are known to occur in the study area (DEE, 2019). Numerous daytime searches were conducted throughout the study area. No camps were found and therefore the survey effort is not applicable.
Little Bentwing-bat <i>Miniopterus australis</i>	Species Credit for Breeding Habitat	<i>Caves, tunnels, mines or other structures known or suspected to be used by M.australis including species records in the NSW BioNet Atlas with microhabitat code 'IC –in cave'; observation type code 'E nest-roost'; with numbers of individuals &gt;500; or from the scientific literature.</i>	Harp trap	Dec – Feb	8	4	There are no <i>BioNet Atlas</i> (OEH, 2019b) records of these two species in the study area with 'microhabitat code 'IC –in cave'; observation type code 'E nest-roost'; with numbers of individuals >500'. No caves, tunnels or disused underground mines occur in the study area. The study area does not appear to provide the deep (often limestone) cave habitat required by these species for maternity roosts. Sub-optimal potential habitat was only detected at Site 1 (old quarry and a nearby rocky hill) and a small rocky escarpment near the corner of Edderton Road and Golden Highway.
Eastern Bentwing-bat <i>Miniopterus schreibersii oceanensis</i>	Species Credit for Breeding Habitat	<i>Caves, tunnels, mines or other structures known or suspected to be used by M.schreibersii oceanensis including species records in the NSW BioNet Atlas with microhabitat code 'IC –in cave'; observation type code 'E nest-roost'; with numbers of individuals &gt;500; or from the scientific literature</i>	Harp trap	Dec – Feb	8	4	Harp trapping was carried out at Site 1 in early December 2018 using two harp traps over four nights and repeated two weeks later in December 2018 using four harp traps over three nights (i.e. a total effort of 20 trap-nights over 7 nights). Rocky crevices at Site 1 and along a small escarpment near corner of Edderton Road and Golden Highway were inspected for bat roosts. Acoustic recording devices at Site 1 for 16 detector nights in total in January and December 2018 and at small escarpment Edderton Road/Golden Highway for two detector nights in November 2018.
Large-eared Pied Bat <i>Chalinolobus dwyeri</i>	Species Credit	<i>The PCTs associated with the species (as per the Threatened Biodiversity Data Collection) within 100m of rocky areas containing caves, or overhangs or crevices, cliffs or escarpments, or old mines, tunnels, culverts, derelict concrete buildings. Traps should be set in woodlands, valley floors, riparian areas and relatively fertile parts of the subject land where possible.</i>	Harp trap or mist net	Mid Nov – end Jan	16	4	No caves, cliffs, escarpments, tunnels, disused underground mines or derelict buildings occur in the study area. The study area does not appear to provide the deep sandstone overhang with domed roof habitat required by these species for maternity roosts. Sub-optimal potential habitat was only detected at Site 1 (old quarry and a nearby rocky hill) and a small rocky escarpment near the corner of Edderton Road and Golden Highway. Harp trapping was carried out at Site 1 in early December 2018 using two harp traps over four nights and repeated two weeks later in late December 2018 using four harp traps over three nights (i.e. a total effort of 20 trap-nights over 7 nights).
			Acoustic detection	Mid Nov – end Jan	16	4	Rocky crevices at Site 1 and along a small escarpment near corner of Edderton Road and Golden Highway were inspected for bat roosts. Acoustic recording devices at Site 1 for 16 detector nights in total in January and December 2018 and at the small escarpment near Edderton Road/Golden Highway for two detector nights in November 2018. 65 harp trap nights were carried out across the study area in January, November and December 2018. Acoustic recording (34 trap nights) were conducted in January, November and December 2018 at a number of sites across the study area. Culverts were inspected at Sites 10 and 17.

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 11 (Continued): 'Species Credit' Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method (OEH, 2018)

Species	Credit Type <sup>1</sup>	Potential Habitat <sup>#</sup>	Survey Method	Survey Period	Required Survey Effort <sup>#</sup>	Required Minimum Number of Days <sup>#</sup>	Actual Survey Details
Southern Myotis <i>Myotis macropus</i>	Species Credit	<i>The range of PCTs associated with the species (as per the Threatened Biodiversity Data Collection) within 200 meters of any medium to large permanent creeks, rivers, lakes or other waterways (i.e. with pools/ stretches 3m or wider)</i>	Harp trap or mist net	Oct – Mar	16	4	No medium to large permanent creeks, rivers or lakes occur in the study area. The study area does contain a number of farm and mine water dams. Some of these dams are within 200 m of relevant PCTs associated with this species in the study area. Harp trapping was carried out next to dam/ponds at Sites 1, 2, 3, 5 and a mine dam (known as Savoy Dam) in November and December 2018. 65 harp trap nights were carried out across the study area in January, November and December 2018. Acoustic recording (34 trap nights) were conducted in January, November and December 2018 at a number of sites across the study area. Culverts were inspected at Sites 10 and 17.
			Roost search	Oct – Mar	1 per structure	30 min per feature	
			Acoustic detection	Oct – Mar	16	4	
Eastern Cave Bat <i>Vespadelus troughtoni</i>	Species Credit	<i>The PCTs associated with the species (as per the Threatened Biodiversity Data Collection) within 100m of rocky areas, caves, overhangs crevices, cliffs and escarpments, or old mines or tunnels, old buildings and sheds within the potential habitat. Traps should be set in woodlands, valley floors, riparian areas and relatively fertile parts of the subject land where possible.</i>	Harp trap or mist net	Nov – end Jan	16	4	No caves, cliffs, escarpments, tunnels, disused underground mines or derelict buildings occur in the study area. Sub-optimal potential habitat was only detected at Site 1 (old quarry and a nearby rocky hill) and a small rocky escarpment near corner of Edderton Road and Golden Highway. Harp trapping was carried out at Site 1 in early December 2018 using two harp traps over four nights and repeated two weeks later in December 2018 using four harp traps over three nights (i.e. a total effort of 20 trap-nights over 7 nights). Rocky crevices at Site 1 and along a small escarpment near corner of Edderton Road and Golden Highway were inspected for bat roosts. Acoustic recording devices at Site 1 for 16 detector nights in total in January and December 2018 and at small escarpment Edderton Road/Golden Highway for two detector nights in November 2018. 65 harp trap nights were carried out across the study area in January, November and December 2018. Acoustic recording (34 trap nights) were conducted in January, November and December 2018 at a number of sites across the study area.
			Roost search	Nov – end Jan	1 per structure	30 min per feature	
			Acoustic detection	Nov – end Jan	16	4	

<sup>1</sup> Biodiversity credit class under the *Threatened Biodiversity Data Collection* (OEH, 2019a) (current as at March 2019).

<sup>#</sup> 'Species Credit' *Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018).

### ***Ultrasonic Bat Detection - Anabat***

Electronic detectors were used to collect ultrasonic calls of microbat species. Detectors used included: Anabat SD1 and Express detectors (Titley Scientific, Brisbane QLD), and an SMBAT2+ Songmeter and Echo Meter Touch 2 PRO (Wildlife Acoustics, Maynard, MA, USA).

Detectors were used both statically (i.e. set and left in one location overnight) and actively (carried around) when undertaking nocturnal searches.

Static recording detectors were placed in an area of habitat, left for a minimum of two nights, placed at ground level or off the ground aiming along potential microbat flyways that microbats could use to forage and navigate their way through woodland areas. For active recording, Anabats were carried when spotlighting surveys were undertaken recording as the surveyor passes through habitat.

Electronic bat call recording units were deployed during the January 2018 survey period as follows:

- Site 1a – one unit placed for two nights (22-23 January 2018).
- Site 2a – one unit placed for two nights (23-24 January 2018).
- Site 3a – one unit placed for two nights (24-25 January 2018).
- Site 4a – one unit placed for two nights (25-26 January 2018).
- Site 5a – one unit placed for two nights (23-24 January 2018).
- Site 6a – one unit placed for two nights (23-24 January 2018).
- Site 7a – one unit placed for two nights (23-24 January 2018).
- Site 10a:
  - One unit placed for two nights at culvert (25-26 January 2018).
  - One unit placed at dam for 0.25 hours (6 June 2018).
- Site 11a – one unit placed for one night (26 January 2018).
- Two units were placed at an observed bat roost tree between Sites 5 and 10 (Site IncA) at dusk on 27 January 2018 for approximately 0.5 hour as several bats emerged from the tree.
- A mobile unit was carried across several sites over several nights during nocturnal work including Site 1 (22 January 2018 for 0.25 hour), Site 2 (23 January 2018 for 1 hour) and Site 5 (24 January 2018 for 0.25 hour).
- Site 1b Quarry Pond – one detector placed for four nights, one detector placed for three nights (3-7 December 2018).
- Site 1c Power Easement – one detector placed for four nights, one detector placed for three nights (17-21 December 2018).
- Rocky escarpment near corner of Golden Highway and Edderton Road (Site IncB) – one detector placed for two nights (5-7 December 2018).

During the June 2018 survey period a bat detector unit was deployed for approximately 0.25 hour to assist the identification of two bats observed foraging over a farm dam at Site 10.

Figure 5 shows the bat survey sites.

### **Microbat Habitat Searches**

Culverts were visually inspected during the day for roosting bats as follows:

- Sites 1a and 1b various dates in November and December 2018.
- Site 10a – Site 10 culvert – 26 January 2018.
- Site 17a – Site 17 culvert – 12 November and 20 December 2018.
- Site 17b – Rail Loop Dam culvert – 20 December 2018.
- Site IncB – various dates November and December 2018.

Figure 5 shows the bat survey sites.

### **Nocturnal Call Playback**

The playback of pre-recorded calls of threatened nocturnal species was carried out at dusk or after dark using digital MP3 players coupled to loudhailers or portable speakers.

After an initial listening period of ten minutes, each call was played for a total of five minutes, followed by a five-minute listening period, with the last listening period followed by at least ten minutes of spotlighting.

Species targeted (in order of call-playback) were the Koala, Squirrel Glider, Yellow-bellied Glider (*Petaurus australis*), Powerful Owl (*Ninox strenua*), Masked Owl (*Tyto novaehollandiae*), Barking Owl, Sooty Owl (*Tyto tenebricosa*) and Bush Stone-curlew (*Burhinus grallarius*). During the November/December 2018 surveys, calls of Green and Golden Bell Frog (*Litoria aurea*) were also broadcast at sites with potential habitat (dams, ponds, drainage lines with sedges and reeds). Any fauna responding were identified either by characteristic call or direct observation using spotlights.

Figure 4 shows the mammal survey sites which includes call-playback.

Call playback was carried out as follows:

- Site 1:
  - Site 1a – Power Easement – 22 January 2018.
  - Site 1a – Power Easement – 27 January 2018.
  - Site 1a – Power Easement – 17 September 2018.
  - Site 1b – Quarry Pond – 3 December 2018.
  - Site 1b – Quarry Pond – 5 December 2018.
  - Site 1b – Quarry Pond – 6 December 2018.
  - Site 1a – Power Easement – 17 December 2018.
  - Site 1a – Power Easement – 18 December 2018.
  - Site 1a – Power Easement – 21 December 2018.
- Site 2:
  - Site 2a – 23 January 2018.
  - Site 2a – 26 January 2018.
  - Site 2b/3b – 18 September 2018 in paddock on eastern side (note that the call playback survey at this location also covered both Sites 2 and 3, due to the volume at which the calls were played).
  - Site 2c – 12 November 2018.

- Site 2c – 12 November 2018.
- Site 2c – Dam 29 – 14 November 2018.
- Site 3:
  - Site 3a – 23 January 2018 (note that the call playback survey at this location also covered Sites 2 and 3, located within 500 m, due to the volume at which the calls were played).
  - Site 3a – 26 January 2018 (note that the call playback survey at this location also covered Site 2, located downslope within 500 m, due to the volume at the which the calls were played).
  - Site 2b/3b – 18 September 2018 in paddock on eastern side (note that this location covered both Sites 2 and 3).
  - Site 3c – 12 November 2018.
  - Site 3c – 14 November 2018.
- Site 4:
  - Site 4a – 26 January 2018.
  - Site 4a – 4 December 2018.
  - Site 4a – 6 December 2018.
- Site 5:
  - Site 5a – 24 January 2018.
  - Site 5a – 5 June 2018.
  - Site 5b – 19 September 2018.
  - Site 5a – 12 November 2018.
  - Site 5a – 13 November 2018
  - Site 5b – 15 November 2018.
  - Site 5b – 20 November 2018.
- Site 6:
  - Site 6a – 13 November 2018.
  - Site 6c – 15 November 2018 (note that the call playback survey at this location also covered Site 18, located within 500 m downslope, due to the volume at which the calls were played).
  - Site 6b – 5 December 2018 (note that the call playback survey at this location also covered Site 7 located within 800 m upslope, and Site 18, approximately 800 m downslope, due to the volume at which the calls were played).
- Site 7:
  - Site 7a – 25 January 2018 (note that the call playback survey at this location also covered Site 6, located downslope within 500 m, due to the volume at which the calls were played).
  - Site 7a – 14 November 2018 (note that the call playback survey at this location also covered Sites 6, 8, 13, 14 and 18, located downslope within 0.5-2 km, as the survey was on a very still night and domestic dogs from 'Edderton' property 2 km away to north-west were heard responding to calls).
  - Site 7b – 21 November 2018.
  - Site 7b – 22 November 2018.

- Site 8:
  - Site 8a – 4 December 2018 (note that the call playback survey at this location also covered Site 13, located approximately 600 m downstream, due to the volume at which the calls were played).
- Site 10:
  - Site 10a – 6 June 2018.
  - Site 10b – 19 December 2018.
- Site 11:
  - January, June, November, December 2018 sessions covered by nearby Site 5a call-playback sessions (see above).
- Site 13:
  - Covered by call-playback surveys conducted at Sites 7a, 8a and 14b within 600m of this site.
- Site 14:
  - Site 14a – 4 June 2018.
  - Site 14b – 5 December 2018 (note that the call playback survey at this location also covered Site 13, located approximately 500 m downslope, due to the volume at which the calls were played).
  - Site 14b – 19 December 2018 (note that the call playback survey at this location also covered Site 13, located approximately 500 m downslope, due to the volume at which the calls were played).
- Site 15a – 28 August 2018.
- Site 16a – 29 August 2018.
- Site 17:
  - Site 17a – 4 December 2018.
  - Site 17a – 6 December 2018.
- Site 18:
  - Site 18a – 4 December 2018.
  - Site 18a – 6 December 2018.
- a mine dam (known as Savoy Dam):
  - Savoy – 3 December 2018.
  - Savoy – 6 December 2018.

### **Spotlighting**

Spotlighting was conducted on foot by using powerful LED hand-held torches and/or headlamps. It was undertaken in conjunction with call-playback and nocturnal searches for fauna at several sites over several nights and survey periods as follows:

- Site 1:
  - Site 1a – Power Easement – 22 January 2018 (five observers for 1.25 hours each).
  - Site 1a – Power Easement – 27 January 2018 (five observers for 1.25 hours each).
  - Site 1a – Power Easement – 17 September 2018 (two observers for 1.25 hours each).



- Site 1c – Quarry Pond – 3 December 2018 (two observers for 1.5 hours each).
- Site 1c – Quarry Pond – 5 December 2018 (two observers for 0.25 hours each).
- Site 1c – Quarry Pond – 6 December 2018 (two observers for 0.25 hours each).
- Site 1a – Power Easement – 17 December 2018 (two observers for 1 hour each).
- Site 1a – Power Easement – 18 December 2018 (two observers for 1 hour each).
- Site 1a – Power Easement – 20 December 2018 (two observers for 0.25 hours each).
- Site 1a – Power Easement – 21 December 2018 (two observers for 0.5 hour each).
- Site 2:
  - Site 2a – 23 January 2018 (two observers for 1 hour each).
  - Site 2a – 26 January 2018 (three observers for 1 hour each).
  - Site 2a – 27 January 2018 (four observers for 1 hour each).
  - Site 2b/3b – 18 September 2018 (two observers for 0.625 hours each).
  - Site 2c – 12 November 2018 (two observers for 1 hour each).
  - Site 2c – 13 November 2018 (two observers for 1.25 hours each).
  - Site 2c – 14 November 2018 (three observers for 0.75 hours each).
- Site 3:
  - Site 3a – 23 January 2018 (three observers for 1 hour each).
  - Site 3a – 26 January 2018 (two observers for 1 hour each).
  - Site 3a – 27 January 2018 (one observer for 0.5 hour).
  - Site 2b/3b – 18 September 2018 (two observers for 0.625 hours each).
  - Site 3c – 12 November 2018 (two observers for 1.75 hours each).
  - Site 3c – 14 November 2018 (three observers for 1.25 hours each).
- Site 4:
  - Site 4a – 26 January 2018 (five observers for 0.75 hours each).
  - Site 4a – 4 December 2018 (two observers for 0.75 hours each).
  - Site 4a – 6 December 2018 (two observers for 1 hours each).
  - Site 4a – 6 December 2018 (two observers for 0.25 hours each).
- Site 5:
  - Site 5a – 24 January 2018 (three observers for 1.25 hours each).
  - Site 5a – 5 June 2018 (two observers for 2 hours each).
  - Site 5b – 19 September 2018 (two observers for 1.25 hours each).
  - Site 5a – 12 November 2018 (two observers for 1.5 hours each).
  - Site 5a – 13 November 2018 (two observers for 0.5 hours each).
  - Site 5b – 15 November 2018 (three observers for 1.25 hours each).
  - Site 5b – 20 November 2018 (two observers for 1.25 hours each).
  - Site 5b – 20 December 2018 (two observers for 0.25 hours each).

- Site 6:
  - Site 6a – 25 January 2018 (three observers for 1.5 hours each).
  - Site 6a – 13 November 2018 (two observers for 1 hour each).
  - Site 6b – 14 November 2018 (one observer for 1.25 hours).
  - Site 6c – 15 November 2018 (three observers for 1.25 hours each).
  - Site 6b – 5 December 2018 (three observers for 1.25 hours each).
- Site 7:
  - Site 7a – 25 January 2018 (two observers for 2 hours each, three observers for 1 hour).
  - Site 7a – 14 November 2018 (two observers for 1.25 hours each).
  - Site 7b – 21 November 2018 (three observers for 1.25 hours each).
  - Site 7b – 22 November 2018 (three observers for 1.25 hours each).
- Site 8a – 4 December 2018 (two observers for 0.25 hours each).
- Site 10:
  - Site 10a – 6 June 2018 (two observers for 1.25 hours each).
  - Site 10b – 19 December 2018 (two observers for 1.25 hours each).
- Site 11:
  - Site 11a – 24 January 2018 (two observers for 1.25 hours each).
  - Site 11a – 13 November 2018 (two observers for 1.25 hours each).
  - Site 11a – 20 December 2018 (two observers for 0.25 hours each).
- Site 14:
  - Site 14a – 4 June 2018 (two observers for 1.5 hours each).
  - Site 14b – 5 December 2018 (three observers for 0.5 hours each).
  - Site 14b – 19 December 2018 (two observers for 2 hours each).
- Site 15a – 28 August 2018 (two observers for 1.5 hours each).
- Site 16a – 29 August 2018 (two observers for 1.75 hours each).
- Site 17:
  - Site 17a – 4 December 2018 (two observers for 1.5 hours each).
  - Site 17a – 6 December 2018 (one observer for 0.75 hours).
- Site 18:
  - Site 18a – 15 November 2018 (two observers for 1.5 hours each).
  - Site 18a – 4 December 2018 (two observers for 0.75 hours each).
  - Site 18a – 6 December 2018 (two observers for 0.75 hours each).
- a mine dam (known as Savoy Dam):
  - Savoy – 3 December 2018 (two observers for 1 hour each).
  - Savoy – 6 December 2018 (two observers for 0.5 hours each).

Figure 4 shows the mammal survey sites which includes spotlighting.

### **Koala Scat Searches**

One preferred food species listed in NSW *State Environmental Planning Policy No. 44 – Koala Habitat Protection* (SEPP 44) Schedule 2 Koala feed trees was observed in the study area during fauna surveys, namely White Box (*Eucalyptus albens*).

Additional Koala food species to that scheduled in SEPP 44 are listed in the *NSW State Recovery Plan for the Koala* (DECC, 2008b). The study area is located within the Central Coast Koala Management Area where the following listed secondary food tree species were observed: Fuzzy box (*Eucalyptus conica*), Yellow Box (*E. melliodora*) and Grey Box (*E. moluccana*) (Hunter Eco, 2019).

General Koala scat searches were undertaken at several sites as part of diurnal and nocturnal fauna searches and are included in those times. Any potential Koala scats detected were firstly examined to see if they were composed of finely chewed Eucalyptus leaves and to help differentiate from Brush-tailed Possum (*Trichosurus vulpecula*) scats (which can be superficially similar). Where such scats were detected then they were sent to expert Barbara Triggs for further identification and targeted surveys using the *Spot Assessment Technique for determining localised levels of habitat use by Koalas* (Phillips and Callaghan, 2011), were carried out.

### **Searches for Reptiles and Amphibians – Active Searches, Pitfall Traps and Artificial Shelter Habitat**

Active searches were conducted at selected/preferred sites located at representative habitat components across the study area. This included potential shelter, refuge, foraging, over-wintering and breeding habitat for the range of species detected and searched for. Inspected habitat features included ground logs/timber, surface rock, cow pats, rock shelters, rock outcrops/crevices, decorticated bark, mature/old growth trees and stags with accessible crevices/fissures/hollows, culverts, dams, riparian zones (ponded sections of creeks and creek banks), soaks and man-made refuge habitats, where present, at each survey site and across the study area.

Further opportunistic searches including searches of other suitable microhabitat features encountered whilst traversing between survey plots – this approach targeted species known to have specific habitat/micro-habitat preferences not apparent within the survey plots chosen. Similarly, during road/track traverses (diurnal and nocturnal) scans were made for species that were active or more active at certain times of the day.

Surveys for amphibians were undertaken in accordance with the OEH *Hygiene Protocol for the Control of Disease in Frogs* (DECC, 2008a).

Initial habitat surveys were carried out for the threatened Green and Golden Bell Frog at 54 dams/ponds/drainage lines within the study area (Figure 6), searching for habitat features known to be favoured by this species (still water with some growth of Cumbungi (*Typha* sp) and/or other sedges such as *Phragmites australis*, *Juncus acutus*).

Targeted surveys were then carried out at 11 locations which contained at least some habitat features for this species. Surveys consisted of diurnal and/or nocturnal active searches for frogs and tadpoles, call-playback and listening (Figure 6) as follows:

- Location 31 – Quarry Pond: 16 November 2018 and 3-6 December 2018.
- Location 29 – Dam: 12-14 November 2018.
- Location 28 – Dam: 12-14 November 2018.
- Location 43 – Drainage Line: 5-6 and 19 December 2018.
- Location 32 – Pond Complex: 20 December 2018.
- Location 34 – Dam: 13 November 2018 and 20 December 2018.

## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

- Location 48 – Concrete Pond #1: 12 November 2018, 3 and 20 December 2018.
- Location 50 – Concrete Pond #2: 20 December 2018.
- Location 44 – Drainage Line with Ponds: 12 November 2018, 3 and 20 December 2018.
- Location 33 – Railway Loop Dams: 20 December 2018.
- Location 51 – Workshop Dam 1: 20 December 2018.
- Location 52 – Workshop Dam 2: 20 December 2018.

All ecologists recorded herpetofauna as they were encountered during the survey periods however the following information is in relation to the dedicated herpetofauna specialists (Henry Cook, Ross Wellington and Alex Dudley) on the ecological team.

Diurnal surveys were generally conducted between dawn and midday or until conditions became too hot. Evening surveys were generally run between dusk and 22:00-23:00 hours. The time spent on a site was determined by the habitat quality and the species to be targeted, so that survey effort was increased for higher quality sites compared with lower quality sites.

Specific targeted surveys were conducted for Striped Legless Lizard (*Delma impar*), Pink-tailed Legless Lizard (*Aprasia parapulchella*) and Pale-headed Snake (*Hoplocephalus bitorquatus*).

In January 2018 the total survey effort over the 6 days from the 22-27 January 2018 was 48.25 hours. Diurnal survey effort was 12.75 hours and nocturnal survey effort was 35.5 hours. The survey effort for each site in January 2018 is summarised below with a break down showing time spent between diurnal and nocturnal surveys.

**Table 12: Herpetofauna Survey Effort for January 2018 Survey Period**

Site	1	2	3	4	5	6	7	10	11	Totals
Diurnal (hrs)	3.75	1.50	2.25	1.50	1.00	0.50	0.50	1.00	0.75	<b>12.75</b>
Nocturnal (hrs)	20.00	7.50	1.00	1.00	Nil	2.00	2.00	Nil	2.00	<b>35.5</b>
<b>Totals (hrs)</b>	<b>23.75</b>	<b>9.00</b>	<b>3.25</b>	<b>2.5</b>	<b>1.00</b>	<b>2.5</b>	<b>2.5</b>	<b>1.00</b>	<b>2.75</b>	<b>48.25</b>

No habitat searches for reptiles and amphibians were carried out during the cooler June, August and September 2018 survey periods but if any species were incidentally encountered, they were recorded.

In November 2018 diurnal survey effort was at least 136.25 hours and nocturnal survey effort was at least 19 hours. The survey effort for each site in November 2018 is summarised below with a break down showing time spent between diurnal and nocturnal surveys.

**Table 13: Herpetofauna Survey Effort for November 2018 Survey Period**

Site	1	2	3	4	5	6	7	9	10	17	18	Incidental	Totals
Diurnal (hrs)	4.5	23	8	6	36	8	7.5	1	1.75	12	4.5	24	<b>136.25</b>
Nocturnal (hrs)	-	5.75	1.25	-	5.25	1.25	2.5	-	-	-	3	-	<b>19.00</b>
<b>Totals (hrs)</b>	<b>4.5</b>	<b>28.75</b>	<b>9.25</b>	<b>6</b>	<b>41.25</b>	<b>9.25</b>	<b>10</b>	<b>1</b>	<b>1.75</b>	<b>12</b>	<b>7.5</b>	<b>24</b>	<b>155.25</b>

In December 2018 diurnal survey effort was at least 84.25 hours and nocturnal survey effort was at least 16 hours. The survey effort for each site in December 2018 is summarised below with a break down showing time spent between diurnal and nocturnal surveys.

**Table 14: Herpetofauna Survey Effort for December 2018 Survey Period**

Site	1	3	4	6	7	10	11	13	14	16	17	18
Diurnal (hrs)	11.0	12.75	14.25	0.75	1.75	-	10.0	1.5	1.5	1	8.0	3.25
Nocturnal (hrs)	5	-	-	1.25	-	1.25	0.25	-	3.5	-	3.75	-
<b>Totals (hrs)</b>	<b>16</b>	<b>12.75</b>	<b>14.25</b>	<b>2.0</b>	<b>1.75</b>	<b>1.25</b>	<b>10.25</b>	<b>1.5</b>	<b>5.0</b>	<b>1</b>	<b>11.75</b>	<b>3.25</b>

Site	Savoy Dam	Incidental	Totals
Diurnal (hrs)	6.0	12.5	<b>84.25</b>
Nocturnal (hrs)	1.0	-	<b>16.00</b>
<b>Totals (hrs)</b>	<b>7.0</b>	<b>12.5</b>	<b>100.25</b>

During the November/December 2018 survey periods two additional reptile survey techniques were employed being Pitfall Traps and Artificial Shelter Habitat (see Figure 4 for locations).

Pitfall trap grids

Pitfall trap grids consisting of six 150 mm diameter, 600 mm deep end capped PVC pipes with 300 mm high drift fencing were deployed at Sites 2d, 3a and 5b for four days/nights from 19-23 November 2018 (Figure 4). Pits were deployed and placed 5 m apart in a straight line with a continuous run of drift fence placed along the pitfalls and extending for a few metres beyond the end pitfalls as per DEC (2004a). A piece of polystyrene foam together with leaf litter was placed in the bottom of each pit. In the event of inclement weather or when not in use, the top end of the pits was capped. In addition, at least two reptile net funnel traps were placed on each pitfall drift fence line. Pitfalls and funnels were typically checked at dawn, in the afternoon and after sunset and any animals identified and released.

Artificial shelter habitat

Artificial shelter habitat consisting of grids of second-hand terracotta roofing tiles (approximately 40 cm x 30 cm) were deployed at Sites 4a, 6, 7a and 17 to specifically target Striped Legless Lizard as per SEWPaC (2011c) (Figure 4). Arrays at Sites 6b, 7a and 17 consisted of 50 tiles, at 5 metre spacing between tiles, arranged in a grid of 10 tiles by five. 70 tiles were deployed at Site 4a. At each relevant site, tiles were placed in DNG adjacent to grassy woodland / open forest where grass cover was not too sparse. Wherever possible a northerly aspect was chosen. Artificial shelter habitat was deployed from 13 November to 23 December 2018 (39 days/nights) and checked one to two times a week during survey periods when ambient temperatures were not too high.

Targeted habitat surveys

Based on advice from the OEH, targeted habitat surveys for the Pink-tailed Legless Lizard were carried out by two researchers over two days from 30 April 2019 to 1 May 2019 to map rocky areas that provide potential habitat. As such, surveys specifically targeted areas of lightly imbedded surface rock within PCT 1606 and 1606 DNG (where the Pink-tailed Legless Lizard was recorded during the 2018 surveys) within the proposed underground mining area and associated surface disturbance areas (including within the vicinity of the proposed Edderton Road realignment). In order to create the species polygon, a 50 m zone was applied around the rocky areas, as request by OEH.

Areas of lightly imbedded surface rock were mapped within the target vegetation community type either on foot or via 4WD using a GPSKit device with an accuracy of +/- 5-10 m.

### **Opportunistic Observations**

All fauna observed or heard opportunistically during the field surveys (including travelling between sites in the broader area) were recorded. Characteristic signs, tracks, trails and other indirect evidence of fauna species from all fauna groups were also recorded. Any observed predator scats and/or owl pellets containing bone and fur material were collected and sent for analysis to expert Barbara Triggs (Genoa, Victoria).

### 2.3.3 Survey Effort

Table 15 provides a summary of the survey techniques and effort employed at each of the survey sites.

### 2.3.4 Limitations

Despite below average rainfall conditions (Section 2.3.1), a number of fauna surveys have been carried in the study area since the early 2000s and therefore it is likely that the fauna and habitats present (or potentially present) are well understood.

### 2.3.5 Nomenclature

Primary sources of literature accessed for nomenclature includes:

- *CSIRO list of Australian Vertebrates* (Clayton et al, 2006);
- *Birds - Systematics and Taxonomy of Australian Birds* (Christidis and Boles, 2008);
- *Mammals - The Mammals of Australia, Third Edition*, (Van Dyck and Strahan, 2008);
- *Bats - Australian Bats, Second Edition*, (Churchill, 2009) and *A current taxonomic list of Australian Chiropteran* (Reardon, Armstrong, and Jackson, 2015); and
- *Amphibians/Reptiles - Reptiles and Amphibians of Australia, Seventh Edition*, (Cogger, 2018).

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 15: Summary of Survey Techniques and Effort Used at Each Site within the Study Area

Site (Figure 4)	Habitat Survey (hrs)	Diurnal Bird Survey (hrs)	Nocturnal Bird Surveys (hrs)	Ground "A" Elliott Trapping (small) (trap nights)	Arboreal "B" Elliott Trapping (trap nights)	Cage Trapping (trap nights)	Hair Tubes (trap nights)	Camera Trapping (trap nights)	Nest boxes (trap nights)	Harp Trapping (trap nights)	Ultrasonic Bat Detection - Anabat (detector nights)	Habitat searches for cave roosting bats (hrs)	Nocturnal Call Playback (sessions)	Spotlighting (hrs)	Diurnal Searches for Reptiles and Amphibians (hrs)	Nocturnal Searches for Reptiles and Amphibians (hrs)(nights)	Pitfall Trapping (trap nights)	Artificial Shelters (trap nights)
1, 1a and 1b	2	25.45	10.5 (14.5*)	100	40	4	365	140	660	23	16.25	6	9	24.5	19.25	25 (>2 nights)	-	-
2, 2a, 2b/3b, 2c and 2d	2	29.2	9.8 (7.45*)	100	40	4	608	138	530	16	3	-	6	17	24.5	13.25 (4.00*) (>2 nights)	24	-
3, 3a, 2b/3b and 3c	2	26	1.8 (11.2*)	100	40	8	600	138	665		2	-	5	14	23	2.25 (10.75*) (>2 nights)	24	-
4 and 4a	2	22.25	3 (4.75*)	-	-	-	-	-	-	-	2	-	3	7.75	21.75	1 (6.75*) (>2 nights)	-	2730
5, 5a and 5b	2	43.5	12.5	-	40	-	8	4	-	16	2	-	7	21	37	5.25 (16.75*) (>2 nights)	24	-
6, 6a, 6b and 6c	2	17	4.25 (11*)	-	40	-	8	25	-	2	2	-	5	15.25	9.25	4.5 (10.75*) (>2 nights)	-	1950
7, 7a, 7b and 7c	2	14.5	17*	100	40	-	-	20	-	-	2	-	5	17	9.75	4.5 (12.5*) (>2 nights)	-	1950
8 and 8a	2	0.5	-	-	-	-	-	-	-	-	-	-	2	0.5	-	1.5* (1 night)	-	-
9	2	1.00	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-
10, 10a and 10b	2	3.75	5*	-	-	-	-	-	-	-	2.5	1	2	5	2.75	1.25 (3.75*) (2 nights)	-	-
11 and 11a	2	16.25	1.5 (4*)	-	-	-	-	-	-	-	1	-	4	5.5	10.75	2.25 (3.25*) (>2 nights)	-	-
12	2	1.50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
13	2	1.75	-	-	-	-	-	-	-	-	-	-	4	-	1.5	-	-	-
14a and 14b	2	4.00	4.5 (4*)	-	-	-	-	-	-	-	-	-	4	8.5	1.5	3.5 (5.00*) (>2 nights)	-	-
15 and 15a	2	7.50	1.5 (1.5*)	-	-	-	-	-	-	-	-	-	1	3	-	3* (1 night)	-	-
16, 16a, 16b and 16c	2	6.50	1.7 (1.55*)	-	-	-	760	-	-	-	-	-	1	3.5	1	3.25* (1 night)	-	-
17	2	20	3.75	-	-	-	-	30	-	-	-	3	2	3.75	20	3.75 (2 nights)	-	1950
18 and 18a	2	7.75	3 (3*)	-	40	-	360	-	-	-	-	-	5	6	7.75	3 (3*) (> 2 nights)	-	-
A mine dam (known as Savoy Dam)	2	6	1 (2*)	-	-	-	-	-	-	8	-	-	2	3	6	1 (2*) (2 nights)	-	-
Rocky Escarpment near corner Edderton Road and Golden Highway (IncB)	-	-	-	-	-	-	-	-	-	-	2	6	-	-	-	-	-	-
Incidental (including IncA)	-	36.5	-	-	-	-	-	266	-	-	1	-	-	-	36.5	-	-	-
<b>Totals</b>	<b>38</b>	<b>290.90</b>	<b>145.75</b>	<b>400</b>	<b>280</b>	<b>16</b>	<b>2,709</b>	<b>623</b>	<b>1,855</b>	<b>65</b>	<b>35.75</b>	<b>16</b>	<b>67</b>	<b>155.25</b>	<b>233.25</b>	<b>156.75</b>	<b>72</b>	<b>8,580</b>

\* Additional hours undertaken during activities other than the dedicated nocturnal bird surveys by Tony Saunders or dedicated herpetofauna nocturnal surveys by Henry Cook, Ross Wellington and Alex Dudley.

### 2.3.6 Targeted Searches for Threatened Fauna

Threatened fauna species listed under the BC Act and/or EPBC Act which are known or likely to occur in the study area were specifically targeted during the surveys (Table 16). Threatened fauna species were targeted in accordance with the survey timing, techniques and effort described within the relevant survey guidelines listed in Section 2.2.

Table 1 provides a list of threatened fauna species-specifically targeted during the surveys (although the surveys were designed to obtain an inventory of all native and introduced fauna species present not only the threatened species listed).

The following species listed in Table 1 did not have any potential habitat in the study area and are therefore not included in Table 16: Booroolong Frog (*Litoria booroolongensis*) and Green-thighed Frog (*Litoria brevipalmata*).

### 2.3.7 Species Credit Species Habitat Polygon Mapping

Species credit species habitat polygon maps have been produced in accordance with the BAM (OEH, 2017) and the '*Species Credit Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method*' (OEH, 2018).

Where a survey confirms the species is present or likely to use the habitat in the study area, a 'species polygon' is produced that shows the area of suitable fauna species habitat for a species credit species. Species polygons were prepared for the Pink-tailed Legless Lizard, Striped Legless Lizard, Squirrel Glider and Southern Myotis based on the results of the survey.

The 'species polygons' were mapped using a best available ortho-rectified aerial image and contain the suitable habitat features or habitat components associated with that species on the subject land. A GPS was used to confirm the location of the species polygon.

Hunter Eco (2019) undertook a paddock tree assessment for Squirrel Glider connectivity from woodland patches, conducted using maximum separation between canopies of 50 m potential gliding distance. To achieve this paddock trees were digitally extracted from enhanced high-resolution aerial imagery into a vector drawing. A Distance Network with maximum distance 50 m was applied to the extracted canopies and those trees were grouped that were within 50 m or less from each other, and similarly connected to woodland patches. Hunter Eco (2019) notes that a limitation of this approach was that all paddock trees were extracted irrespective of height. This then would include regrowth trees that were likely to be too short for a glider to attain maximum gliding distance from.



MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 16: Targeted Searches for Conservation Significant Fauna Species

Common Name	Scientific Name	Conservation Status			Survey Guideline Requirements	Survey Timing (OEH, 2019a)	Survey Techniques and Effort undertaken by Future Ecology
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>			
<b>Amphibians</b>							
Green and Golden Bell Frog	<i>Litoria aurea</i>	V	E	S	NSW (DECC, 2009): Combination of tadpole surveys, call surveys (this species has a distinctive call) and active searching both during the day and night. Small areas of habitat (less than 0.3 ha) should be surveyed for a minimum of one hour on three separate occasions during the species' activity period. Larger areas, which may include whole wetlands and lagoon margins, are more difficult to survey and require a minimum of three separate four-hourly searches during the species' activity period. Commonwealth guidelines (DEWHA, 2010a) similar to NSW but require 4 separate visits from September to March.	January to March, November and December	<b>The survey guideline requirements and timing were met.</b> Approximately 42 dams/ponds have been identified in the study area as occurring within or adjacent to PCTs 1598, 1604, 1606, 1691, 1692 and 1731 (Hunter Eco, 2019) (Figure 6). All were inspected in November/December 2018 and most were found to not provide suitable habitat for this species due to lack of fringing and aquatic vegetation. Four dams/ponds/drainage lines which did provide some potential habitat were intensively surveyed at least three times for at least one person hour on each occasion in accordance with DECC (2009) guidelines. This includes locations 28, 29, 31 and 43. Survey techniques included diurnal and nocturnal active searches and call-playback. Less intensive surveys involving 1-2 diurnal call-playback sessions were also carried out at locations 32, 33, 34, 44, 45, 48, and 49.
<b>Reptiles</b>							
Pink-tailed Legless Lizard	<i>Aprasia parapulchella</i>	V	V	S	Diurnal habitat searches (which includes overturning of rocks) in spring and early summer (SEWPaC, 2011b).	September to November	<b>The survey guideline requirements and timing were met.</b> Diurnal habitat searches (which included the overturning of rocks) were undertaken at several sites in November 2018 (Table 13). In addition to the above, pitfall trapping and placement of artificial shelter habitat in November 2018 could have also detected this species. Further, although outside the survey timing by OEH (2019a), diurnal habitat searches (which included the overturning of rocks) was also undertaken in January 2018 and December 2018 (Tables 12 and 14).

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 16 (Continued): Targeted Searches for Conservation Significant Fauna Species

Common Name	Scientific Name	Conservation Status			Survey Guideline Requirements	Survey Timing (OEH, 2019a)	Survey Techniques and Effort undertaken by Future Ecology
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>			
Striped Legless Lizard	<i>Delma impar</i>	V	V	S	NSW guidelines (DEC, 2004a): Pitfall trapping over 24 trap nights, preferably using six traps for a minimum of four consecutive nights per stratification unit up to 100 ha. Commonwealth Guidelines and Referral Guidelines (SEWPaC 2011a:b): Either artificial shelter sites or pitfall trapping as primary technique but artificial sites are preferred where there is surface rock. Habitat searches are secondary. Artificial shelter: up to 10 grids of 50 tiles on north facing slopes in habitat areas >30 ha as a minimum at least 50 pitfall configurations should be used for sites greater than 25 ha, using 2-5 pitfalls, 5m fence per configuration. Survey timing: September to May (SEWPaC, 2011b)	September to December	<b>The survey guideline requirements and timing were met.</b> This species was surveyed via multiple methods, namely pitfall trapping, artificial shelter sites and active searching. Pitfall trapping was undertaken over 24 trap nights, using six traps for four consecutive nights, at Sites 2d, 3a and 5b in November 2018 (Figure 4). In addition, four grids of 50 artificial shelter sites (total of 200) were installed at Sites 6a, 7a and 17, with 70 tiles deployed at Site 4a, in November 2018 and checked several times through December 2018 (Figure 4). As a secondary measure, diurnal and nocturnal active searches were carried out in November/December 2018 at several sites with suitable habitat including the above sites. Further, diurnal habitat searches were also undertaken in January 2018, within the survey timing recommended by SEWPaC (2011b).
Pale-headed Snake	<i>Hoplocephalus bitorquatus</i>	-	V	S	No species-specific requirement defined. General survey guidelines (DEC, 2004a) for nocturnally active snakes such as this species require thirty-minute searches on two separate nights per stratification unit.	January, February, March, November and December	<b>The survey guideline requirements and timing were met.</b> Diurnal and nocturnal habitat searches for reptiles (which targeted woodland habitat in close proximity to watercourses) were undertaken within general fauna survey sites 1, 2, 3, 4, 5, 6, 7, 10 and 11 and for at least 30 minutes on at least one day/night (some sites surveyed for two day/nights) in January 2018 (Table 12 and Figure 4). Additional nocturnal reptile searches for the Pale-headed Snake were undertaken at several sites with suitable habitat (hollow trees) within PCTs 1604, 1655, 1606, 1691 and 1692 and particularly focused on Sites 3 and 5 were conducted on at least two separate nights in November/December 2018. Total survey effort for most sites during warm seasons exceeded two nights of survey and 30 mins per session. Rain fell in January, November and December 2018 (Table 2).

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 16 (Continued): Targeted Searches for Conservation Significant Fauna Species

Common Name	Scientific Name	Conservation Status			Survey Guideline Requirements	Survey Timing (OEH, 2019a)	Survey Techniques and Effort undertaken by Future Ecology
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>			
<b>Birds</b>							
Freckled Duck	<i>Stictonetta naevosa</i>	-	V	E	No species-specific requirement defined. Diurnal bird surveys as per DEC (2004a) would be appropriate for these species.	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> Diurnal bird surveys were undertaken in areas of suitable habitat i.e. dams.
Australasian Bittern	<i>Botaurus poiciloptilus</i>	E	E	E	NSW (DEC, 2004a): No species-specific requirement defined. Diurnal and nocturnal surveys would be appropriate for this species. Federal (DEWHA, 2010c): Observation of targeted foraging habitat within wetlands in the early morning or early evening. Detection by sightings and unsolicited calls. Area searches in suitable habitat for sightings, nests, indicative footprints and feathers.	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> Quality wetland habitat is not present in study area. Diurnal and nocturnal surveys undertaken in dams in study area would have detected this species if it were to occur.
Black Falcon	<i>Falco subniger</i>	-	V	E	No species-specific requirement defined (DEC, 2004a), except for Red Goshawk where searches for its distinctive nest and area searches (80 hours over 10 days) is recommended (DEWHA, 2010c). Diurnal bird surveys would be appropriate for these species. Some species have seasonal survey requirements under BAM.	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> Diurnal bird surveys were undertaken, particularly in woodland habitat in close proximity to watercourses. The diurnal bird surveys exceeded 80 hours over 10 days. Raptor surveys from high points in study area would have detected these species, if present. Particular attention was given to searches for and inspection of raptor nests to determine use and what species were using them were undertaken. In accordance with BAM seasonal surveys for Square-tailed Kite were undertaken in January, September, November and December 2018, White-bellied Sea-eagle in August, September, November and December 2018, and Little Eagle in August and September 2018.
Square-tailed Kite	<i>Lophoictinia isura</i>	-	V	S/E		January, September, October, November and December	
White-bellied Sea-eagle	<i>Haliaeetus leucogaster</i>	Ma	V	S/E		July, August, September, October, November and December	
Spotted Harrier	<i>Circus assimilis</i>	-	V	E		None	
Red Goshawk	<i>Erythroriorchis radiatus</i>	V	CE	S		All year	
Little Eagle	<i>Hieraaetus morphnoides</i>	-	V	S/E		August, September and October	

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 16 (Continued): Targeted Searches for Conservation Significant Fauna Species

Common Name	Scientific Name	Conservation Status			Survey Guideline Requirements	Survey Timing (OEH, 2019a)	Survey Techniques and Effort undertaken by Future Ecology
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>			
Bush Stone-curlew	<i>Burhinus grallarius</i>	-	E	S	No species-specific requirement defined. Nocturnal bird surveys incorporating call-playback and spotlighting would be appropriate for this species.	All year	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> Numerous nocturnal bird surveys incorporating call-playback and spotlighting were carried out at various sites at various times of the year for this species.
Australian Painted Snipe	<i>Rostratula australis</i>	E	E	E	Area searches or transects; targeted stationary observations at dawn and dusk of suitable foraging locations within wetlands (DEWHA, 2010c).	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> Quality wetland habitat not present in study area. Diurnal and nocturnal surveys undertaken in dams in study area would have covered this species. Targeted stationary observations were generally not undertaken as no habitat for this species (e.g. wetlands, lakes, swamps and clay pans) is present within the study area.
Eastern Curlew	<i>Numenius madagascariensis</i>	CE	-	S/E	No species-specific requirement defined. Diurnal bird surveys would be appropriate for this species.	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> Suitable habitat (intertidal estuarine mudflats and saltmarsh) is not present in study area. Numerous diurnal bird surveys were carried out throughout the study area from January to December 2018.
Curlew Sandpiper	<i>Calidris ferruginea</i>	CE	E	S/E	No species-specific requirement defined. Diurnal bird surveys would be appropriate for this species.	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> Numerous diurnal bird surveys were carried out throughout the study area from January to December 2018 but suitable habitat (intertidal estuarine mudflats, saltmarsh) is not present in study area.
Glossy Black-cockatoo	<i>Calyptorhynchus lathami</i>	-	V	S/E	No species-specific requirement defined (DEC, 2004a). DEWHA (2010c) has some recommended survey techniques for Glossy Black Cockatoo (which is listed as nationally endangered in South Australia) and these techniques are also relevant to this species wherever it occurs. Diurnal surveys for all species would be appropriate together with searches for chewed Casuarina cones under trees for signs of Glossy Black Cockatoo (DEWHA, 2010c).	March, April, May, June, July and August	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> Numerous diurnal bird surveys were carried out throughout the study area from January to December 2018 including within the required seasonal survey periods for Glossy Black Cockatoo and Gang-gang Cockatoo. Chewed cones (Glossy Black Cockatoo) were searched for under casuarina trees whenever encountered. Wherever hollows suitable for nesting by Glossy Black Cockatoo or Gang-gang Cockatoo were detected they were searched (from the ground) for signs of nesting, feathers etc.
Gang-gang Cockatoo	<i>Callocephalon fimbriatum</i>	-	V	S/E		January, October, November and December	
Little Lorikeet	<i>Glossopsitta pusilla</i>	-	V	E		None	
Turquoise Parrot	<i>Neophema pulchella</i>	-	V	E		None	

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 16 (Continued): Targeted Searches for Conservation Significant Fauna Species

Common Name	Scientific Name	Conservation Status			Survey Guideline Requirements	Survey Timing (OEH, 2019a)	Survey Techniques and Effort undertaken by Future Ecology
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>			
Swift Parrot	<i>Lathamus discolor</i>	CE	E	S/E	DEWHA (2010c) recommends area searches or transect surveys of suitable habitat (20 hours over 8 days), preferably in the early morning and afternoon when birds are most active and vocal. Detection by sighting or call. Slow-moving vehicle transects also effective in expansive areas, detecting loud, distinctive 'clinking' call that can be heard over noise of engine. Targeted surveys of patches of heavily flowering eucalypts may be useful.  Timing: surveys on the mainland should be conducted between March and July (DEWHA, 2010c).	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b>  Specific diurnal surveys for this species were carried out in June and August 2018 targeting patches of flowering box trees at that time. In June 23.75 hours were spent on diurnal surveys over 4 days in August 14 hours was spent over 3 days (i.e. a total of 36.75 hours over 7 days).
Eastern Grass Owl	<i>Tyto longimembris</i>	-	V	E	DEC (2004a) requires nocturnal surveys for all species incorporating quiet listening, spotlighting and call-playback. A minimum of five visits per site is suggested for Powerful Owl and Barking Owl and eight visits for Masked Owl. DEC (2004a) states that the surveys can occur any time of year.	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b>  At least five nocturnal surveys per relevant site were carried out during the survey timing (OEH, 2019a), with additional sessions conducted outside of the OEH (2019a) survey period (May to August) but within the DEC (2004a) survey period (any time of year). A total of 69 call playback sessions were undertaken (Table 15).  Wherever hollows suitable for nesting were detected they were searched (from the ground) for signs of nesting, feathers, pellets, remains of prey etc.
Masked Owl	<i>Tyto novaehollandiae</i>	-	V	S/E		May, June, July and August	
Powerful Owl	<i>Ninox strenua</i>	-	V	S/E		May, June, July and August	
Barking Owl	<i>Ninox connivens</i>	-	V	S/E		May, June, July, August, September, October, November and December	
Brown Treecreeper (eastern subspecies)	<i>Climacteris picumnus victoriae</i>	-	V	E	No species-specific requirement defined. Diurnal bird surveys would be appropriate for this species.	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b>  Numerous diurnal bird surveys were carried out throughout the study area from January to December 2018.
Speckled Warbler	<i>Chthonicola sagittata</i>	-	V	E		None	

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 16 (Continued): Targeted Searches for Conservation Significant Fauna Species

Common Name	Scientific Name	Conservation Status			Survey Guideline Requirements	Survey Timing (OEH, 2019a)	Survey Techniques and Effort undertaken by Future Ecology
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>			
Black-chinned Honeyeater (eastern subspecies)	<i>Melithreptus gularis gularis</i>	-	V	E	No species-specific requirement defined. Diurnal bird surveys would be appropriate for this species.	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> Numerous diurnal bird surveys were carried out throughout the study area from January to December 2018.
Regent Honeyeater	<i>Anthochaera phrygia</i>	CE	CE	S/E	Diurnal bird surveys undertaken for 20 hours over 10 days in areas of less than 50 ha (DEWHA, 2010c). The species is most conspicuous in the breeding season (primarily between September and November) (DEWHA, 2010c). Targeted searches of woodland patches with heavily flowering trees may be useful as well as call playback (DEWHA, 2010c).	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> At least 20 hours per site of diurnal surveys for this species were carried out at several sites in September, November and December 2018 (an excess of 10 days). Numerous diurnal bird surveys were also completed outside of the required survey timing period.
Painted Honeyeater	<i>Grantiella picta</i>	V	V	E	No species-specific requirement defined. Diurnal bird surveys would be appropriate for this species.	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> Numerous diurnal bird surveys were carried out throughout the study area from January to December 2018.
Hooded Robin (south-eastern form)	<i>Melanodryas cucullata cucullata</i>	-	V	E		None	
Flame Robin	<i>Petroica phoenicea</i>	-	V	E		None	
Scarlet Robin	<i>Petroica boodang</i>	-	V	E		None	
Grey-crowned Babbler (eastern subspecies)	<i>Pomatostomus temporalis temporalis</i>	-	V	E		None	
Varied Sittella	<i>Daphoenositta chrysoptera</i>	-	V	E		None	
Dusky Woodswallow	<i>Artamus cyanopterus cyanopterus</i>	-	V	E		None	
Diamond Firetail	<i>Stagonopleura guttata</i>	-	V	E		None	

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 16 (Continued): Targeted Searches for Conservation Significant Fauna Species

Common Name	Scientific Name	Conservation Status			Survey Guideline Requirements	Survey Timing (OEH, 2019a)	Survey Techniques and Effort undertaken by Future Ecology
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>			
<b>Mammals</b>							
Spotted-tailed Quoll	<i>Dasyurus maculatus maculatus</i> (south-eastern mainland population)	E	V	E	Habitat surveys (for potentially suitable habitat resources and signs of activity, scats and latrines), hair tubes and camera trapping (SEWPaC, 2011a). May to August is the optimal survey period for this species (SEWPaC, 2011a). 40 hair tubes (funnels) per 100 ha sampling units spread 100m apart from May to August (SEWPaC, 2011a).	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> A total of 5,004 hair tube trap nights and 623 camera trap nights were completed over the study area from January to June and November to December 2018. A total of 16 cage and 280 arboreal Elliott B trap nights were carried out in January and November 2018. A total of 1,855 nest box nights were carried out from January to June 2018. Numerous nocturnal spotlighting surveys were carried out throughout the study area. Predator scats were collected and analysed whenever detected.
Brush-tailed Phascogale	<i>Phascogale tapoatafa</i>	-	V	S	No species-specific requirement defined. Nocturnal spotlighting surveys, wildlife cameras, nest boxes, hair tubes and arboreal Elliott trapping would all be appropriate for this species.	All year	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> The survey techniques and effort discussed above for Spotted-tail Quoll would also have covered this species.
Common Planigale	<i>Planigale maculata</i>	-	V	S	No species-specific requirement defined. Pitfall trapping, artificial shelter habitat, nocturnal spotlighting surveys, wildlife cameras, hair tubes and Elliott trapping would all be appropriate for this species.	All year	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> The survey techniques and effort discussed above for Spotted-tail Quoll would also have covered this species. In addition, a total of 24 nights of pitfall trapping took place at each of 3 sites in the study area (total of 72 trap nights). Also, a total of 400 Elliott A terrestrial trap nights were carried out. This species may also have been potentially detected via the artificial shelter habitats (roofing tiles).
Koala	<i>Phascolarctos cinereus</i>	V	V	S/E	Federal guidelines (Commonwealth of Australia, 2014) discuss direct and indirect methods depending on density of animals and purpose of study. For low density populations indirect methods (signs) are recommended including searches for scratchings, scats etc. For higher density populations call-playback, spotlighting, wildlife cameras and other methods are recommended. Direct observation surveys should be undertaken between August and January (Commonwealth of Australia, 2014).	All year	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> Numerous spotlighting and call-playback sessions were undertaken across the study area from January to December 2018 including during the recommended target period between August and January. A total of 623 camera trap nights were completed over the study area from January to December 2018. Searches for scratchings on tree trunks and scats were undertaken as part of general ecological surveys on a daily basis.

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 16 (Continued): Targeted Searches for Conservation Significant Fauna Species

Common Name	Scientific Name	Conservation Status			Survey Guideline Requirements	Survey Timing (OEH, 2019a)	Survey Techniques and Effort undertaken by Future Ecology
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>			
Eastern Pygmy-possum	<i>Cercartetus nanus</i>	-	V	S	No species-specific requirement defined. Pitfall trapping, hair tubes, nest boxes, nocturnal spotlighting surveys, wildlife cameras, and Elliott trapping would all be appropriate for this species.	January, February, March, October, November and December	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> The survey techniques and effort discussed above for Spotted-tail Quoll would also have covered this species. In addition, a total of 24 nights of pitfall trapping took place at each of 3 sites in the study area (total of 72 trap nights). Also, a total of 400 Elliott A terrestrial trap nights were carried out.
Yellow-bellied Glider	<i>Petaurus australis</i>	-	V	E	No species-specific requirement defined. Hair tubes, nest boxes, nocturnal spotlighting and call-playback surveys, wildlife cameras, and Elliott trapping would all be appropriate for these species.	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> The survey techniques and effort discussed above for Spotted-tail Quoll would also have covered these species. In addition, numerous call-playback surveys were conducted across the study area over the survey period.
Squirrel Glider	<i>Petaurus norfolcensis</i>	-	V	S		All year	
Greater Glider	<i>Petauroides volans</i>	V	-	S		All year	
Brush-tailed Rock-wallaby	<i>Petrogale penicillata</i>	V	E	S	Thorough daytime searches for signs and habitat resources are considered an adequate form of survey method for detecting the brush-tailed rock wallaby, as long as all suitable rocky habitat including mid-level ledges and holes are inspected for signs of activity (SEWPaC, 2011a). Camera traps are also considered to be suitable (SEWPaC, 2011a).	All year	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> The rocky hill at Site 1 and the small escarpment near the main entrance off Edderton Road / The Golden Highway provide the only potential (but sub-optimal) habitat for this species in the study area. They were thoroughly searched for signs of this species on a number of occasions. Camera traps were also used.
Grey-headed Flying-fox, Little Bentwing-bat, Eastern Bentwing-bat, Large-eared Pied Bat, Southern Myotis, Eastern Cave Bat	-	-	-	-	Refer to Table 11.	-	Refer to Table 11.



MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 16 (Continued): Targeted Searches for Conservation Significant Fauna Species

Common Name	Scientific Name	Conservation Status			Survey Guideline Requirements	Survey Timing (OEH, 2019a)	Survey Techniques and Effort undertaken by Future Ecology
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>			
Yellow-bellied Sheathtail bat, Eastern Freetail-bat, Northern Freetail-bat, Corben's Long-eared Bat, Eastern False Pipistrelle, Greater Broad-nosed Bat	Various	-	-	E	Harp trapping and ultrasonic bat detection (Anabat) (DEC, 2004a; DEWHA, 2010b)	Oct to March	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> A total of 69 harp trap nights and 34 Anabat detection nights were undertaken (Table 15).
New Holland Mouse	<i>Pseudomys novaehollandiae</i>	V	-	E	No species-specific requirement defined. Pitfall trapping, artificial shelter habitat, nocturnal spotlighting surveys, wildlife cameras, hair tubes and Elliott trapping would all be appropriate for this species.	None	<b>The survey guideline requirements were met and surveys undertaken at an appropriate time.</b> The survey techniques and effort discussed above for Spotted-tail Quoll would also have covered this species. In addition, a total of 24 nights of pitfall trapping took place at each of 3 sites in the study area (total of 72 trap nights). Also, a total of 400 Elliott A terrestrial trap nights were carried out. This species may also have been potentially detected via the artificial shelter habitats (roofing tiles).

<sup>1</sup> Threatened species status under the EPBC Act (current as at March 2019). V = Vulnerable, E = Endangered, CE = Critically Endangered.

<sup>2</sup> Threatened species status under the BC Act (current as at March 2019). V = Vulnerable, E = Endangered, CE = Critically Endangered.

<sup>3</sup> Biodiversity credit class under the *Threatened Biodiversity Data Collection* (OEH, 2019a) (current as at March 2019), E = Ecosystem, S = Species.

## 3 Survey Results

### 3.1 Fauna Habitat Types

Vegetation within the study area was described and mapped by Hunter Eco (2019). Broad fauna habitat types in the study area have been described and mapped on Figures 7a and 7b based on the vegetation mapping by Hunter Eco (2019) and include:

- Dry Sclerophyll Forest.
- Grassy Woodlands.
- Forested Wetlands.
- Derived Native Grassland (DNG).
- Planted Trees.
- Cultivation.
- Waterbody/Dam.
- Woodland Rehabilitation.
- Pasture Rehabilitation.
- Infrastructure/Cleared Land.

The broad fauna habitat types are described below while a summary of habitat features, habitat types and dominant flora species observed at each survey site is presented in **Appendix B**.

Livestock grazing has effectively removed the shrub layer from parts of the study area and suppressed the ground-cover layer. The growth and flowering of groundcover species observed in November and December 2018 surveys following sporadic rainfall and removal of cattle was very noticeable.


Some relatively small areas of scattered surface rock were observed throughout the study area including at most survey sites. Observed surface rock was generally small (<50 cm diameter) and rocks were mostly located directly on soil rather than rock on rock. The exception to this was at Sites 1 and IncB (Figure 4). A rocky rise and associated escarpment were located at Site 1 with a smaller escarpment located at Site IncB. An old quarry was located at Site 1b with some exposed vertical rock faces, quarried rock and spoil.

#### **Dry Sclerophyll Forest**

The dry sclerophyll forest in the study area comprises Red Gum - Ironbark - Apple shrubby woodland (PCT 1607), White Box - Ironbark - Red Gum shrubby forest (PCT 1606) and Slaty Box shrubby woodland (PCT 1655) (Hunter Eco, 2019).

This habitat type consisted of remnant patches of native dominated vegetation containing a Eucalypt dominated canopy (Figure 7a).

The common canopy species included White Box (*Eucalyptus albens*), Blakely's Red Gum (*E. blakelyi*), Bulloak (*Allocasuarina luehmannii*) and Slaty Gum (*E. dawsonii*). If a sub-canopy was present it was generally patchy and dominated by Bulloak (*Allocasuarina luehmannii*) and/or Kurrajong (*Brachychiton populneus*).




- LEGEND**
- Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - 1 Dry Sclerophyll Forest
  - 2 Grassy Woodland
  - 3 Forested Wetland
  - 4 Derived Native Grassland
  - A Planted Trees
  - B Cultivation
  - C Waterbody/Dam
  - D Woodland Rehabilitation
  - E Pasture Rehabilitation
  - F Infrastructure/Cleared Land

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019);  
 Orthophoto Mosaic: 2018, 2016, 2011

**MALABAR COAL**  
 MAXWELL PROJECT  
 Broad Fauna Habitat Types  
 - Maxwell Underground

**Figure 7a**



- LEGEND**
- Biodiversity Assessment Development Footprint
  - 1 Dry Sclerophyll Forest
  - 2 Grassy Woodland
  - 3 Forested Wetland
  - 4 Derived Native Grassland
  - C Waterbody/Dam
  - D Woodland Rehabilitation
  - E Pasture Rehabilitation
  - F Infrastructure/Cleared Land

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019);  
 Orthophoto Mosaic: 2018, 2016, 2011

  
**MAXWELL PROJECT**  
**Broad Fauna Habitat Types**  
**- Maxwell Infrastructure**

**Figure 7b**

Where a shrub layer was present it was generally very sparse with common species being Cooba (*Acacia salicina*), Mock Olive (*Notelaea spp.*), Blackthorn (*Bursaria spinosa*) and the weed species African Boxthorn (*Lycium ferocissimum*).

The groundcover layer was typically sparse. It was typically co-dominated by a variety of grass and forb species depending on the site. Common species included Slender Bamboo Grass (*Austrostipa verticillata*), Purple Wire-grass (*Aristida ramosa*), and Small-leaf Bluebush (*Maireana microphylla*).

Generally, this habitat type contained areas of more complex vegetation and a greater number of fauna habitat features such as leaf litter, fallen timber, hollow logs, hollow-bearing trees and dead trees.

Dead trees (stags) and hollow-bearing trees were present at most sites but generally in low relative abundance.

All sites within this habitat type exhibited some degree of disturbance as a result of agricultural activities; including grazing, clearing, removal of fallen timber, removal or thinning of shrub and sub-canopy layers. Generally, the connectivity for this habitat type was low to moderate across the study area due to historical clearing to support agriculture.

### **Grassy Woodlands**

The grassy woodlands in the study area comprises Bull Oak grassy woodland (PCT 1692), Yellow Box - Apple grassy woodland (PCT 1693), Fuzzy Box woodland (PCT 201), Ironbark - Grey Box grassy woodland (PCT 1691), Weeping Myall woodland (PCT 116), and Grey Box - Spotted Gum - Narrow-leaved Ironbark woodland (PCT 1604) (Hunter Eco, 2019).

This habitat type consisted of remnant patches of native dominated vegetation which contained a more spread out Eucalypt dominated canopy and a grass/forb dominated understorey with fewer shrubs.

The common canopy species included Narrow-leaved Ironbark (*Eucalyptus crebra*), Grey Box (*E. moluccana*), Bulloak, Yellow Box (*E. melliodora*), Spotted Gum (*Corymbia maculata*) and Fuzzy Box (*E. conica*).

The groundcover layer is typically sparse. It is typically co-dominated by a variety of grass and forb species depending on the site. Common species include Slender Bamboo Grass (*Austrostipa verticillata*), Purple Wire-grass (*Aristida ramosa*), and Small-leaf Bluebush (*Maireana microphylla*).

Generally, this habitat type contained a greater number of fauna habitat features such as leaf litter, fallen timber, hollow logs, hollow-bearing trees, dead trees, and areas of more complex vegetation.

Dead trees (stags) and hollow-bearing trees were present at most sites but generally in low relative abundance.

All sites within this habitat type showed some degree of disturbance including grazing, clearing, removal of fallen timber, removal or thinning of shrub and sub-canopy layers. Generally, the connectivity for this habitat type was low to moderate across the study area which has been heavily cleared for agricultural.

### **Derived Native Grassland**

This habitat type consists of open grassy areas between patches of Dry Sclerophyll Forest and Grassy Woodlands. Cover was sparse to moderately dense. Typical species included Slender Bamboo Grass, Purple Wire-grass, and Small-leaf Bluebush.

Fauna habitat features were generally poor within this habitat type as it only provided open areas for some species. Generally fallen timber, hollow logs etc. were absent.

This habitat type was probably formed by and subject to ongoing disturbance from grazing and other agricultural practices.

Connectivity in this habitat type was generally moderate to high as it formed large connected areas across the study area landscape.

Scattered or remnant trees (paddock trees) were occasionally present in this broad fauna habitat type.

It was found adjacent to or interspersed with most sites and was the most abundant habitat type within the study area.

### **Forested Wetlands**

The forested wetland comprises Swamp Oak forest (PCT 1731) and Hunter Lowland Red Gum Forest (PCT 1598) (Hunter Eco, 2019). This habitat type was associated with parts of the Saddlers Creek drainage line and consisted of a mixture of riparian remnant old growth trees and Swamp Oak (*Casuarina glauca*) regeneration (Figure 7a).

The common canopy species included Swamp Oak (*Casuarina glauca*), Bulloak (*Allocasuarina luehmannii*), Rough-barked Apple (*Angophora floribunda*) and Yellow Box (*Eucalyptus melliodora*). If a sub-canopy was present it was dominated by Swamp Oak, Bulloak and/or Cooba (*Acacia salicina*). If a shrub layer was present it typically included Silver Cassia (*Senna artemisioides*), Small-leaf Bluebush (*Maireana microphylla*) and the weeds African Boxthorn (*Lycium Ferocissimum*) and Common Prickly Pear (*Opuntia stricta*).

The groundcover layer was mostly sparse. It was typically co-dominated by a variety of grass and forb species depending on the site. Common species included *Aristida ramosa*, *Austrostipa verticillata*, *Cynodon dactylon* and *Phalaris aquatica*.

It was in poor condition due to sparse cover, grazing pressure, erosion and dry conditions.

It provided few tree hollows, fallen logs and dead standing trees.

It was poorly connected except along watercourse where the riparian zone varies in thickness and quality and with some open sections with little woody cover.

### **Planted Trees**

This occurred as a narrow 20 m wide strip of planted trees along the frontage of the study area with the Golden Highway near the Edderton Road intersection (Figure 7a). The dominant planted tree was *Eucalyptus dawsonii* (Slaty Gum) (Hunter Eco, 2019).

Habitat value is low due to narrow area of planting, young age of planted trees and lack of features such as tree hollows, dead standing tree and fallen logs.

### **Cultivation**

There was an area of cultivated land west of Edderton Road (Figure 7a). It contained no native tree or shrub species and has little habitat value.

### **Waterbody/Dam**

There were 52 waterbodies/dams within the study area with most being relatively small farm dams associated with grazing activity (Figures 7a and 7b). There were several larger waterbodies associated with old mining pits and dams associated with the previous Drayton open cut coal mine.

Saddlers Creek and Saltwater Creek and their tributaries were located in the southern part of the study area. They appeared to have no permanently flowing water and only a few small ponds observed at some locations during the survey periods.

Most waterbodies/dams lacked aquatic vegetation and/or dense fringing vegetation and provided lower quality habitat for most species. Grazing has degraded most of the dams and suppressed growth of aquatic plants and fringing sedges.

### **Woodland Rehabilitation**

The location of woodland rehabilitation of previous open cut mining areas at the Maxwell Infrastructure is shown on Figure 7b.

Hunter Eco (2019) describes the woodland rehabilitation as comprising a low native species diversity and high weed diversity including five high threat weed species; native species were present in low numbers. Native canopy species were *Acacia saligna*, *Eucalyptus cladocalyx* (Sugar Gum) and a variety of Acacia species.

The woodland rehabilitation provides lower quality fauna habitat compared to intact woodland sites (Malabar, 2018), except perhaps movement and foraging habitat for more mobile species.

### **Pasture Rehabilitation**

The location of pasture rehabilitation of previous open cut mining areas at the Maxwell Infrastructure is shown on Figure 7b.

Hunter Eco (2019) describes the pasture rehabilitation as comprising of a low native flora species diversity and high weed diversity including eight high threat weed species; native species were present in low numbers. *Melinis repens* (Red Natal Grass) was the dominant weed species and *Hyparrhenia hirta* (Coolatai Grass) was the dominant high threat weed species.

The pasture rehabilitation provides lower quality fauna habitat compared to intact woodland sites (Malabar, 2018), except perhaps movement and foraging habitat for more mobile species.

### **Infrastructure/Cleared Land**

Cleared land and infrastructure areas associated with the Maxwell Infrastructure are shown on Figures 7a and 7b.

## 3.2 Fauna Species

A total of 227 fauna species were recorded in the study area during the 2018 surveys including 8 amphibian, 22 reptile, 148 bird, and 49 mammal species. 12 of the recorded species were exotics.

**Appendix A** contains the full list of fauna species recorded during the survey periods.

## 3.3 Threatened Fauna Species Listed under the BC Act

### 3.3.1 Threatened Fauna Species Recorded During the Surveys

A total of 25 threatened fauna species listed under the BC Act (all listed as vulnerable) were recorded by Future Ecology in the study area during the current surveys, as well as the Square-tailed Kite which was observed during vegetation surveys (Colin Driscoll, Hunter Eco, pers. obs.) (Table 17). Five of the species recorded by Future Ecology are listed under the EPBC Act (all listed as vulnerable) (Table 17).

**Table 17: Threatened Fauna Species Recorded During this Study**

Common Name	Scientific Name	Conservation Status		Credit Class <sup>3</sup>
		BC Act	EPBC Act <sup>2</sup>	
<b>Reptiles</b>				
Pink-tailed Legless Lizard	<i>Aprasia parapulchella</i>	V	V	S
Striped Legless Lizard	<i>Delma impar</i>	V	V	S
<b>Birds</b>				
Square-tailed Kite	<i>Lophoictinia isura</i>	V	-	E*
White-bellied Sea-eagle	<i>Haliaeetus leucogaster</i>	V	-	E*
Spotted Harrier	<i>Circus assimilis</i>	V	-	E
Little Eagle	<i>Hieraaetus morphnoides</i>	V	-	E*
Glossy Black-Cockatoo	<i>Calyptorhynchus lathami</i>	V	-	E*
Little Lorikeet	<i>Glossopsitta pusilla</i>	V	-	E
Brown Treecreeper (eastern subspecies)	<i>Climacteris picumnus victoriae</i>	V	-	E
Speckled Warbler	<i>Chthonicola sagittata</i>	V	-	E
Black-chinned Honeyeater (eastern subspecies)	<i>Melithreptus gularis gularis</i>	V	-	E
Painted Honeyeater	<i>Grantiella picta</i>	V	V	E
Flame Robin	<i>Petroica phoenicea</i>	V	-	E
Scarlet Robin	<i>Petroica boodang</i>	V	-	E
Grey-crowned Babbler (eastern subspecies)	<i>Pomatostomus temporalis temporalis</i>	V	-	E
Varied Sittella	<i>Daphoenositta chrysoptera</i>	V	-	E
Dusky Woodswallow	<i>Artamus cyanopterus cyanopterus</i>	V	-	E
<b>Mammals</b>				
Squirrel Glider	<i>Petaurus norfolcensis</i>	V	-	S
Grey-headed Flying-fox	<i>Pteropus poliocephalus</i>	V	V	E*
Yellow-bellied Sheath-tail-bat	<i>Saccolaimus flaviventris</i>	V	-	E
Eastern Freetail-bat	<i>Mormopterus norfolkensis</i>	V	-	E
Little Bentwing-bat	<i>Miniopterus australis</i>	V	-	E*
Eastern Bentwing-bat	<i>Miniopterus schreibersii oceanensis</i>	V	-	E*
Large-eared Pied Bat	<i>Chalinolobus dwyeri</i>	V	V	S^
Southern Myotis	<i>Myotis macropus</i>	V	-	S

<sup>1</sup> Conservation status under the BC Act (current as at March 2019). V = Vulnerable.

<sup>2</sup> Conservation status under the EPBC Act (current as at March 2019). V = Vulnerable.

<sup>3</sup> Biodiversity credit class under the *Threatened Biodiversity Data Collection* (OEH, 2019a) (current as at March 2019). E = Ecosystem, S = Species

\* This species is a dual credit species, however, no breeding habitat was recorded and therefore it is considered an ecosystem credit species in the study area.

^ This species is a species credit species, however, no breeding habitat was recorded and therefore no species credits would be generated from the study area.



Figures 8 to 11 show the locations of threatened fauna species records.

Four of the species in Table 17 are considered 'species credit species' in the study area (Figure 12), namely:

- Pink-tailed Legless Lizard;
- Striped Legless Lizard;
- Squirrel Glider; and
- Southern Myotis.

Figure 13 shows the locations of threatened fauna species listed under the EPBC Act.

All of the species in Table 17 are discussed below.

### **Pink-tailed Legless Lizard (*Aprasia parapulchella*)**

This species is listed as 'vulnerable' under the BC Act and EPBC Act. It is classified as a 'Species Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

This small, distinctive legless lizard lives beneath lightly embedded rocks on sloping, generally well-drained soils in derived grasslands and grassy woodlands (Wong et al., 2011; OEH, 2019c). It has a specialised diet and lives in the burrows of small ants; the eggs and larvae of which it relies on for food (Wong et al., 2011). It currently has a patch-work distribution along the foothills of the western slopes of the Great Dividing Range between Bendigo in Victoria and Gunnedah in NSW with populations mostly fragmented and isolated from one another (Wong et al., 2011). The population density is low except in the most favourable habitats (Wong et al., 2011), with the most abundant populations known from the Australian Capital Territory (ACT) and in NSW between Yass and Cooma (ACT Government, 2017a).

This species has not been previously recorded in the study area (Table 1). A single adult specimen of Pink-tailed Legless Lizard was found beneath a rock in grassland at Site 5 in November 2018 (Figure 8; Plate 1). In addition, a single slough (shed skin) of the same species was also found beneath a rock in the same area. It was not recorded in any pitfall traps or artificial shelter habitat.


This represents the first record of this species in the Muswellbrook LGA and represents an eastern range extension of its current known distribution with the closest record being from Goulburn River National Park about 80 km to the west of the study area and dated 2000 (OEH, 2019b).

The vegetation where this species was found on site is White Box - Ironbark - Red Gum shrubby forest PCT1606 - Derived Native Grassland (Hunter Eco, 2019).

There are no PCTs published in the *Threatened Biodiversity Data Collection* (OEH, 2019a) as being associated with this species within the Sydney Basin – Hunter IBRA sub-region. The *Threatened Biodiversity Data Collection* (OEH, 2019a) describes a habitat constraint for this species as rocky areas or within 50 m of rocky areas (Table 4).

Throughout its range this species generally occupies sites with a grassy ground layer with little or no leaf litter, and relatively low tree and shrub cover (Wong et al., 2011).

This species can be hard to detect and may only occur in low numbers across the study area. It is most likely to occur in similar grassy habitat to that found at Site 5 at locations with a good cover of native grasses and forbs as well as lightly-embedded rocks. It is unlikely to be found at any locations where the ground layer vegetation has been subject to pasture improvement and/or heavy stock concentrations (Wong et al., 2011).




- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
- Threatened Species**
- ▲ Green and Golden Bell Frog
  - Pink-tailed Legless Lizard
  - Striped Legless Lizard

Reference:  
 13. OEH (2019)  
 15. Future Ecology (2019)  
 Note: There are no references 1 - 12 and 14 on this figure.

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011



**Figure 8**




- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
- Threatened Species**
- Black Falcon
  - Square-tailed Kite
  - White-bellied Sea-Eagle
  - Spotted Harrier
  - Little Eagle
  - Bush Stone-curlew
  - Glossy Black-Cockatoo
  - Little Lorikeet
  - Swift Parrot
  - Barking Owl
  - Brown Treecreeper (eastern subspecies)
  - Speckled Warbler
  - Black-chinned Honeyeater (eastern subspecies)
  - Painted Honeyeater
  - Hooded Robin (south-eastern form)
  - Flame Robin
  - Scarlet Robin
  - Grey-crowned Babbler (eastern subspecies)
  - Varied Sittella
  - Dusky Woodswallow
  - Diamond Firetail

- Reference:**
- |                               |                                     |
|-------------------------------|-------------------------------------|
| 1. Ecotone (2000)             | 13. OEH (2019)                      |
| 2. Umwelt (2005)              | 14. Hunter Eco (pers. comm. (2019)) |
| 3. Umwelt (2006b)             | 15. Future Ecology (2019)           |
| 5. Umwelt (2007b)             | 16. Eco Logical Australia (2014)    |
| 6. Cumberland Ecology (2009a) | 17. Eco Logical Australia (2015)    |
| 7. Cumberland Ecology (2010)  | 18. Eco Logical Australia (2016a)   |
| 8. Cumberland Ecology (2012)  | 20. Eco Logical Australia (2016b)   |
| 12. Birdlife Australia (2017) |                                     |
- Note: There are no references 4, 9-11 and 19 on this figure.

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019);  
 MSEC (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011

**Figure 9**



- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
- Threatened Species**
- ▲ Spotted-tailed Quoll
  - △ Brush-tailed Phascogale
  - ▲ Koala
  - ▲ Squirrel Glider


**Reference:**

- 5. Umwelt (2007b)
- 6. Cumberland Ecology (2009a)
- 7. Cumberland Ecology (2010)
- 9. HVEC (pers. Comm. (2012))
- 13. OEH (2019)
- 15. Future Ecology (2019)
- 16. Eco Logical Australia (2014)
- 18. Eco Logical Australia (2016a)
- 19. Eco Logical Australia (2016b)

*Note: There are no references 1 - 4, 8, 10 - 12, 14 and 17 on this figure.*

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011

**Figure 10**



**LEGEND**

- Maxwell Project Exploration Licence Boundary
- Maxwell Project Mining and Coal Lease Boundary
- Indicative Extent of Underground Development
- Biodiversity Assessment Development Footprint
- Ecology Study Area
- Existing Conservation/Offset Area

**Threatened Species**

- ▽ Grey-headed Flying-fox
- ▽ Yellow-bellied Sheath-tail-bat
- ▽ Northern Freetail-bat
- ▽ Eastern Freetail-bat
- ▽ Eastern Bentwing-bat
- ▽ Corben's Long-eared Bat
- ▽ Large-eared Pied Bat
- ▽ Little Bentwing-bat
- ▽ Eastern False Pipistrelle
- ▽ Southern Myotis
- ▽ Greater Broad-nosed Bat
- ▽ Eastern Cave Bat

**Reference:**


1. Ecolone (2000)
3. Umwelt (2006b)
4. Umwelt (2007a)
6. Cumberland Ecology (2009a)
8. Cumberland Ecology (2012)
11. Niche (2012)
13. OEH (2019)
15. Future Ecology (2019)
16. Eco Logical Australia (2014)
17. Eco Logical Australia (2015)
18. Eco Logical Australia (2016a)
19. Eco Logical Australia (2016b)
20. Eco Logical Australia (2017)

*Note: There are no references 2, 5, 7, 9, 10, 12 and 14 on this figure. (Some records by Niche (2012) were possible/probable records).*

*Source: © NSW Department of Planning and Environment (2019); NSW Department of Finance, Services & Innovation (2019) Orthophoto Mosaic: 2018, 2016, 2011*



**Figure 11**



**LEGEND**

- Maxwell Project Exploration Licence Boundary
- Maxwell Project Mining and Coal Lease Boundary
- Indicative Extent of Underground Development
- Biodiversity Assessment Development Footprint
- Ecology Study Area
- Existing Conservation/Offset Area

**Threatened Species**

- Pink-tailed Legless Lizard
- Striped Legless Lizard
- ▲ Squirrel Glider
- ▲ Southern Myotis


**Reference:**

1. Ecotone (2000)
5. Umwelt (2007b)
6. Cumberland Ecology (2009a)
7. Cumberland Ecology (2010)
8. Cumberland Ecology (2012)
13. OEH (2019)
15. Future Ecology (2019)
16. Eco Logical Australia (2014)
18. Eco Logical Australia (2016a)
19. Eco Logical Australia (2016b)
20. Eco Logical Australia (2017)

*Note: There are no references 2 - 4, 9 - 12, 14 and 17 on this figure.*

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011

**Figure 12**



- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
- Threatened Species**
- Green and Golden Bell Frog
  - Pink-tailed Legless Lizard
  - Striped Legless Lizard
  - Swift Parrot
  - Painted Honeyeater
  - Spotted-tailed Quoll
  - Koala
  - Grey-headed Flying-fox
  - Large-eared Pied Bat
  - Corben's Long-eared Bat

**Reference:**

5. Umwelt (2007b)
6. Cumberland Ecology (2009a)
7. Cumberland Ecology (2010)
13. OEH (2019)
15. Future Ecology (2019)
16. Eco Logical Australia (2014)
18. Eco Logical Australia (2016a)
19. Eco Logical Australia (2016b)
20. Eco Logical Australia (2017)

Note: There are no references 1 - 4, 8 - 12, 14 and 17 on this figure.

Source: © NSW Department of Planning and Environment (2019); NSW Department of Finance, Services & Innovation (2019) Orthophoto Mosaic: 2018, 2016, 2011

**MALABAR COAL**

**MAXWELL PROJECT**

**Threatened Species Listed under the EPBC Act**



**Figure 13**

To further refine the potential habitat within the study area, in accordance with advice from the OEH, rocky areas were identified in PCT 1606 and 1606 DNG (where the Pink-tailed Legless Lizard was previously recorded during the 2018 surveys) (Section 2.3.3). In order to create the species polygon, a 50 m zone was applied around the rocky areas, as requested by OEH.

Figure 14 shows the species polygon (extent of habitat) for the Pink-tailed Legless Lizard in study area.



**Plate 1: Pink-tailed Legless Lizard at Site 5, November 2018 (Henry Cook).**

**Striped Legless Lizard (*Delma impar*)**

This species is listed as 'vulnerable' under the BC Act and EPBC Act. It is classified as a 'Species Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).


In NSW, the Striped Legless Lizard mostly occurs in the Southern Tablelands and the South West Slopes regions. It also occurs in the ACT, Victoria and south-eastern South Australia (OEH, 2019c).

Striped Legless Lizards are easily distinguished from other members of the genus *Delma* by the combination of two supranasal scales (rather than four), the first upper labial scale being partially fused to the nasal scale, the third upper labial scale being below the eye and the possession of a linear pattern (Smith and Robertson, 1999; Wilson and Swan, 2017; Cogger, 2018).

It is mainly found in Natural Temperate Grassland dominated by perennial, tussock-forming grasses such as Kangaroo Grass *Themeda australis*, spear-grasses *Austrostipa* spp. and poa tussocks *Poa* spp., and occasionally wallaby grasses *Austrodanthonia* spp., but has also been observed in grasslands that have a high exotic component and occasionally in open Box-Gum Woodland provided the tree cover is low (ACT Government, 2017b; OEH, 2019c). It is sometimes found in grasslands with significant amounts of surface rocks, which are used for shelter (OEH, 2019c).

This species has not been previously recorded in the study area (Table 1). 26 observations of Striped Legless Lizard as either living specimens (16 individuals) or sloughs (shed skins) (10 sloughs) were made at several locations within the study area in November and December 2018 (Figure 8; Plate 2). The majority of the observations were made under cow pats (dung) in open grassy areas with a good cover of native grasses and herbs. There was one observation within a pitfall trap at Site 3 and one slough and an individual animal observed beneath artificial shelter habitat at Site 6.






- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
  - Threatened Species**
  - Pink-tailed Legless Lizard
  - SPECIES POLYGON MAPPING**
  - 2. White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)
  - 2a. White Box - Ironbark - Red Gum Shrubby Forest - Derived Native Grassland (PCT1606)
  - Other**
  - Rocky Area
  - Rocky Area 50 m Zone

*Reference:*  
 15. Future Ecology (2019)  
*Note: There are no references 1 - 14 on this figure.*

*Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011*

  
**MAXWELL PROJECT**  
 Pink-tailed Legless Lizard  
 Species Polygon



**Figure 14**

There are only five previous records of this species from the Upper Hunter area, all from near Muswellbrook Common and dated from the year 2013. This is approximately 15 km north-east of the study area. The Upper Hunter population appears to be disjunct from other recorded populations which occur greater than 200 km to the south.

Prior to commencement of surveys Muswellbrook Common was inspected by two ecologists from Future Ecology and was found to be composed of a thick cover of largely exotic grasses and forbs together with some rocks and dumped materials. The vegetation where this species was found in the study area is mapped as the following PCTs (Hunter Eco, 2019):

- PCT 1606 (3 observations).
- PCT 1606 - DNG (18 observations).
- PCT 1655 (1 observation).
- PCT 1655 DNG (2 observations).
- PCT 1692 (1 observation).
- An unmapped location just outside of study area with adjacent mapped vegetation being PCT 1606 and PCT 1606 DNG (1 observation).

The following PCTs are published in the *Threatened Biodiversity Data Collection* (OEH, 2019a) as being associated with this species within the Sydney Basin – Hunter IBRA sub-region:

- PCT 1655.
- PCT 1604.
- PCT 1691.
- PCT 1692.
- PCT 1693.

The density of native grass and forbs across the study area would fluctuate due to rainfall and grazing pressure. It was noted that the property was de-stocked around August-September 2018. During surveys in November-December 2018 a number of previously unseen forb species were conspicuous due to flowering and grass cover seemed to be subjectively denser than previous surveys. Given that most observations were made under cow pats then cattle may form an important role in microhabitat creation for this species in the study area but conversely intense grazing pressure, pasture improvement and ploughing are known to be deleterious to Striped Legless Lizard (ACT Government, 2017b).

Given the paucity of previous records of this species in the Upper Hunter and the lack of research on locally preferred vegetation, it is possible that all of the published associated PCTs together with the non-associated PCTs in which it was observed to occur in the study area would form potential habitat for this species across the study area, namely:

- PCT 1655.
- PCT 1655 DNG.
- PCT 1604.
- PCT 1606.
- PCT 1606 DNG.
- PCT 1691.
- PCT 1691 DNG.

- PCT 1692.
- PCT 1693.
- PCT 1693 DNG.

Figure 15 shows the species polygon (extent of habitat) for the Striped Legless Lizard in study area.



**Plate 2: Striped Legless Lizard, November-December 2018 (Alex Dudley).**

**Square-tailed Kite (*Lophoictinia isura*)**

This species is listed as 'vulnerable' in NSW and it is not nationally listed. It is classified as a 'Species/Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).


This species has not been previously recorded in the study area (Table 1). A single individual of this species was observed at Site 5 (within PCT 1655) in September 2018 during vegetation surveys (Colin Driscoll Hunter Eco, pers. obs.). (Figure 9). It was not observed during any other survey period.

A raptor nest located within 200 m of this observation was checked several times during September, November and December 2018 by Future Ecology and while it appeared to be in use in September due to fresh prey remains and whitewash below the nest no raptor species was ever observed in or near the nest. During the November-December surveys the nest appeared to be unoccupied.

Several additional raptor nests were detected across the study area during current surveys and if occupied the only occupants observed were Wedge-tailed Eagles.

There are additional records of this species outside the study area (ALA, 2018) where it appears to have been observed in cleared agricultural land and disturbed native vegetation.

No evidence of breeding habitat for this species was recorded and therefore this species is regarded as an Ecosystem Credit Species within the study area. The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which occur in the study area would provide potential habitat (Appendix C).



- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
  - Threatened Species**
  - Striped Legless Lizard
  - SPECIES POLYGON MAPPING**
  - Dry Sclerophyll Forests (Shrub/grass sub-formation)
  - 2. White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)
  - 2a. White Box - Ironbark - Red Gum Shrubby Forest - Derived Native Grassland (PCT1606)
  - Dry Sclerophyll Forests (Shrubby sub-formation)
  - 3. Slaty Box Shrubby Woodland (PCT1655)
  - 3a. Slaty Box Shrubby Woodland - Derived Native Grassland (PCT1655)
  - Grassy Woodlands
  - 6. Bull Oak Grassy Woodland (PCT1692)
  - 7. Yellow Box - Apple Grassy Woodland (PCT1693)
  - 7a. Yellow Box - Apple Grassy Woodland - Derived Native Grassland (PCT1693)
  - 9. Ironbark - Grey Box Grassy Woodland (PCT1691)
  - 9a. Ironbark - Grey Box Grassy Woodland - Derived Native Grassland (PCT1691)
  - 11. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)

*Reference:*  
 15. Future Ecology (2019)  
*Note:* There are no references 1 - 14 on this figure.

*Source:* © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011

**Figure 15**

**White-bellied Sea-Eagle (*Haliaeetus leucogaster*)**

This species is listed as 'vulnerable' in NSW and is nationally listed as a Marine species. It is classified as a 'Species/Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

This species has not been previously recorded in the study area (Table 1). Two individuals were observed gliding across Site 15 (over PCT 1604) during the August 2018 surveys (Figure 9). They were not observed to land or to perch.

There are additional records of this species outside the study area (Birdlife Australia, 2018; ALA, 2018; OEH, 2019b) where it appears to have been observed over water (Lake Liddell), cleared agricultural land and native vegetation as well as near the Mt Arthur Mine.

A large raptor nest was detected at Site 15 during the August 2018 survey but it was observed to be occupied by a Wedge-tailed Eagle. Several additional raptor nests were detected across the study area during current surveys and if occupied the only occupants observed were Wedge-tailed Eagles.

No evidence of breeding habitat for this species was recorded and therefore this species is regarded as an Ecosystem Credit Species within the study area. The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which occur in the study area would provide potential habitat (Appendix C) together with the non-associated PCT 1604.

**Spotted Harrier (*Circus assimilis*)**

This species is listed as 'vulnerable' in NSW but it is not nationally listed. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

This species was observed as single individuals at several locations in the study area including Sites 5 and 18 during the November-December 2018 survey period mostly foraging over open grassland (Figure 9; Plate 3). It is indeterminant whether the same or several individuals were observed over the survey period.

The mapped PCTs in which it was observed during November 2018 were PCT1606 DNG and PCT1691.

Several raptor nests were detected across the study area during current surveys and if occupied the only occupants observed were Wedge-tailed Eagles.

The PCTs in the *Threatened Biodiversity Data Collection* (OEH, 2019a) associated with this highly mobile species which have been mapped in the study area would provide potential habitat (Appendix C) (i.e. PCT 1731) together with two additional non-associated PCTs in which it was observed, namely PCT 1606 DNG and PCT 1691.



**Plate 3: Spotted Harrier, November 2018 (Alex Dudley).**

**Little Eagle (*Hieraaetus morphnoides*)**

This species is listed as 'vulnerable' in NSW and it is not nationally listed. It is classified as a 'Species/Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

This species was observed as single individuals at two locations during the November 2018 survey (Figure 9).

Two different individuals may have been observed based on morphology – one individual appeared to be a light morph and the other a dark morph.

The mapped PCT in which it was observed during November 2018 was PCT 1606 and 1606 DNG. Mapped PCTs at previous observations of this species by others within the study area include PCT 1655 DNG (Cumberland Ecology, 2012). It was also recorded by Umwelt (2007b) within the study area but it is not clear in what PCT (note this record is not shown on Figure 9 as the location was not reported).

There are additional records of this species outside the study area where it appears to have been observed in both cleared agricultural land, rural residential land and disturbed native vegetation.

Several raptor nests were detected across the study area during current surveys and if occupied the only occupants observed were Wedge-tailed Eagles.

No evidence of breeding habitat for this species was recorded and therefore this species is regarded as an Ecosystem Credit Species within the study area. The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which occur in the study area would provide potential habitat (Appendix C).

**Glossy Black Cockatoo (*Calyptorhynchus lathamii*)**

This species is listed as 'vulnerable' in NSW and it is not nationally listed. It is classified as a 'Species/Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

This species has not been previously recorded in the study area (Table 1). It was detected indirectly at Site 16 in August 2018 (Figure 9), via the observation of chewed cones of a she-oak species *Allocasuarina gymnanthera*, a known food species for Glossy Black Cockatoo (DEC, 2004b). Chewed cones were observed under several trees at two nearby locations within Site 16. The second location was about 20 m north of the actual northern boundary of the study area and Site 16. The actual birds were never observed and no nesting activity was observed despite the presence of suitable large tree hollows at several locations within the study area.

The mapped PCT in which it was observed within the study area was PCT 1604.

The vegetation at the second location just outside the northern boundary of the study area and within 20 m of Site 16 is not mapped but the adjacent vegetation includes:

- PCT 1598.
- PCT 1604.

There are no previous observations of this species by others within the study area.

No evidence of breeding habitat for this species was recorded and therefore this species is regarded as an Ecosystem Credit Species within the study area. The published PCTs for this species which occur in the study area would provide potential habitat (Appendix C).

#### **Little Lorikeet (*Glossopsitta pusilla*)**

This species is listed as 'vulnerable' in NSW and it is not nationally listed. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

This species has been previously recorded in the study area by Hanson Bailey (2007) within PCT 1604 Grey Box – Spotted Gum – Narrow-leaved Ironbark Woodland (Table 1). This species was recorded in January 2018 and June 2018 (most observations during flowering of Grey Box/White Box). It was observed at Site 1 (six individuals), 3 (two individuals), 5 (six individuals), 6 (no. of individuals not recorded), 7 (one individual) and immediately adjacent to Site 11 (four individuals) but about 40 m outside of study area (Figure 9).

The mapped PCTs in which it was observed include:

- PCT 201.
- PCT 1655.
- PCT 1606 (just outside of study area).
- PCT 1607.
- PCT 1691.

The PCTs in the *Threatened Biodiversity Data Collection* (OEH, 2019a) associated with this highly mobile species which have been mapped in the study area would provide potential habitat (Appendix C).

#### **Brown Treecreeper (eastern subspecies) (*Climacteris picumnus victoriae*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was detected at Sites 5 and 11 during the 2018 survey periods (Figure 9):

- Site 5 (one to six individuals) in January, June, September and November 2018.
- Site 11 (one to two individuals) in June 2018.

The mapped PCTs in which it was observed include:

- PCT 1655.
- PCT 1606 and PCT 1606 DNG.
- PCT 1691.

Mapped PCTs at previous observations of this species by others within the study area (Figure 9) include:

- PCT 1606 DNG (OEH, 2019b).
- PCT 1691.
- PCT 1655 (Cumberland Ecology, 2012).

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C).

#### **Speckled Warbler (*Chthonicola sagittata*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was detected at several locations within the study area during the 2018 survey period as follows (Figure 9):

- Site 1 (one to three individuals) in January, June, September and November 2018.
- Site 2 (one to four individuals) in January, June and November 2018.
- Site 3 (two to three individuals) in January, June and November 2018.
- Site 5 (one to four individuals) in January, June and November 2018.
- Site 6 (one to five individuals) in January, June and November 2018.
- Site 7 (one to five individuals) in June and November 2018.
- Site 11 (two individuals) in June and November 2018.
- Site 13 (one individual) in June 2018.
- Site 15 (two to three individuals) in August 2018.

The mapped PCTs in which it was observed include:

- PCT 201 and 201 DNG;
- PCT 1655 DNG;
- PCT 1604;
- PCT 1606 and 1606 DNG;
- PCT 1607 and 1607 DNG;
- PCT 1691; and
- PCT 1692.



It has been previously observed within the study area by others including Cumberland Ecology (2012) with records from what are now Sites 2, 6, 10 and about 20 m from Site 16 just outside the study area (Eco Logical Australia, 2015) (Figure 9).

Mapped PCTs at previous observations of this species by others within the study area include:

- PCT 1606 (Cumberland Ecology, 2012);
- PCT 1691 (Cumberland Ecology, 2012; OEH, 2019b); and
- PCT 1598 (Eco Logical Australia, 2015) – just outside of Site 16 and study area.

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C).

#### **Black-chinned Honeyeater (eastern subspecies) (*Melithreptus gularis gularis*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

Two individuals of this species were observed mating at Site 6 in June 2018, within PCT 1691 (Figure 9).

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C).

#### **Painted Honeyeater (*Grantiella picta*)**

This species is listed as 'vulnerable' in NSW and is nationally listed as vulnerable. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

This species has not been previously recorded in the study area (Table 1). A single individual was detected at Site 1a in January 2018, within PCT 1607 (Figure 9).

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C) (Figure 16).

#### **Flame Robin (*Petroica phoenicea*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).


This species has not been previously recorded in the study area (Table 1). It was detected during the 2018 survey period as follows (Figure 9):

- Site 1 (one individual) in June 2018; and
- Site 15 (one individual on two dates) in August 2018.

The mapped PCTs in which it was observed were:

- PCT 1604; and
- PCT 1607.


The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C).



- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
  - Threatened Species**
  - Painted Honeyeater
  - POTENTIAL HABITAT MAPPING**
  - Dry Sclerophyll Forests (Shrub/grass sub-formation)**
  - 1. Red Gum - Ironbark - Apple Shrubby Woodland (PCT1607)
  - 2. White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)
  - Dry Sclerophyll Forests (Shrubby sub-formation)**
  - 3. Slaty Box Shrubby Woodland (PCT1655)
  - Grassy Woodlands**
  - 6. Bull Oak Grassy Woodland (PCT1692)
  - 8. Fuzzy Box Woodland (PCT201)
  - 9. Ironbark - Grey Box Grassy Woodland (PCT1691)
  - 10. Weeping Myall Woodland (PCT116)
  - 11. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)

**Reference:**  
 15. Future Ecology (2019)  
**Note:** There are no references 1 - 14 on this figure.

**Source:** © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011



**Figure 16**

**Scarlet Robin (*Petroica boodang*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was detected during the 2018 survey period as follows (Figure 9):

- Site 6 (one individual) in June 2018;
- Site 7 (one individual on two separate days) in June 2018; and
- Site 11 (one individual) in June 2018.

The mapped PCTs in which it was observed were:

- PCT 1606;
- PCT 1655 DNG; and
- PCT 1691.

It has been previously observed within the study area by others including Cumberland Ecology (2012) with records from what are now Sites 2, 6 and 10 (Figure 9). Mapped PCTs at previous observations of this species by others within the study area include:

- PCT 1606 (Cumberland Ecology, 2012); and
- PCT 1691 (Cumberland Ecology, 2012; OEH, 2019b).

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C).

**Grey-crowned Babbler (eastern subspecies) (*Pomatostomus temporalis temporalis*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was detected at several locations within the study area during the 2018 survey period as follows (Figure 9):

- Site 3 (3 to 13 individuals) in January and December 2018.
- Site 4 (3 to 9 individuals) in January, November and December 2018.
- Site 5 (3 to 7 individuals) in January and November 2018.
- Site 6 (2 to 5 individuals) in January, June, September and November 2018.
- Site 7 (2 to 9 individuals) in January, June, September, November and December 2018.
- Site 15 (3 to 6 individuals) in August 2018.
- Site 18 (five individuals) in November 2018.
- Incidental (5 to 8 individuals) in January 2018.

The mapped PCTs in which it was observed include:

- PCT 201;
- PCT 1655 and PCT 1655 DNG;
- PCT 1604;

- PCT 1606 and PCT 1606 DNG;
- PCT 1691;
- PCT 1692;
- PCT 1693; and
- PCT 1731.

It has been previously observed within the study area by others with records from what are now Site 4, just south of Site 5, near Site 9, just north of what is now Site 18 (OEH, 2019b; Cumberland Ecology, 2012:2015) (Figure 9).

Mapped PCTs at previous observations of this species by others within the study area include:

- No PCT – Planted Trees (Cumberland Ecology, 2012).
- PCT 1691 (Cumberland Ecology, 2012).
- PCT 1693 and PCT 1731 (Cumberland Ecology, 2012) – just outside of Site 18 and study area.

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C) together with additional non-associated PCTs in which it was also observed to occur within the study area, namely: PCT 1693 and PCT 1731.

#### **Varied Sittella (*Daphoenositta chrysoptera*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was detected at several locations within the study area during the 2018 survey period as follows (Figure 9):

- Site 2 (3 to 12 individuals) in January and November 2018.
- Site 7 (1 to 3 individuals) in June and November 2018.
- 50 m outside Site 11 and study area (five individuals) in June 2018.
- Site 15 (three individuals) in August 2018.

The mapped PCTs in which it was observed include:

- PCT 1604;
- PCT 1606 and 1606 DNG;
- PCT 1655 and 1655 DNG;
- PCT 1691; and
- PCT 1692.

It has been previously observed within the study area by others with a record from around 1 km north of Site 1 within the Maxwell Infrastructure area (Eco Logical Australia, 2015) (Figure 9).

Mapped PCTs at previous observations of this species by others within the study area include PCT 1598 (Eco Logical Australia, 2015).

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C) together with additional non-associated PCTs in which it was also observed to occur within the study area, namely: PCT 1598 and PCT 1692.

#### **Dusky Woodswallow (*Artamus cyanopterus cyanopterus*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

This species has not been previously recorded in the study area (Table 1). It was detected at several locations within the study area during the 2018 survey period as follows (Figure 9):

- Site 5 (no. of individuals not recorded) in January 2018.
- Site 6 (no. of individuals not recorded) in January 2018.
- Site 15 (13 individuals) in August 2018.
- Site 17 (1 to 3 individuals) in November and December 2018.

The mapped PCTs in which it was observed include:

- PCT 201;
- PCT 1606 DNG;
- PCT 1655; and
- PCT 1604.

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C) together with additional non-associated PCTs in which it was also observed to occur within the study area, namely: PCT 1604 and PCT 1606 DNG.

#### **Squirrel Glider (*Petaurus norfolcensis*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Species Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was detected at several locations within the study area during the 2018 survey period as follows (Figure 10):

- Site 11 (two individuals recorded on same occasion via spotlight) in January 2018.
- Site 5 (one individual recorded via spotlight on two occasions) in November 2018.
- Site 6 (two individuals recorded on same occasion via spotlight) in November 2018.

The mapped PCTs in which it was observed include:

- PCT 1655; and
- PCT 1606.

It has been previously observed within the study area by others with records from just north of what is now Site 17, at Site 17, and around 750 m north-west of what is now Site 1 all within the Maxwell Infrastructure area (Cumberland Ecology, 2009a; Eco Logical Australia, 2015, 2016a, 2016b; OEH, 2019b) (Figure 10).

It is not clear as to how many individual Squirrel Gliders have been previously recorded by others within the study area but Cumberland Ecology (2009a) states that five Squirrel Gliders were recorded via trapping in February 2007 in what is now Site 17. Eco Logical Australia (2016b) do not state how many individual Squirrel Gliders were recorded only the areas in which they were recorded.

There are additional records of this species outside the study area (Umwelt, 2006b, 2007b; Cumberland Ecology, 2009a, 2010; OEH, 2019b) where it appears to have been recorded in cleared agricultural land, native vegetation, disturbed native vegetation, and the edge of disturbed mining lands (note that some of these records are not shown on Figure 10 as the locations were not reported). Some of the records are from within 250 m of the north-west corner of the study area between the Maxwell Infrastructure and the Mt Arthur Mine and around 750 m north-west of what is now Site 1 (Cumberland Ecology, 2009a). Adjacent mapped vegetation in the study area is PCT 1598 and PCT 1606 DNG and woodland rehabilitation. There is an additional record from about 15 m east of the study area and what is now Site 16 (OEH, 2019b). The adjacent mapped vegetation is PCT 1598 and PCT 1604.

This species is likely to occur in low numbers throughout the study area in associated PCTs (201, 1655, 1606). It was also recorded in the study area in the non-associated PCTs 1598 and 1604. It does not require large vegetation remnants to occur as it has a relatively small mean home range of 3-9 ha in coastal habitats and 3-4 ha in productive inland habitat fragments (NSW Scientific Committee, 2008). It is more likely to occur in vegetation patches with suitable microhabitat components (Smith, 2002; Smith and Murray, 2003; NSW Scientific Committee, 2008; Beyer et al., 2008; Crane et al., 2013). including:

- large healthy eucalypt trees close to drainage lines with a preference for Yellow Box (*E. melliodora*) when eucalypts are not flowering and large healthy eucalypts on ridges and upper slopes when eucalypts are in flower;
- abundant tree hollows with small openings (<=5 cm) for denning in living or dead trees; and
- a variety of food resources including nectar, pollen, sap and invertebrates and most importantly a reliable supply of winter nectar and pollen in either the canopy or understorey plant species.

Figure 17 shows the species polygon (extent of habitat) for the Squirrel Glider in the study area.

#### **Grey-headed Flying-fox (*Pteropus poliocephalus*)**

This species is listed as 'vulnerable' under the BC Act and EPBC Act. It is classified as a 'Species/Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).


This species has been previously recorded in the study area by Hansen Bailey (2007) (Table 1). It was detected foraging in flowering White Box (*Eucalyptus albens*) trees within the study area during the 2018 survey period as follows (Figure 11):

- Site 5 (two individuals recorded) in June 2018.
- Site 11 (one individual recorded) in June 2018.

The mapped PCTs in which it was observed were PCT 1606 and PCT 1606 DNG.

It has not been previously observed within the study area by others except for a single individual observed flying over the woodland adjacent to the Maxwell Infrastructure office building (Hansen Bailey, 2007). The vegetation here has been mapped as PCT 1604.

There are some additional records of this species outside the study area including records from the adjacent Mt Arthur Mine (Niche, 2012; OEH, 2019b) where vegetation was previously listed as or currently appears to be rehabilitation grassland, rehabilitation woodland, remnant woodland, and disturbed native vegetation.



- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
  - Threatened Species**
  - ▲ Squirrel Glider
  - SPECIES POLYGON MAPPING**
  - 2. White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)
  - Dry Sclerophyll Forests (Shrubby sub-formation)
  - 3. Slaty Box Shrubby Woodland (PCT1655)
  - Forested Wetlands
  - 5. Hunter Lowland Red Gum Forest (PCT1598)
  - Grassy Woodlands
  - 8. Fuzzy Box Woodland (PCT201)
  - 11. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)
  - Other
  - Paddock Trees that may provide Potential Habitat

**Reference:**

- 5. Umwelt (2007b)
- 6. Cumberland Ecology (2009a)
- 7. Cumberland Ecology (2010)
- 13. OEH (2019)
- 15. Future Ecology (2019)
- 16. Eco Logical Australia (2014)
- 18. Eco Logical Australia (2016a)
- 19. Eco Logical Australia (2016b)

*Note: There are no references 1 - 4, 8 - 12, 14 and 17 on this figure.*

**Source:** © NSW Department of Planning and Environment (2019); NSW Department of Finance, Services & Innovation (2019) Orthophoto Mosaic: 2018, 2016, 2011

**Figure 17**

No camps of this species were recorded within the study area during current surveys. Hansen Bailey (2007) state that no camps were observed in the study area during their field surveys over 10 years ago.

The closest mapped camp of this species to the study area are from Muswellbrook township (DEE, 2019), approximately 8 km north of the study area.

Given that no camps of this species occur within the study area it will be considered as an Ecosystem Credit Species under the BAM. The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C) (Figure 18).

#### **Yellow-bellied Sheathtail-bat (*Saccolaimus flaviventris*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was detected at one location within the study area during the 2018 survey period with a definite confidence level, namely Site 2 (via acoustic recording) in January 2018. (Figure 11).

The mapped PCT in which it was recorded was PCT1606. It has been previously observed within the study area by others (Cumberland Ecology, 2011; OEH, 2018), with records from what is now Site 5, 7 as well as within old mine workings within the Maxwell Infrastructure area. Currently mapped vegetation at these observations include PCT1176 and PCT1606 (DNG).

There are a few additional records of this species outside the study area from the adjacent Mt Arthur Mine area (Cumberland Ecology, 2009a, 2012; OEH, 2019b) where it appears to have been recorded in disturbed native vegetation and what is now disturbed mining lands.

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C).


#### **Eastern Freetail-bat (*Mormopterus norfolkensis*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was detected at one location within the study area during the 2018 survey period with a definite confidence level at a small rocky escarpment near the main entrance off Golden Highway/Edderton Road (via acoustic recording) in December 2018 (Figure 11). The mapped PCT in which it was recorded was PCT 1606.

It has been previously observed within the study area by others with records from the south-west corner of the Maxwell Infrastructure (Eco Logical Australia, 2017; OEH, 2019b), within 20m of the north-east corner of the study area adjacent to what is now Site 16, just south of what is now Site 5 (OEH, 2019b), southern and central western parts of the Maxwell Infrastructure area (Eco Logical Australia, 2017), Site 6 (Ecotone, 2000), Sites 6, 7 and 12 (Cumberland Ecology, 2012) (Figure 11).





- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
  - Threatened Species**
  - ▽ Grey-headed Flying-fox
  - POTENTIAL HABITAT MAPPING**
  - Dry Sclerophyll Forests (Shrub/grass sub-formation)
  - 2. White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)
  - Dry Sclerophyll Forests (Shrubby sub-formation)
  - 3. Slaty Box Shrubby Woodland (PCT1655)
  - Grassy Woodlands
  - 6. Bull Oak Grassy Woodland (PCT1692)
  - 9. Ironbark - Grey Box Grassy Woodland (PCT1691)
  - 11. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)

**Reference:**  
 11. Niche (2012)  
 13. OEH (2019)  
 15. Future Ecology (2019)  
*Note: There are no references 1 - 10, 12 and 14 on this figure.*

**Source:** © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011

**Figure 18**

Mapped PCTs at previous observations of this species by others within the study area include:

- PCT1598 (Eco Logical Australia, 2017);
- PCT1655 (Cumberland Ecology, 2012);
- PCT1606 (Cumberland Ecology, 2012);
- PCT1691 (Ecotone, 2000); and
- PCT1693 (Cumberland Ecology, 2012).

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C) together with additional non-associated PCTs in which it was also observed to occur within the study area, namely: PCTs 1598, 1606, 1693.

### **Little Bentwing-bat (*Miniopterus australis*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as a 'Species/Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was detected at several locations within the study area during the 2018 survey period via acoustic recording with a definite confidence level as follows (Figure 11):

- Site 2 in January 2018.
- Site 5 in January 2018.
- Site 10 in January 2018.
- Site 11 in January 2018.

The mapped PCT in which it was recorded was PCT 1606 and 1606 DNG.

It has been previously observed within the study area by others with records from the south-west corner of the Maxwell Infrastructure area (Eco Logical Australia, 2017; OEH, 2019b), within 20 m of the north-east corner of the study area adjacent to what is now Site 16, and central western part of the Maxwell Infrastructure area (Eco Logical Australia, 2017) (Figure 11).

Mapped PCTs at previous observations of this species by others within the study area include PCT 1598 (Eco Logical Australia, 2017; OEH, 2019b).

Only five nursery sites /maternity colonies are known in Australia, most typically limestone caves and in NSW they share maternity roosts with the Eastern Bentwing-bat (Churchill, 2009; OEH, 2019c).

The study area provides little in the way of potential maternity roost habitat defined as caves, underground mines or tunnels by the 'Species Credit' *Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018). In addition, there no records of this species within or adjacent to the study area indicating that records were obtained within caves, roosts or where observed numbers exceeded 500 individuals.

The study area has a few minor sandstone overhangs and crevices at Site 1 and a small rocky escarpment near the entrance to the Plashett property at the corner of Golden Highway and Edderton Road. There were also some crevices associated with the old volcanic rock quarry at Site 1. There was no sign of any maternity roosts at any of these sites (including actual bats entering/exiting overhangs and crevices, guano, staining, meal remains, capture of lactating females, high numbers of calls recorded on acoustic devices) despite roost searches, harp-trapping and acoustic monitoring taking place during the summer breeding period for this species.

Due to the absence of breeding habitat, this species is considered an ecosystem credit species in the study area. The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C) together with additional non-associated PCTs in which it was also observed to occur within the study area, namely: PCTs 1598 and 1606.

### **Eastern Bentwing-bat (*Miniopterus schreibersii oceanensis*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as a 'Species/Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was detected at several locations within the study area during the 2018 survey period via acoustic recording with a definite confidence level as follows (Figure 11):

- Site 1 in January 2018.
- Site 2 in January 2018.
- Site 4 in January 2018.
- Site 5 in January 2018.
- Site 10 in January 2018.
- Site 11 in January 2018.

The mapped PCT in which it was recorded were:

- PCT 1606 and 1606 DNG; and
- PCT 1607 DNG.

It has been previously observed within the study area by others with records from just south of Site 5 (OEH, 2019b), at Sites 6, 7 and 12 (Cumberland Ecology, 2012), the north-east sections of the Maxwell Infrastructure area (OEH, 2019b), within 20m of the north-east corner of the study area adjacent to what is now Site 16 (Eco Logical Australia, 2015-2017; OEH, 2019b), in rail loop area north of Site 17 (Umwelt, 2006b), and south-western and south central parts of the Maxwell Infrastructure area (Eco Logical Australia, 2014-2017; OEH, 2019b) (Figure 11).

Mapped PCTs at previous observations of this species by others within the study area include:

- PCT1598 (Eco Logical, 2014-2017);
- PCT1655 (Cumberland Ecology, 2012);
- PCT1604 (Umwelt, 2006b);
- PCT1606 (Cumberland Ecology, 2012); and
- PCT1693 (Cumberland Ecology, 2012).

There are additional records of this species outside the study area (Umwelt, 2006b; Cumberland Ecology, 2009a; Niche, 2012; OEH, 2019b) where it appears to have been recorded in agricultural land, native vegetation, rehabilitation woodland, disturbed native vegetation and active mining disturbance areas with several records associated with the Mt Arthur Mine (note that some of these records are not shown on Figure 11 as the locations were not recorded).

It is known from at least three complex limestone cave (Karst) systems in NSW including Abercrombie, Jenolan and Wombeyan Karst Conservation Reserves and in NSW they share maternity roosts with the Eastern Bentwing-bat (National Parks and Wildlife Service, 2019).

The study area provides little in the way of potential maternity roost habitat defined as caves, underground mines or tunnels by the ‘Species Credit’ *Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018). In addition, there no records of this species within or adjacent to the study area indicating that records were obtained within caves, roosts or where observed numbers exceeded 500 individuals.

The study area has a few minor sandstone overhangs and crevices at Site 1 and a small rocky escarpment near the entrance to the Plashett property at the corner of Golden Highway and Edderton Road. There were also some crevices associated with the old volcanic rock quarry at Site 1. There was no sign of any maternity roosts at any of these sites (including actual bats entering/exiting overhangs and crevices, guano, staining, meal remains, capture of lactating females, high numbers of calls recorded on acoustic devices) despite roost searches, harp-trapping and acoustic monitoring taking place during the summer breeding period for this species.

Due to the absence of breeding habitat, this species is considered an ecosystem credit species in the study area. The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C) together with additional non-associated PCTs in which it was also observed to occur within the study area, namely: PCTs 1598, 1607 and 1693.

#### **Large-eared Pied Bat (*Chalinobolus dwyeri*)**

This species is listed as ‘vulnerable’ under the BC Act and EPBC Act. It is classified as a ‘Species Credit Species’ in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was detected at several locations within the study area during the 2018 survey period via acoustic recording with a definite confidence level as follows (Figure 11):

- Site 1 in January 2018 and December 2018.
- Site 2 in January 2018.
- Site 3 in January 2018.
- Site 4 in January 2018.
- Site 10 in January 2018.
- Site 11 in January 2018.

The mapped PCT in which it was recorded were:

- PCT 1606 and 1606 DNG; and
- PCT 1607 and 1607 DNG.

It has been previously observed within the study area by others with records from Site 7 (Cumberland Ecology, 2012), the south-west corner of the Maxwell Infrastructure area (Eco Logical Australia, 2017) and central western part of the Maxwell Infrastructure area (Eco Logical Australia, 2017) (Figure 11).

Mapped PCTs at previous observations of this species by others within the study area include:

- PCT1655 (Cumberland Ecology, 2012); and
- PCT1598 (Eco Logical Australia, 2017).

There are additional records of this species outside the study area (Cumberland Ecology, 2012; OEH, 2019b) where it appears to have been recorded in disturbed native vegetation. There is an additional record from Eco Logical Australia (2015) although it is not stated where (note this record is not shown on Figure 11 as the location was not reported).

Potential breeding habitat for this species is defined as: “The PCTs associated with the species (as per the TBDC) within 100 m of rocky areas containing caves, or overhangs or crevices, cliffs or escarpments, or old underground mines, tunnels, culverts, derelict concrete buildings” by the ‘*Species Credit*’ *Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018).

The study area has a few minor sandstone overhangs and crevices at Site 1 and a small rocky escarpment near the entrance to the Plashett property at the corner of Golden Highway and Edderton Road. There were also some crevices and Fairy Martin (*Petrochelidon ariel*) nests (Churchill, 2009) associated with the old volcanic rock quarry at Site 1. None of the overhangs at Site 1 appeared to be deep enough to provide the ‘twilight area’, and high domed ceiling with indentations in which this species prefers to roost (Churchill, 2009; QLD Department of Environment and Resource Management, 2011).

The largest overhang observed was at the small rocky escarpment near the Golden Highway and this had a depth of around 3-4 m with a crevice running through the ceiling. There was no sign of any maternity roosts at any of these sites (including actual bats entering/exiting overhangs and crevices, guano, staining, meal remains, capture of lactating females, high numbers of calls recorded on acoustic devices) despite roost searches, harp-trapping and acoustic monitoring taking place during the summer breeding period for this species (November to end of January). No bats of this species were captured in harp traps during the survey including those placed at the Site 1 Quarry (November 2018) and the Site 1 Powerline Easement (December 2018) just below a rocky escarpment. There was calls recorded from this species at the Site 1 Powerline Easement site but none at the Site 1 Quarry.

In addition, there are no records of this species within or adjacent to the study area indicating that records were obtained within caves, roosts etc.

Therefore, in accordance with ‘*Species Credit*’ *Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018) breeding habitat is not considered present on the subject land because no breeding individuals of the target species were observed. In addition, the proposed impact is not a potential ‘serious and irreversible impact’ (SAII) (OEH, 2018).

### **Southern Myotis (*Myotis macropus*)**

This species is listed as ‘vulnerable’ under the BC Act but is not listed under the EPBC Act. It is classified as a ‘Species Credit Species’ in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

Two individuals were observed foraging over a dam at Site 10 within the study area in June 2018 (Figure 11). The pair were observed for nearly 30 minutes as they foraged within 20 cm of the dam surface occasionally raking its surface. An acoustic recorder was also used to supplement the visual observation. At least some of the calls had the typical characteristics for this species.

The mapped PCT around this dam at Site 10 was PCT 1606 and 1606 DNG.

It was not recorded with a definite confidence level anywhere else in the study area during the current survey period, was not caught in harp traps placed around dams at Site 1, 2, 3 and 5 and a mine dam (known as Savoy Dam) in November/December 2018 and was not observed foraging over any dam (apart from Site 10).

It has been previously observed within the study area by others with records from south of Site 5 (OEH, 2019b), Sites 3, 5 and 10 (Ecotone, 2000), Sites 6, 7 and 12 (Cumberland Ecology, 2012), the south-west corner and western central areas of the Maxwell Infrastructure area (Eco Logical Australia, 2017).

Mapped PCTs at previous observations of this species by others within the study area include:

- PCT 1655 (Ecotone, 2000; Cumberland Ecology, 2012);
- PCT 1606 (Ecotone, 2000; Cumberland Ecology 2012);
- PCT 1606 DNG (OEH, 2019b);
- PCT 1693 (Cumberland Ecology, 2012); and
- PCT 1598 (Eco Logical Australia, 2017).

There are additional records of this species outside the study area (Cumberland Ecology, 2012; Eco Logical Australia, 2017; OEH, 2019b) where it appears to have been recorded in native vegetation, disturbed native vegetation, current disturbed mine workings, a dam and cleared agricultural land (note some of these records are not shown on Figure 11 as the location was not reported).


Potential breeding habitat for this species is defined as: "The range of PCTs associated with the species (as per the TBDC) within 200 meters of any medium to large permanent creeks, rivers, lakes or other waterways (i.e. with pools/ stretches 3m or wider)" by the *'Species Credit' Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018).

The study area has many farm dams but no permanently flowing creeks. No Southern Myotis was caught via harp trapping around the pond in the Old Quarry at Site 1 and at dams at Sites 2, 3, and 5. The study area lacks old wooden bridges typically favoured by this species. Culverts at Sites 10 and 17 and another at the Railway Loop dam were checked visually for roosting bats during the day but no bats, staining or guano was observed. The only potential breeding habitat would be the numerous old hollow-bearing trees which occur near some dams. Little Forest Bat (*Vespadelus vulturnus*) was observed to be possibly roosting in a hollow-bearing tree adjacent to a dam at Site 5 but Southern Myotis was not detected.

In addition, there no records of this species within or adjacent to the study area indicating that records were obtained within roosts etc.

Therefore, in accordance *'Species Credit' Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018) habitat is considered present in the study area in dams and ponds which occur in associated PCTs mapped for the study area, namely 1691, 1604, 1692.

Figure 19 shows the species polygon (extent of habitat) for the Southern Myotis in study area.



- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
  - Threatened Species**
  - ▼ Southern Myotis
  - SPECIES POLYGON MAPPING**
  - 6. Bull Oak Grassy Woodland (PCT1692)
  - 9. Ironbark - Grey Box Grassy Woodland (PCT1691)
  - 11. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)
  - Other
  - Waterbody/Dam
  - Waterbody/Dam 200 m Buffer


**Reference:**

1. Ecotone (2000)
6. Cumberland Ecology (2009a)
8. Cumberland Ecology (2012)
13. OEH (2019)
15. Future Ecology (2019)
20. Eco Logical Australia (2017)

*Note: There are no references 2 - 5, 7, 9 - 12, 14 and 16 - 19 on this figure.*

**Source:** © NSW Department of Planning and Environment (2019); NSW Department of Finance, Services & Innovation (2019) Orthophoto Mosaic: 2018, 2016, 2011

  
**MAXWELL PROJECT**  
 Southern Myotis  
 Species Polygon



**Figure 19**

### 3.3.2 Other Threatened Fauna Species Previously Recorded within the Study Area

Some additional threatened species which were not detected by Future Ecology during current surveys have been previously detected within the study area by others (Ecotone, 2000; Cumberland Ecology, 2009a:2012; Eco Logical Australia, 2017; OEH, 2018) and are listed in Table 18.

Figures 8 to 11 show the locations of threatened fauna species records (based on the surveys detailed in this report, previous surveys and database records) within the study area and surrounds. Unconfirmed records (those which are possible or probable) are not shown on the figures or the table below.

**Table 18: Threatened Species Recorded by Others in the Study Area but not Future Ecology**

Common Name	Scientific Name	Conservation Status		Credit Class <sup>3</sup>	Previous Studies <sup>8</sup>
		BC Act <sup>1</sup>	EPBC Act <sup>2</sup>		
<b>Birds</b>					
Swift Parrot	<i>Lathamus discolor</i>	E	CE	E*	A
Barking Owl	<i>Ninox connivens</i>	V	-	E <sup>&gt;</sup>	B
Hooded Robin (south-eastern form)	<i>Melanodryas cucullata cucullata</i>	V	-	E	B
Diamond Firetail	<i>Stagonopleura guttata</i>	V	-	E	A, B, G
<b>Mammals</b>					
Spotted-tailed Quoll	<i>Dasyurus maculatus maculatus</i> (south-eastern mainland population)	V	E	E	C, D
Northern Freetail-bat	<i>Mormopterus lumsdenae</i>	V	-	E	F
Corben's Long-eared Bat	<i>Nyctophilus corbeni</i>	V	V	E	B
Eastern False Pipistrelle	<i>Falsistrellus tasmaniensis</i>	V	-	E	D, E
Greater Broad-nosed Bat	<i>Scoteanax rueppellii</i>	V	-	E	B, C, D, G
Eastern Cave Bat	<i>Vespadelus trougtoni</i>	V	-	S <sup>^</sup>	A, F, G

<sup>1</sup> Conservation status under the BC Act (current as at March 2019). V = Vulnerable, E = Endangered.

<sup>2</sup> Conservation status under the EPBC Act (current as at March 2019). V = Vulnerable, E = Endangered, CE = Critically Endangered.

<sup>3</sup> Biodiversity credit class under the Threatened Biodiversity Data Collection (OEH, 2019a) (current as at March 2019). E = Ecosystem, S = Species.

\* This species is classed an ecosystem credit species in the study area based on no important habitat mapping within the study area by OEH.

> This species is a dual credit species, however, no breeding habitat was recorded and therefore it is considered an Ecosystem credit species in the study area.

<sup>^</sup> This species is a species credit species, however, no breeding habitat was recorded.

<sup>8</sup> Study area previous survey references:

A – Cumberland Ecology (2009a) and/or Cumberland Ecology (2012).

B – Ecotone (2000).

C – Eco Logical Australia (2016a).

D – Eco Logical Australia (2016b).

E – Eco Logical Australia (2015).

F – Eco Logical Australia (2017).

G – Hansen Bailey (2007).

#### **Swift Parrot (*Lathamus discolor*)**

This species is listed as 'endangered' in NSW and 'critically endangered' nationally. It is classified as a 'Species/Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a). Swift Parrot in the study area are classed as ecosystem credit species based on no important habitat mapping within the study area by OEH.

It was not recorded by Future Ecology during surveys in 2018 including surveys in June 2018 during some flowering of White Box/Grey Box (*Eucalyptus albens/moluccana*) in the study area but has been previously recorded in 2011 on what is now Site 5 (Cumberland Ecology, 2012) (Figure 9). Two individuals were detected; one observed foraging on mistletoe and Grey Box and the second individual was heard calling (Cumberland Ecology, 2012). The sightings were in PCT 1691.



There were few if any records of this species from the Upper Hunter during the winter months of 2018 but approximately 200 Swift Parrots were recorded in the Lower Hunter in May 2018 (Mick Roderick and Alex Berryman pers. obs. 29/5 in #234487 of Birdline NSW, 2019).

The PCTs in the *Threatened Biodiversity Data Collection* (OEH, 2019a) associated with this highly mobile species which have been mapped in the study area would provide potential habitat (Appendix C) (Figure 20).

### **Barking Owl (*Ninox connivens*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as a 'Species/Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was not detected during current 2018 surveys within the study area despite numerous nocturnal call-playback and spotlighting sessions throughout the year, but has been previously detected in 2000 (Ecotone, 2000) at what is now Site 5 (Figure 9). There is little information about this observation other than this species was 'tentatively recorded' in 2000 within the study area (Cumberland Ecology, 2012).

There are also two additional records of this species from the study area within 1 km of the Ecotone observation (Figure 9). The *Bionet Atlas* record (OEH, 2019b) is also from the year 2000 and this species was listed as observed (rather than heard). There is not much detail of the third record other than its coordinates (ALA, 2018). Given that all three records from the study area are within 1 km of each other and two are from the year 2000 they could represent the same observation/record or at least the same survey as Ecotone (2000).

Mapped PCTs at previous observations of this species by others within the study area include PCT 1606 DNG (Ecotone, 2000; ALA, 2018; OEH, 2019b).

There are no other records of this species from within the immediate vicinity of the study area.

Given that this species has not been recorded within the study area (or immediate vicinity) since 2000 and there was no indication of nesting/breeding, this species is regarded as an Ecosystem Credit Species within the study area. The published PCTs for this species which occur in the study area would provide potential habitat (Appendix C).

### **Hooded Robin (south-eastern form) (*Melanodryas cucullata cucullata*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).


It was not detected during Future Ecology surveys in 2018. It has been previously recorded by others within the study area just south of what is now Site 5 (OEH, 2019b) (Figure 9).

The *Bionet Atlas* record (OEH, 2019b) lists four individuals as detected on 4/02/2000 and appears to be the same record from Ecotone (2000) discussed in Cumberland Ecology (2012).

The mapped PCT in which it was observed was PCT1606 DNG.

There are no additional records of this species within or nearby the study area.


The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C).



- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
  - Threatened Species**
  - Swift Parrot
  - POTENTIAL HABITAT MAPPING**
  - Dry Sclerophyll Forests (Shrub/grass sub-formation)
  - 2. White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)
  - Dry Sclerophyll Forests (Shrubby sub-formation)
  - 3. Slaty Box Shrubby Woodland (PCT1655)
  - Grassy Woodlands
  - 6. Bull Oak Grassy Woodland (PCT1692)
  - 8. Fuzzy Box Woodland (PCT201)
  - 9. Ironbark - Grey Box Grassy Woodland (PCT1691)
  - 11. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)

Reference:  
 8. Cumberland Ecology (2012)  
 Note: There are no references 1 - 7 on this figure.

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011



**Figure 20**

**Diamond Firetail (*Stagonopleura guttata*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was not detected during Future Ecology surveys in 2018. It has been previously recorded by others within the study area as follows (Figure 9):

- At what is now Site 5 (Cumberland Ecology, 2012) – two individuals were detected.
- At the Rail Loop Dam, west of what is now Site 17 (Hansen Bailey, 2007) – three individuals were detected.
- South of what is now Site 5 (OEH, 2019b).
- An unknown location within the study area (Ecotone, 2000) (note this record is not shown on Figure 9 as the location was not reported).

The mapped PCTs in which it was observed was:

- PCT 1604 (Hansen Bailey, 2007).
- PCT 1655 (Cumberland Ecology, 2012).
- PCT 1606 DNG (OEH, 2019b).

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C) together with additional non-associated PCTs in which it was also observed to occur within the study area, namely: PCT 1691.

**Spotted-tail Quoll (*Dasyurus maculatus*) (south-eastern mainland population)**

This species is listed as 'vulnerable' in NSW and is listed as 'endangered' nationally. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was not detected during Future Ecology surveys in 2018.


It has been previously observed within the study area by others with a record from around 1 km north of Site 1 within the Maxwell Infrastructure area (Eco Logical Australia, 2016a:2016b; OEH 2019b) (Figure 10).

Mapped PCTs at previous observations of this species by others within the study area include PCT 1598 (Eco Logical Australia, 2016a:2016b), one individual was observed via wildlife camera.

There are additional records of this species outside the study area (ALA, 2018; OEH, 2019b) where it appears to have been recorded in cleared agricultural land, disturbed native vegetation, residential land (Jerrys Plains village), and on a road (New England Highway). The Spotted-tail Quoll was also tentatively recorded during the first half of 2016 by a HVEC staff member on the main access road to the Mt Arthur Mine offices (Hunter Eco, 2013).

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C).

Figure 21 shows the potential habitat for the Spotted-tail Quoll in the study area.



- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
- Threatened Species**
- ▲ Spotted-tailed Quoll
- POTENTIAL HABITAT MAPPING**
- Dry Sclerophyll Forests (Shrub/grass sub-formation)**
- 1. Red Gum - Ironbark - Apple Shrubby Woodland (PCT1607)
  - 1a. Red Gum - Ironbark - Apple Shrubby Woodland - Derived Native Grassland (PCT1607)
  - 2. White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)
  - 2a. White Box - Ironbark - Red Gum Shrubby Forest - Derived Native Grassland (PCT1606)
- Dry Sclerophyll Forests (Shrubby sub-formation)**
- 3. Slaty Box Shrubby Woodland (PCT1655)
  - 3a. Slaty Box Shrubby Woodland - Derived Native Grassland (PCT1655)
- Forested Wetlands**
- 4. Swamp Oak Forest (PCT1731)
  - 5. Hunter Lowland Red Gum Forest (PCT1598)
- Grassy Woodlands**
- 6. Bull Oak Grassy Woodland (PCT1692)
  - 7. Yellow Box - Apple Grassy Woodland (PCT1693)
  - 7a. Yellow Box - Apple Grassy Woodland - Derived Native Grassland (PCT1693)
  - 8. Fuzzy Box Woodland (PCT201)
  - 8a. Fuzzy Box Woodland - Derived Native Grassland (PCT201)
  - 9. Ironbark - Grey Box Grassy Woodland (PCT1691)
  - 9a. Ironbark - Grey Box Grassy Woodland - Derived Native Grassland (PCT1691)
  - 11. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)
  - 11a. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland - Derived Native Grassland (PCT1604)

Reference:  
 13. OEH (2019)  
 18. Eco Logical Australia (2016a)  
 19. Eco Logical Australia (2016b)  
 Note: There are no references 1 - 13 and 15 - 17 on this figure.

**MALABAR COAL**

**MAXWELL PROJECT**

**Spotted-tailed Quoll  
Potential Habitat**

Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011

**Figure 21**

### **Northern Freetail-bat (*Mormopterus lumsdenae*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was not detected during the 2018 survey period by Future Ecology.

It has been previously observed within the study area by others with a single record from the south-west corner of the Maxwell Infrastructure area (Eco Logical Australia, 2017) (Figure 11). There are no details supplied as to how this species was detected but it is assumed that it was recorded on an acoustic device as part of the annual monitoring of the Maxwell Infrastructure area by Eco Logical Australia as such devices were used on the previous monitoring sessions (Eco Logical, 2014-2016).

The location where this species was recorded is mapped as PCT 1598 (Eco Logical Australia, 2017).

There are no additional records of this species outside the study area or within the Muswellbrook LGA (OEH, 2019b).

Given its current published distributional range of this species it is unlikely that this species occurs in the study area; positive identification would need to be made from a caught individual to confirm its occurrence locally.

Given that it does not occur in the region then, are no PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

### **Corben's Long-eared Bat (*Nyctophilus corbeni*)**

This species is listed as 'vulnerable' under the BC Act and EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was not detected with a definite confidence level during the 2018 survey period by Future Ecology.

It has been previously observed within the study area by others with records south-west of Site 5 (OEH, 2019b), and south of Site 5, including Sites 5 and Site 6 (Ecotone, 2000) (Figure 11). There is no information available as to if this species was identified by live trapping or by call recording. A record in the *Bionet Atlas* (OEH, 2019b) dated the year 2000 from Saddlers Creek is probably an Ecotone (2000) record and states the detection method as 'M' or miscellaneous.

Mapped PCTs at previous observations of this species by others within the study area include:

- PCT 1655 (Ecotone, 2000).
- PCT 1606 DNG (OEH, 2019b; Ecotone, 2000).
- PCT 1691 (Ecotone, 2000).

There is one additional record of this species outside the study area (where it occurs about 190m west of Site 2 (Ecotone, 2000) in disturbed native vegetation. Nearby mapped native vegetation includes PCT1606 DNG and PCT1692.

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat together with an additional non-associated PCT in which it was also observed to occur within the study area, namely: PCT 1691.

### **Eastern False Pipistrelle (*Falsistrellus tasmaniensis*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was not detected within the study area during the 2018 survey period with a definite confidence level.

It has been previously observed within the study area by others with records from the south-west corner of the Maxwell Infrastructure area and within 20m of the north-east corner of the study area adjacent to what is now Site 16 (Eco Logical Australia, 2015:2016b; OEH, 2019b) (Figure 11).

Mapped PCTs at previous observations of this species by others within the study area include PCT 1598 (Eco Logical Australia, 2015:2016b; OEH, 2019b).

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C) together with an additional non-associated PCT in which it was also observed to occur within the study area, namely: PCT 1598.

### **Greater Broad-nosed Bat (*Scoteanax rueppellii*)**

This species is listed as 'vulnerable' under the BC Act and is not listed under the EPBC Act. It is classified as an 'Ecosystem Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was not detected within the study area during the 2018 survey period with a definite confidence level.

It has been previously observed within the study area by others with records from just south-west of Site 5 (OEH, 2019b), Site 6, 10 (Ecotone, 2000), the south-west corner of the Maxwell Infrastructure area and within 20m of the north-east corner of the study area adjacent to what is now Site 16 (Eco Logical Australia, 2016a:2016b; OEH, 2019b) (Figure 11).

Mapped PCTs at previous observations of this species by others within the study area include:

- PCT 1598 (Eco Logical Australia, 2016a:2016b).
- PCT 1606 (Ecotone, 2000).
- PCT 1691 (Ecotone, 2000).

The PCTs assigned to this species in the *Threatened Biodiversity Data Collection* (OEH, 2019a) which have been mapped in the study area would provide potential habitat (Appendix C) together with an additional non-associated PCT in which it was also observed to occur within the study area, namely: PCT 1598.

### **Eastern Cave Bat (*Vespadelus troughtoni*)**

This species is listed as 'vulnerable' under the BC Act but is not listed under the EPBC Act. It is classified as a 'Species Credit Species' in the *Threatened Biodiversity Data Collection* (OEH, 2019a).

It was not detected within the study area during the 2018 survey period.

It has been previously observed within the study area by others with records from Sites 6, 7 and 12 (Cumberland Ecology, 2012), the south-west corner of the Maxwell Infrastructure area (Eco Logical Australia, 2017) (Figure 11), north-east corner of the Maxwell Infrastructure area and at what is now Site 17 (Hansen Bailey, 2007).

Mapped PCTs at previous observations of this species by others within the study area include:

- PCT 1655 (Cumberland Ecology, 2012).
- PCT 1606 (Cumberland Ecology, 2012).
- PCT 1693 (Cumberland Ecology, 2012).

- PCT 1598 (Eco Logical Australia, 2017).
- PCT 1604 (Hansen Bailey, 2007).

There are additional records of this species outside the study area (Cumberland Ecology, 2009a; Niche, 2012; OEH, 2019b) where it appears to have been recorded in disturbed native vegetation, current disturbed mine workings and cleared agricultural land.

Potential breeding habitat for this species is defined as: "The PCTs associated with the species (as per the TBDC) within 100m of rocky areas, caves, overhangs crevices, cliffs and escarpments, or old underground mines or tunnels, old buildings and sheds within the potential habitat" by the '*Species credit*' *Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018).

The study area has a few minor sandstone overhangs and crevices at Site 1 and a small rocky escarpment near the entrance to the Plashett property at the corner of Golden Highway and Edderton Road. There were also some crevices and Fairy Martin (*Petrochelidon ariel*) nests (Churchill, 2009) associated with the old volcanic rock quarry at Site 1. None of the overhangs observed had domed ceiling with indentations in which this species prefers to roost (Churchill, 2009). The largest overhang observed was at the small rocky escarpment near the Golden Highway and this had a depth of around 3-4 m with a crevice running through the ceiling. There was no sign of any maternity roosts at any of these sites (including actual bats entering/exiting overhangs and crevices, guano, staining, meal remains, capture of lactating females, high numbers of calls recorded on acoustic devices) despite roost searches, harp-trapping and acoustic monitoring taking place during the summer breeding period for this species (November to end of January). No bats of this species were captured in harp traps during the survey including those placed at the Site 1 Quarry (November 2018) and the Site 1 Powerline Easement (December 2018) just below a rocky escarpment. There were no calls of this species recorded with a definite confidence level.

In addition, there no records of this species within or adjacent to the study area indicating that records were obtained within caves, roosts etc.

Therefore, in accordance with '*Species Credit*' *Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method* (OEH, 2018) breeding habitat is not considered present on the subject land because despite there being potential breeding habitat no breeding individuals of the target species were observed. In addition, the proposed impact is not a potential 'serious and irreversible impact' (SAII) (OEH, 2018).

### 3.3.3 Other Threatened Fauna Species Not Recorded in the Study Area

A number of threatened fauna not recorded in the study area during past or present surveys are discussed in Table 19.

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 19: Other Threatened Fauna Species Not Recorded in the Study Area

Common Name	Scientific Name	Conservation Status		Credit Class <sup>3</sup>	Survey Result
		BC Act <sup>1</sup>	EPBC Act <sup>2</sup>		
<b>Amphibians</b>					
Green and Golden Bell Frog	<i>Litoria aurea</i>	E	V	S	Not recorded, despite targeted surveys.
Booroolong Frog	<i>Litoria booroolongensis</i>	E	E	S	No potential habitat as preferred habitat of permanent western flowing rocky streams (OEH, 2019a) are not present within the study area. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Green-thighed Frog	<i>Litoria brevipalmata</i>	V	-	S	No potential habitat as preferred habitat of rainforest and moist eucalypt forest (OEH, 2019a) not present in study area.
<b>Reptiles</b>					
Pale-headed Snake	<i>Hoplocephalus bitorquatus</i>	V	-	S	Not recorded, despite targeted surveys.
<b>Birds</b>					
Freckled Duck	<i>Stictonetta naevosa</i>	V	-	E	Not recorded, despite targeted surveys. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Australasian Bittern	<i>Botaurus poiciloptilus</i>	E	E	E	Some marginal habitat present (e.g. farm and mine dams) but large permanent freshwater wetlands with tall, dense vegetation, particularly bullrushes ( <i>Typha</i> spp.) and spikerushes ( <i>Eleocharis</i> spp.) (OEH, 2019a) are absent. Not recorded despite several surveys over several years since year 2000. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Black Falcon	<i>Falco subniger</i>	V	-	E	Not recorded, despite targeted surveys. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Red Goshawk	<i>Erythrotriorchis radiatus</i>	CE	V	S	Not recorded, despite targeted surveys. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Bush Stone-curlew	<i>Burhinus grallarius</i>	E	-	S	Not recorded, despite targeted surveys.
Australian Painted Snipe	<i>Rostratula australis</i>	E	E	E	Some marginal habitat present in the form of ephemeral shallow, freshwater terrestrial wetlands (Birdlife Australia, 2018) but not recorded despite several surveys over several years since year 2000. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Eastern Curlew	<i>Numenius madagascariensis</i>	-	CE	S/E	No potential habitat, as preferred estuarine intertidal mudflat habitat (OEH, 2019a) is not present in study area. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Curlew Sandpiper	<i>Calidris ferruginea</i>	E	CE	S/E	No potential habitat, as preferred estuarine intertidal mudflat habitat (OEH, 2019a) is not present in study area. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Gang-gang Cockatoo	<i>Callocephalon fimbriatum</i>	V	-	E <sup>^</sup>	Not recorded, despite targeted surveys.
Turquoise Parrot	<i>Neophema pulchella</i>	V	-	E	Not recorded, despite targeted surveys.
Eastern Grass Owl	<i>Tyto longimembris</i>	V	-	E	Not recorded, despite targeted surveys.
Masked Owl	<i>Tyto novaehollandiae</i>	V	-	E <sup>^</sup>	Not recorded, despite targeted surveys.
Powerful Owl	<i>Ninox strenua</i>	V	-	E <sup>^</sup>	Not recorded, despite targeted surveys.



MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 19 (Continued): Other Threatened Fauna Species Not Recorded in the Study Area

Common Name	Scientific Name	Conservation Status		Credit Class <sup>3</sup>	Survey Result
		BC Act <sup>1</sup>	EPBC Act <sup>2</sup>		
Regent Honeyeater	<i>Anthochaera phrygia</i>	CE	CE	E*	Not recorded, despite targeted surveys. Regent Honeyeater in the study area are classed as ecosystem credit species based on no important habitat mapping within the study area by OEH.
<b>Mammals</b>					
Brush-tailed Phascogale	<i>Phascogale tapoatafa</i>	V	-	S	Not recorded, despite targeted surveys.
Common Planigale	<i>Planigale maculata</i>	V	-	S	Not recorded, despite targeted surveys.
Koala	<i>Phascolarctos cinereus</i>	V	V	E <sup>^</sup>	Not recorded, despite targeted surveys.
Eastern Pygmy-possum	<i>Cercartetus nanus</i>	V	-	S	Not recorded, despite targeted surveys.
Yellow-bellied Glider	<i>Petaurus australis</i>	V	-	E	Not recorded, despite targeted surveys.
Greater Glider	<i>Petauroides volans</i>	-	V	S	Not recorded, despite targeted surveys. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Brush-tailed Rock-wallaby	<i>Petrogale penicillata</i>	E	V	S	No potential habitat as preferred rocky escarpment habitat with complex structures such as fissures, caves and ledges absent from study area. The two minor and relatively simple rocky escarpment habitats present in the study area were the subject of targeted surveys and this species was not recorded.
New Holland Mouse	<i>Pseudomys novaehollandiae</i>	-	V	E	Not recorded, despite targeted surveys.

<sup>1</sup> Conservation status under the BC Act (current as at March 2019). V = Vulnerable, E = Endangered, CE = Critically Endangered

<sup>2</sup> Conservation status under the EPBC Act (current as at March 2019). V = Vulnerable, E = Endangered, CE = Critically Endangered

<sup>3</sup> Biodiversity credit class under the Threatened Biodiversity Data Collection (OEH, 2019a) (current as at March 2019). E = Ecosystem, S = Species.

\* This species is classed an ecosystem credit species in the study area based on no important habitat mapping within the study area by OEH.

<sup>^</sup> This species is a dual credit species, however, no core habitat is present therefore it is considered an ecosystem credit species in the study area.

### 3.4 Potential Koala Habitat – SEPP 44

There are two relevant definitions that apply when considering Koala habitat under SEPP 44:

- 'potential koala habitat' means areas of native vegetation where the trees of the types listed in Schedule 2 constitute at least 15% of the total number of trees in the upper or lower strata of the tree component; and
- 'core koala habitat' means an area of land with a resident population of koalas, evidenced by attributes such as breeding females (that is, females with young) and recent sightings and historical records of a population.

Koala preferred feed tree species listed in SEPP 44 are:

- Grey Gum (*Eucalyptus punctata*);
- Forest Red Gum (*E. tereticornis*);
- Swamp Mahogany (*E. robusta*);
- Tallowwood (*E. microcorys*);
- Ribbon or Manna Gum (*E. viminalis*);
- River Red Gum (*E. camaldulensis*);
- Broad-leaved Scribbly Gum (*E. haemastoma*);
- Scribbly Gum (*E. signata*);
- White Box (*E. albens*); and
- Bimble Box or Poplar Box (*E. populnea*).

#### *Koala Potential Habitat*

Hunter Eco (2019) undertook a survey of potential koala food trees in the study area. Of the SEPP 44 preferred feed trees, two occur in the study area, namely Forest Red Gum, which is part of PCT 1598 mapped in only a few small locations, and White Box, which is part of PCT 1606 (Figure 22). PCT 1598 and PCT 1606 provide 'potential koala habitat' as defined by SEPP 44 because areas of native vegetation where the trees of the types listed in Schedule 2 constitute at least 15% of the total number of trees in the upper or lower strata of the tree component.

The following additional Koala food tree species (recognised by Department of Planning and Environment, 2018) were identified in the study area (Hunter Eco, 2019):

- Grey Box (*E. moluccana*) within PCT 1604;
- Yellow Box (*E. melliodora*) within PCT 1693;
- Blakely's Red Gum (*E. blakelyi*) within PCT 1607 and PCT 1606; and
- Fuzzy Box (*E. conica*) with PCT 201.

The *Threatened Biodiversity Data Collection* (OEH, 2019a) also recognises PCT 1655 could provide potential habitat. However, the occurrence of PCT1655 in the study area only contains Slaty Box (*E. dawsonii*) which is not a recognised koala food tree.


Potential koala habitat is mapped on Figure 22.

*Koala Presence*

No 'core koala habitat' occurs in the study area. The Koala was not detected during the 2018 survey period by Future Ecology and it has not been previously recorded within the study area during past studies. There are a few additional records of this species outside the study area including from:

- disturbed mining land at the Mt Arthur Mine about 3 km west of the study area (HVEC Personnel pers. Comms., 2012 in Hunter Eco, 2013);
- disturbed native vegetation about 2.2 km north-east of study area dated 2006 and with an accuracy of 10 km (OEH, 2019b); and
- disturbed native vegetation / cleared powerline easement about 1.9 km east of study area dated from 1954 (OEH, 2019b).

There are 24 records of this species within the Muswellbrook LGA (OEH, 2019b). If this species does occur in the locality it is likely to be in very low numbers and/or only occurs occasionally.



- LEGEND**
- Maxwell Project Exploration Licence Boundary
  - Maxwell Project Mining and Coal Lease Boundary
  - Indicative Extent of Underground Development
  - Biodiversity Assessment Development Footprint
  - Ecology Study Area
  - Existing Conservation/Offset Area
  - Threatened Species**
  - ▲ Koala
  - POTENTIAL HABITAT MAPPING**
  - 1. Red Gum - Ironbark - Apple Shrubby Woodland (PCT1607)
  - 2. White Box - Ironbark - Red Gum Shrubby Forest (PCT1606)
  - 3. Slaty Box Shrubby Woodland (PCT1655)
  - 5. Hunter Lowland Red Gum Forest (PCT1598)
  - Grassy Woodlands
  - 7. Yellow Box - Apple Grassy Woodland (PCT1693)
  - 8. Fuzzy Box Woodland (PCT201)
  - 11. Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland (PCT1604)

*Reference:*  
 9. HVEC (pers. Comm. (2012)  
 13. OEH (2019)  
*Note: There are no references 1 - 8 and 10 - 12 on this figure.*

*Source: © NSW Department of Planning and Environment (2019);  
 NSW Department of Finance, Services & Innovation (2019)  
 Orthophoto Mosaic: 2018, 2016, 2011*

**Figure 22**

### 3.5 Threatened Fauna Species Listed under the EPBC Act

Records of threatened fauna species listed under the EPBC Act are shown on Figure 13. Five threatened fauna species listed under the EPBC Act were recorded during the surveys, namely, the Pink-tailed Legless Lizard, Striped Legless Lizard, Painted Honeyeater, Grey-headed Flying-fox and Large-eared Pied Bat. Two additional threatened fauna species listed under the EPBC Act were previously recorded in the study area during other surveys, namely, the Swift Parrot and Spotted-tailed Quoll (south-eastern mainland population).

The Corben's Long-eared Bat may also have been recorded in the study area nearly 20 years ago but the record is uncertain as the detection method is not known. This species was not recorded with certainty during the present survey (Section 3.3.2).

The potentially relevant threatened species under the EPBC Act are listed in Table 20.

**Table 20: Threatened Fauna Species Listed under the EPBC Act**

Common Name	Scientific Name	Conservation Status		Survey Result
		BC Act <sup>1</sup>	EPBC Act <sup>2</sup>	
<b>Amphibians</b>				
Green and Golden Bell Frog	<i>Litoria aurea</i>	E	V	Not recorded, despite targeted during past and present surveys.
Booroolong Frog	<i>Litoria booroolongensis</i>	E	E	No potential habitat, as preferred habitat of permanent western flowing rocky streams (OEH, 2019a) are not present within the study area. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
<b>Reptiles</b>				
Pink-tailed Legless Lizard	<i>Aprasia parapulchella</i>	V	V	Recorded during this survey (Section 3.3.1; Figure 14).
Striped Legless Lizard	<i>Delma impar</i>	V	V	Recorded during this survey (Section 3.3.1; Figure 15).
<b>Birds</b>				
Australasian Bittern	<i>Botaurus poiciloptilus</i>	E	E	Some marginal habitat present (e.g. farm and mine dams) but large permanent freshwater wetlands with tall, dense vegetation, particularly bullrushes ( <i>Typha</i> spp.) and spikerushes ( <i>Eleocharis</i> spp.) (OEH, 2019a) are absent. Not recorded despite several surveys over several years since year 2000. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Red Goshawk	<i>Erythrotriorchis radiatus</i>	CE	V	Not recorded, despite targeted surveys. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Australian Painted Snipe	<i>Rostratula australis</i>	E	E	Some marginal habitat present in the form of ephemeral shallow, freshwater terrestrial wetlands (Birdlife Australia, 2018) but not recorded despite several surveys over several years since year 2000. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Eastern Curlew	<i>Numenius madagascariensis</i>	-	CE	No potential habitat, as preferred estuarine intertidal mudflat habitat (OEH, 2019a) is not present in study area. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Curlew Sandpiper	<i>Calidris ferruginea</i>	E	CE	No potential habitat, as preferred estuarine intertidal mudflat habitat (OEH, 2019a) is not present in study area. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Table 20 (Continued): Threatened Fauna Species Listed under the EPBC Act

Common Name	Scientific Name	Conservation Status		Survey Result
		BC Act <sup>1</sup>	EPBC Act <sup>2</sup>	
Swift Parrot	<i>Lathamus discolor</i>	E	CE	Previously recorded in 2011 on what is now Site 5 (Cumberland Ecology, 2012) (Figure 20). Two individuals were detected; one observed foraging on mistletoe and Grey Box and the second individual was heard calling (Cumberland Ecology, 2012). The sightings were in PCT 1691. Swift Parrot is classed as ecosystem credit species in the study area based on no important habitat mapping within the study area by OEH.
Regent Honeyeater	<i>Anthochaera phrygia</i>	CE	CE	Not recorded, despite targeted surveys. Regent Honeyeater are classed as ecosystem credit species in the study area based on no important habitat mapping within the study area by OEH.
Painted Honeyeater	<i>Grantiella picta</i>	V	V	Recorded during this survey (Section 3.3.1; Figure 16).
<b>Mammals</b>				
Spotted-tailed Quoll	<i>Dasyurus maculatus maculatus</i> (south-eastern mainland population)	V	E	It has been previously observed within the study area by others with a record from around 1 km north of Site 1 within the Maxwell Infrastructure area (Eco Logical Australia, 2016a:2016b; OEH 2019b) (Figure 21).
Koala	<i>Phascolarctos cinereus</i>	V	V	Not recorded, despite targeted surveys (Section 3.4; Figure 22).
Greater Glider	<i>Petauroides volans</i>	-	V	Not recorded, despite targeted surveys. The PCTs in the study area are not recognised as potential habitat for this species in the <i>Threatened Biodiversity Data Collection</i> (OEH, 2019a).
Brush-tailed Rock-wallaby	<i>Petrogale penicillata</i>	E	V	No potential habitat, as preferred rocky escarpment habitat with complex structures such as fissures, caves and ledges is absent from study area. The two minor and relatively simple rocky escarpment habitats present in the study area were the subject of targeted surveys and this species was not recorded.
Grey-headed Flying-fox	<i>Pteropus poliocephalus</i>	V	V	Recorded during this survey (Section 3.3.1; Figure 18).
Corben's Long-eared Bat	<i>Nyctophilus corbeni</i>	V	V	It has been previously observed within the study area by others with records near Sites 5 and 6 (Ecotone, 2000) (Figure 11). There is no information available as to if this species was identified by live trapping or by call recording.
Large-eared Pied Bat	<i>Chalinolobus dwyeri</i>	V	V	Recorded during this survey (Section 3.3.1). No known roosting sites.
New Holland Mouse	<i>Pseudomys novaehollandiae</i>	-	V	Not recorded, despite targeted surveys.

<sup>1</sup> Threatened species status under the BC Act (current as at March 2019). V = Vulnerable, E = Endangered, CE = Critically Endangered.

<sup>2</sup> Threatened species status under the EPBC Act (current as at March 2019). V = Vulnerable, E = Endangered, CE = Critically Endangered.

## 4 Conclusion

Future Ecology has reviewed a number of fauna surveys previously undertaken partly within and/or adjacent to the study area since the year 2000, and then undertaken additional fauna surveys in 2018.

Ten broad fauna habitat types were observed within the study area, comprising three natural habitats (Dry Sclerophyll Forest, Grassy Woodlands, Forested Wetlands) and seven secondary habitats (Derived Native Grassland, Planted Trees, Cultivation, Waterbody/Dam, Woodland Rehabilitation, Pasture Rehabilitation and Infrastructure/Cleared Land). The majority of survey sites were located within the Woodland or Open Forest broad fauna habitat types. Most woodland/forest patches showed evidence of historic and ongoing disturbance from grazing. Most woodland/forest patches were small to medium size (<150 ha), fragmented and lacked structural diversity in terms of subcanopy and understorey layers due to grazing pressure. Connectivity between remnant Woodland/Open Forest habitats was generally poor across the study area. However, some fauna habitat features such as hollow bearing trees, hollow logs, fallen timber, were present at most survey sites.

A total of 227 fauna species were recorded in the study area during the 2018 surveys including 8 amphibian, 22 reptile, 148 bird and 49 mammal species. A total of 25 threatened fauna species listed under the BC Act (all listed as vulnerable) were recorded by Future Ecology in the study area during the current surveys.

Four of the threatened fauna species recorded are considered relevant 'species credit species' in the study area, namely, Pink-tailed Legless Lizard, Striped Legless Lizard, Squirrel Glider and Southern Myotis.

Five threatened fauna species listed under the EPBC Act were recorded during the surveys, namely, the Pink-tailed Legless Lizard, Striped Legless Lizard, Painted Honeyeater, Grey-headed Flying-fox and Large-eared Pied Bat. Two additional threatened fauna species listed under the EPBC Act were previously recorded in the study area during other surveys, namely, the Swift Parrot and Spotted-tailed Quoll (south-eastern mainland population). The Corben's Long-eared Bat may also have been recorded in the study area nearly 20 years ago but the record is uncertain as the detection method is not known.

## 5 References

- Atlas of Living Australia (2018) **Atlas of Living Australia Database with Search Co-ordinates -32.3136 150.7728, -32.3136 150.9855, -32.4929 150.9855, -32.4929 150.7728**, download at <http://www.ala.org.au>. Accessed 9 July 2018.
- Australian Capital Territory Government (2017a) **Action Plan, Pink-tailed Worm Lizard *Aprasia parapulchella***, Threatened Species, ACT Government Canberra.
- Australian Capital Territory Government (2017b) **Action Plan, Striped Legless Lizard *Delma impar***, ACT Government Canberra.
- Beyer GL, Goldingay RL and Sharpe DJ (2008) **The characteristics of squirrel glider (*Petaurus norfolcensis*) den trees in subtropical Australia**. *Australian Journal of Zoology* 56(1) 13-21
- Birdlife Australia (2018) **Birdlife Australia Atlas Database with Search Co-ordinates -32.3136 150.7728, -32.3136 150.9855, -32.4929 150.9855, -32.4929 150.7728**.
- Birdline NSW (2019) **Eremaea Birdlines, Interesting and unusual bird observations**, Archive of records from 2018. Website: <http://www.ereamae.com/BirdlineRecentSightings.aspx?Birdline=2>
- Bureau of Meteorology (2018a) **Weather records from Singleton STP, NSW Automatic Weather Station No.061397**.Website: <http://www.bom.gov.au/climate/dwo/IDCJDW2122.latest.shtml>
- Bureau of Meteorology (2018b) **Drought: 15-month rainfall deficiencies**, report dated 4 July 2018. Viewed on 12/7/2018. Website: <http://www.bom.gov.au/climate/drought/>
- Bureau of Meteorology (2018c) **New South Wales in spring 2018: warm, with rain in some areas, Seasonal Climate Summary for New South Wales** - Product code IDCKGC25R0 dated 3 December 2018. Viewed on 30/1/2019. Website: <http://www.bom.gov.au/climate/current/season/nsw/summary.shtml#recordsTminAvgHigh>
- Bureau of Meteorology (2019a) **Drought report dated 8/1/2019**. Viewed on 30/1/2019. Website: <http://www.bom.gov.au/climate/drought/#tabs=Drought>
- Bureau of Meteorology (2019b) **Australia's climate in 2018 dated 10/1/2019**. Viewed on 30/1/2019. Website: <http://www.bom.gov.au/climate/current/annual/aus/#tabs=Overview>
- Christidis, L and Boles, W. (2008) **Systematics and Taxonomy of Australian Birds**, CSIRO Publishing, Clayton, VIC. ISBN: 9780643065116
- Churchill S. (2009) **Australian Bats**, 2nd Edition, Jacana Books Allen and Unwin, Crows Nest NSW.
- Clayton, M, Wombey, J.C, Mason, I.J, Chesser, R.T, and Wells, A. (2006) **CSIRO list of Australian Vertebrates**.
- Cogger, H. (2018) **Reptiles and Amphibians of Australia, updated 7th edition**, CSIRO Publishing, Clayton VIC, ISBN: 978064309696, October 2018
- Commonwealth of Australia (2014) **EPBC Act Referral Guidelines for the vulnerable koala (combined populations of Queensland, New South Wales and the Australian Capital Territory)**. Department of the Environment, Canberra, 2014.



- Crane, MJ, Lindenmayer, DB and Cunningham, RB (2013) **Use and characteristics of nocturnal habitats of the squirrel glider (*Petaurus norfolcensis*) in Australian temperate woodlands.** *Australian Journal of Zoology* 60(5) 320-329
- Cumberland Ecology (2009a) **Ecological Assessment of Section 75W Modification for Drayton Mine.** Prepared for Hansen Bailey.
- Cumberland Ecology (2009b) **Mt Arthur Coal Consolidation Project Ecological Assessment.** A report prepared by Cumberland Ecology for Hansen Bailey Pty Ltd. August 2009.
- Cumberland Ecology (2010a) **Mt Arthur Coal Flora and Fauna Monitoring Program Spring 2010 Ecological Monitoring Report.** A draft report prepared by Cumberland Ecology for Mt Arthur Coal. October 2010.
- Cumberland Ecology (2012) **Drayton South Coal Project Ecology Impact Assessment,** prepared for Hansen Bailey, Final October 2012.
- Cumberland Ecology (2015a) **Biodiversity Assessment Report for Drayton South Coal Project,** prepared for Hansen Bailey as part of the Environmental Impact Statement for this project, Final April 2015.
- Cumberland Ecology (2015b) **Biodiversity Offset Strategy for Drayton South Coal Project,** prepared for Hansen Bailey as part of the Environmental Impact Statement for this project, Final April 2015.
- Department of Environment and Conservation (2004a) **Threatened Biodiversity Survey and Assessment: Guidelines for developments and activities (working draft),** Biodiversity Conservation Unit, Department of Environment and Conservation, Sydney South NSW, November 2004.
- Department of Environment and Conservation (2004b) **North East New South Wales Glossy Black-cockatoo Feed Trees,** Natural Resource Management Advisory Series: NOTE 2, DEC Coffs Harbour, August 2004.
- Department of Environment and Climate Change (2008a) **NSW State Recovery Plan for the Koala.**
- Department of Environment and Climate Change (2008b) **Hygiene protocol for the control of disease in frogs.** Information Circular Number 6. DECC (NSW), Sydney South.
- Department of Environment and Climate Change (2009) **Threatened species survey and assessment guidelines: field survey methods for fauna Amphibians.** Sydney South, DECC 2009/213
- Department of the Environment and Energy (2018a) **Protected Matters Search within a 15km buffer of point coordinates -32.415270° and 150.858402°** being the nominal centre of the Maxwell Project study area. Data Received: July 2018.
- Department of Environment and Energy (DoEE) (2018b) **SPRAT profiles of relevant Commonwealth listed threatened and/or migratory fauna species.** Australian Government DoEE. Website: <http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl>
- Department of Environment and Energy (2019) **National Flying-fox monitoring viewer.** Australian government DoE. Viewed on 17/2/2019. Website: <http://www.environment.gov.au/webgis-framework/apps/ffc-wide/ffc-wide.jsf>

- Department of Environment, Water, Heritage and Arts (2010a) **Survey guidelines for Australia's threatened frogs.** Guidelines for detecting frogs listed as threatened under the Environment Protection and Biodiversity Conservation Act 1999, Australian Govt. Barton, ACT. Website: <https://www.environment.gov.au/resource/survey-guidelines-australias-threatened-frogs-guidelines-detecting-frogs-listed-threatened>
- Department of Environment, Water, Heritage and Arts (2010b) **Survey guidelines for Australia's threatened bats.** Guidelines for detecting bats listed as threatened under the Environment Protection and Biodiversity Conservation Act 1999, Australian Govt. Barton, ACT. Website: <http://www.environment.gov.au/resource/survey-guidelines-australias-threatened-bats-guidelines-detecting-bats-listed-threatened>
- Department of Environment, Water, Heritage and Arts (2010c) **Survey guidelines for Australia's threatened birds: Guidelines for detecting birds.** listed as threatened under the Environment Protection and Biodiversity Conservation Act 1999, Australian Govt. Barton, ACT. Website: <https://www.environment.gov.au/resource/survey-guidelines-australias-threatened-birds-guidelines-detecting-birds-listed-threatened>
- Department of Planning and Environment (2018) **Review of State Environmental Planning Policy 44 (Koala Habitat Protection).** Website: [http://planspolicies.planning.nsw.gov.au/index.pl?action=view\\_job&job\\_id=8073](http://planspolicies.planning.nsw.gov.au/index.pl?action=view_job&job_id=8073)
- Department of Sustainability, Environment, Water, Population and Communities (2011a) **Survey guidelines for Australia's threatened mammals.** Guidelines for detecting mammals listed as threatened under the Environment Protection and Biodiversity Conservation Act 1999. Australian Govt. Barton, ACT. Website: <https://www.environment.gov.au/resource/survey-guidelines-australias-threatened-mammals-guidelines-detecting-mammals-listed>
- Department of Sustainability, Environment, Water, Population and Communities (2011b) **Survey guidelines for Australia's threatened reptiles.** Guidelines for detecting reptiles listed as threatened under the Environment Protection and Biodiversity Conservation Act 1999. Australian Govt. Barton, ACT. Website: <https://www.environment.gov.au/resource/survey-guidelines-australias-threatened-reptiles-guidelines-detecting-reptiles-listed>
- Department of Sustainability, Environment, Water, Population and Communities (2011c) **Environment Protection and Biodiversity Conservation Act 1999 referral guidelines for the vulnerable striped legless lizard, *Delma impar*.**
- Department of the Environment (2014) **EPBC Act Referral Guidelines for the Vulnerable Koala,** Australian Government, DoE, Canberra.
- Eco Logical Australia (2014) **Drayton Mine 2013 Flora and Fauna Monitoring Report.**
- Eco Logical Australia (2015) **Drayton Mine 2014 Flora and Fauna Monitoring Report.**
- Eco Logical Australia (2016a) **Drayton Coal Mine 2015 Flora and Fauna Monitoring Report.**
- Eco Logical Australia (2016b) **Drayton Coal Mine 2016 Flora and Fauna Monitoring Report.**
- Eco Logical Australia (2017) **Drayton Coal 2017 Spring Biodiversity Monitoring Report.**
- Ecotone (2000) **Flora and Fauna and Threatened Species Assessment for the proposed coal mining area at Saddlers Creek.** Unpublished report by Ecotone Ecological Consultants Ltd, Waratah, NSW
- Geoscience Australia (2018a) **Compute Sunrise, Sunset & Twilight Times,** online information at Geoscience Australia, <http://www.ga.gov.au/geodesy/astro/sunrise.jsp>
-

- Geoscience Australia (2018b) **Compute Moonrise & Moonset Times**, online information at Geoscience Australia, <http://www.ga.gov.au/geodesy/astro/moonrise.jsp>
- Google Inc. (2018) **Google Earth Pro**. Website: <https://www.google.com/earth/download/gep/agree.html>
- Hansen Bailey (2007) **Drayton Mine Extension Flora and Fauna Impact Assessment**.
- Hunter Eco (2013) **Mt Arthur Coal Open Cut 2012 Modification: Flora Survey**. Unpublished Report for Mt Arthur Coal - BHP Billiton,
- Hunter Eco (2019) **Maxwell Project Baseline Flora Report**, prepared for Malabar Coal Limited.
- Lo Cascio, A. (2018) **Identification of echolocation call sequences recorded at Maxwell Project Study Area, Jerry's Plains NSW January 2018**. Unpublished report for Future Ecology, February 2018.
- Lo Cascio, A. (2019) **Identification of echolocation call sequences recorded at Maxwell Project Study Area, Jerry's Plains NSW November-December 2018**. Unpublished report for Future Ecology, February 2019.
- Malabar Coal Limited (2018) **Drayton Mine Annual Environmental Management Report 2017**.
- Museum of Applied Arts and Sciences – Sydney Observation (2018) **Observations**. Website: <https://maas.museum/observations/>
- National Parks and Wildlife Service, NSW (2019) **Eastern bentwing-bat**. Viewed on 18/2/2019. Website: <https://www.nationalparks.nsw.gov.au/plants-and-animals/eastern-bentwing-bat>
- New South Wales Spatial Services (2018) **Six Maps**. Accessed: July 2018. Website: <https://maps.six.nsw.gov.au/>
- Niche Environment and Heritage (2012) **Mt Arthur Coal Fauna Survey Report, October 2010**.
- NSW Scientific Committee (2008) **Squirrel Glider *Petaurus norfolcensis*. Review of current information in NSW**. August 2008. Unpublished report arising from the Review of the Schedules of the Threatened Species Conservation Act 1995. NSW Scientific Committee, Hurstville.
- Office of Environment and Heritage (2017) **Biodiversity Assessment Method**, NSW Government OEH, Sydney, August 2017.
- Office of Environment and Heritage (2018) **'Species Credit' Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method**, Office of Environment and Heritage, Sydney. September 2018.
- Office of Environment and Heritage (2019a) **Threatened Biodiversity Data Collection**. Viewed in March 2019. Website: [https://www.environment.nsw.gov.au/atlaspublicapp/UI\\_Modules/ATLAS\\_/AtlasSearch.arch.aspx](https://www.environment.nsw.gov.au/atlaspublicapp/UI_Modules/ATLAS_/AtlasSearch.arch.aspx)
- Office of Environment and Heritage (2019b) **BioNet Atlas. -32.3136 150.7728, -32.3136 150.9855, -32.4929 150.9855, -32.4929 150.7728**. Viewed in March 2019. Website: [https://www.environment.nsw.gov.au/atlaspublicapp/UI\\_Modules/ATLAS\\_/AtlasSearch.aspx?who=0b679421-e424-47ea-b672-f30693729a7e](https://www.environment.nsw.gov.au/atlaspublicapp/UI_Modules/ATLAS_/AtlasSearch.aspx?who=0b679421-e424-47ea-b672-f30693729a7e)
- Office of Environment and Heritage (2019c) **BioNet Threatened Species Profile Database**. Viewed in March 2019. Website: <https://www.environment.nsw.gov.au/threatenedspeciesapp/>.
- Phillips, S and Callaghan, J. (2011) **The Spot Assessment Technique: a tool for determining localised levels of habitat use by Koalas *Phascolarctos cinereus***. Australian Zoologist volume 35(3).

- QLD Department of Environment and Resource Management (2011) **National recovery plan for the large-eared pied bat *Chalinolobus dwyeri***. Report to the Department of Sustainability, Environment, Water, Population and Communities, Canberra.
- Reardon, T.B., Armstrong, K.N. and Jackson, S.M. (2015) **A current taxonomic list of Australian Chiroptera**. Australasian Bat Society. Version 2015-05-15.  
Website: <http://ausbats.org.au/taxonomic-list/4589345107>
- Smith, AP (2002) **Squirrel Glider (*Petaurus norfolcensis*) Conservation Management Plan**: Wyong Shire, unpublished report prepared for Wyong Shire Council, August 2002.
- Smith, AP and Murray, M. (2003) **Habitat requirements of the squirrel glider (*Petaurus norfolcensis*) and associated possums and gliders on the NSW Central Coast**. *Wildlife Research* 30(3)
- Smith, WJS and Robertson, P (1999) **National Recovery Plan for the Striped Legless Lizard (*Delma impar*)** 1999-2003, NSW National Parks and Wildlife Service and Wildlife Profiles Pty Ltd, June 1999
- Umwelt (2006a) **Mt Arthur Coal Mine Ecological Monitoring Report 2005**. A report prepared by Umwelt (Australia) Pty Limited for Mt Arthur Coal Pty Limited. September 2006.
- Umwelt (2006b) **Ecological Assessment Proposed South Pit Extension**.
- Umwelt (2007a) **Mt Arthur Coal Mine Ecological Monitoring Report 2006**. A report prepared by Umwelt (Australia) Pty Limited for Mt Arthur Coal Pty Limited. January 2007.
- Umwelt (2007b) **Ecological Assessment - Proposed Mt Arthur Underground Project**. A report prepared for Mt Arthur Coal. December 2007.
- United States of America Naval Observatory Astronomical Applications Department (2019) **Phases of the Moon for 2018**. Viewed on 30/1/2019. Website: [https://aa.usno.navy.mil/cgi-bin/aa\\_phases.pl?year=2018andmonth=8andday=12andnump=50andformat=p](https://aa.usno.navy.mil/cgi-bin/aa_phases.pl?year=2018andmonth=8andday=12andnump=50andformat=p)
- Van Dyck, S. and Strahan, R. (2008) **The Mammals of Australia, 3<sup>rd</sup> ed.** Edited by Steve van Dyck and Ronald Strahan. New Holland Publishers, Sydney NSW.
- Wilson, S and Swan, G (2017) **A complete guide to reptiles of Australia**, Fifth Edition, New Holland, August 2017.
- Wong DTY, Jones SR, Osborne WS, Brown GW, Robertson P, Michael DR and Kay GM (2011) **The life history and ecology of the Pink-tailed Worm-lizard *Aprasia parapulchella* Kluge – a review**. *Australian Zoologist* volume 35 (3), pp 927-940.

# Appendix A Fauna Species Detected

Sites 1 to 7

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 1	Observation Type	Confidence Level	Site 2	Observation Type	Confidence Level	Site 3	Observation Type	Confidence Level	Site 4	Observation Type	Confidence Level	Site 5	Observation Type	Confidence Level	Site 6	Observation Type	Confidence Level	Site 7	Observation Type	Confidence Level
Amphibians	Common Eastern Froglet	<i>Crinia signifera</i>							X	W																
Amphibians	Spotted Grass Frog	<i>Limnodynastes tasmaniensis</i>							X	OW								X	OW		X	W				
Amphibians	Dusky Toadlet	<i>Uperoleia fusca</i>																								
Amphibians	Smooth Toadlet	<i>Uperoleia laevigata</i>							X	OW																
Amphibians	Green Tree Frog	<i>Litoria caerulea</i>							X	O		X	O		X	O		X	O							
Amphibians	Eastern Dwarf Tree Frog	<i>Litoria fallax</i>							X	OW																
Amphibians	Broad-palmed Frog	<i>Litoria latopalmeta</i>				X	O		X	OW		X	OW					X	O		X	W				
Amphibians	Peron's Tree Frog	<i>Litoria peronii</i>				X	O,W	PR	X	OW		X	OW					X	OW		X	W		X	W	
Reptiles	Eastern Snake-necked Turtle	<i>Chelodina longicollis</i>				X	K		X	O		X	O					X	O		X	O				
Reptiles	Macquarie Turtle	<i>Emydura macquarii</i>																								
Reptiles	Eastern Stone Gecko	<i>Diplodactylus vittatus</i>				X	O		X	O		X	O		X	O		X	O		X	O,T		X	O	
Reptiles	Robust Velvet Gecko	<i>Nebulifera robusta</i>				X	O		X	O		X	O		X	O		X	O					X	O	
Reptiles	Thick-tailed Gecko	<i>Underwoodisaurus milii</i>				X	O														X	O		X	O	
<b>Reptiles</b>	<b>Pink-tailed Legless Lizard</b>	<b><i>Aprasia parapulchella</i></b>		V	V													X	O							
<b>Reptiles</b>	<b>Striped Legless Lizard</b>	<b><i>Delma impar</i></b>		V	V				X	O, H		X	T		X	H								X	H	
Reptiles	Two-clawed Worm-skink	<i>Anomalopus leuckartii</i>										X	T													
Reptiles	Southern Rainbow-skink	<i>Carlia tetradactyla</i>							X	O		X	O		X	O		X	O,T							
Reptiles	Elegant Snake-eyed Skink	<i>Cryptoblepharus pulcher</i>										X	O					X	O					X	O	
Reptiles	Robust Ctenotus	<i>Ctenotus robustus</i>				X	O		X	O		X	O,T		X	O,T		X	O,T		X	O		X	O	
Reptiles	Tree Skink	<i>Egernia striolata</i>				X	O		X	O		X	O		X	O		X	O		X	O				
Reptiles	Barred-sided Skink	<i>Concinnia tenuis</i>													X	O										
Reptiles	Eastern Ranges Rock-skink	<i>Liopholis modesta</i>				X	O		X	O					X	O								X	O	
Reptiles	South-eastern Morethia Skink	<i>Morethia boulengeri</i>																X	O							
Reptiles	Eastern Blue-tongue	<i>Tiliqua scincoides</i>																								
Reptiles	Eastern Water Dragon	<i>Intellagama lesueurii</i>																								
Reptiles	Eastern Bearded Dragon	<i>Pogona barbata</i>							X	O					X	O		X	O		X	O				
Reptiles	Sand Goanna	<i>Varanus gouldii</i>							X	FB, O																
Reptiles	Lace Monitor	<i>Varanus varius</i>				X	Q		X	Q		X	O		X	O		X	O		X	Q, O				
Reptiles	Brown-snouted Blind Snake	<i>Anilius wiedii</i>				X	O								X	O										
Reptiles	Spotted Black Snake	<i>Pseudechis guttatus</i>							X	O		X	O								X	O		X	O	
Birds	Stubble Quail	<i>Coturnix pectoralis</i>				X	O		X	O																
Birds	Brown Quail	<i>Coturnix ypsilophora</i>							X	O								X	O		X	O				
Birds	Plumed Whistling Duck	<i>Dendrocygna eytoni</i>																								
Birds	Black Swan	<i>Cygnus atratus</i>																								
Birds	Pink-eared Duck	<i>Malacorhynchus membranaceus</i>																								
Birds	Australian Wood Duck	<i>Chenonetta jubata</i>				X	O		X	O								X	O		X	O		X	O	
Birds	Pacific Black Duck	<i>Anas superciliosa</i>				X	O		X	O								X	O		X	O				
Birds	Australasian Shoveler	<i>Anas rhynchotis</i>																								
Birds	Grey Teal	<i>Anas gracilis</i>				X	O		X	O								X	O		X	O		X	O	
Birds	Chestnut Teal	<i>Anas castanea</i>							X	O		X	O													
Birds	Hardhead Duck	<i>Aythya australis</i>				X	O											X	O		X	O				

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 1	Observation Type	Confidence Level	Site 2	Observation Type	Confidence Level	Site 3	Observation Type	Confidence Level	Site 4	Observation Type	Confidence Level	Site 5	Observation Type	Confidence Level	Site 6	Observation Type	Confidence Level	Site 7	Observation Type	Confidence Level
Birds	Musk Duck	<i>Biziura lobata</i>																								
Birds	Australasian Grebe	<i>Tachybaptus novaehollandiae</i>				X	O					X	O					X	O		X	O				
Birds	Hoary-headed Grebe	<i>Poliiocephalus poliocephalus</i>																								
Birds	Straw-necked Ibis	<i>Threskiornis spinicollis</i>																								
Birds	Nankeen Night Heron	<i>Nycticorax caledonicus</i>																X	O							
<b>Birds</b>	<b>Cattle Egret</b>	<b><i>Ardea ibis</i></b>			<b>M</b>																					
Birds	White-necked Heron	<i>Ardea pacifica</i>																								
Birds	White-faced Heron	<i>Egretta novaehollandiae</i>				X	O		X	O		X	O					X	O		X	O				
Birds	Little Pied Cormorant	<i>Microcarbo melanoleucos</i>										X	O													
Birds	Pied Cormorant	<i>Phalacrocorax varius</i>																								
Birds	Australasian Darter	<i>Anhinga novaehollandiae</i>																								
Birds	Nankeen Kestrel	<i>Falco cenchroides</i>				X	OW											X	O					X	O	
Birds	Brown Falcon	<i>Falco berigora</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Birds	Black-shouldered Kite	<i>Elanus axillaris</i>																								
Birds	Black Kite	<i>Milvus migrans</i>																								
<b>Birds</b>	<b>Square-tailed Kite</b>	<b><i>Lophoictinia isura</i></b>		<b>V</b>														X	O							
<b>Birds</b>	<b>White-bellied Sea Eagle</b>	<b><i>Haliaeetus leucogaster</i></b>		<b>V</b>	<b>M</b>																					
<b>Birds</b>	<b>Spotted Harrier</b>	<b><i>Circus assimilis</i></b>		<b>V</b>														X	O							
Birds	Brown Goshawk	<i>Accipiter fasciatus</i>				X	O											X	O		X	O				
Birds	Collared Sparrowhawk	<i>Accipiter cirrocephalus</i>				X	O																			
Birds	Wedge-tailed Eagle	<i>Aquila audax</i>				X	O		X	O, E		X	O		X	O		X	O		X	O		X	O, E	
<b>Birds</b>	<b>Little Eagle</b>	<b><i>Hieraetus morphnoides</i></b>		<b>V</b>																						
Birds	Purple Swamphen	<i>Porphyrio porphyrio</i>																								
Birds	Eurasian Coot	<i>Fulica atra</i>																								
Birds	Black-winged Stilt	<i>Himantopus himantopus</i>																								
Birds	Red-necked Avocet	<i>Recurvirostra novaehollandiae</i>																								
Birds	Banded Lapwing	<i>Vanellus tricolor</i>																								
Birds	Masked Lapwing	<i>Vanellus miles</i>				X	O		X	O		X	O		X	O		X	O					X	O	
Birds	Black-fronted Dotterel	<i>Eseyornis melanops</i>																X	O							
Birds	Common Bronzewing	<i>Phaps chalcoptera</i>				X	O					X	O		X	O		X	O		X	O		X	O	
Birds	Crested Pigeon	<i>Ocyphaps lophotes</i>										X	O		X	O		X	O		X	O		X	O	
Birds	Bar-shouldered Dove	<i>Geopelia humeralis</i>							X	O, W		X	W													
<b>Birds</b>	<b>Glossy Black-Cockatoo</b>	<b><i>Calyptorhynchus lathami</i></b>		<b>V</b>																						
Birds	Yellow-tailed Black-Cockatoo	<i>Calyptorhynchus funereus</i>																								
Birds	Galah	<i>Eolophus roseicapillas</i>				X	O		X	O		X	O		X	O		X	O		X	O		X	O	
Birds	Little Corella	<i>Cacatua sanguinea</i>																X	W							
Birds	Sulphur-crested Cockatoo	<i>Cacatua galerita</i>				X	O		X	O											X	O		X	O	
Birds	Rainbow Lorikeet	<i>Trichoglossus haematodus</i>																								
Birds	Musk Lorikeet	<i>Glossopsitta concinna</i>							X	O		X	O		X	O		X	O		X	O		X	O	
<b>Birds</b>	<b>Little Lorikeet</b>	<b><i>Glossopsitta pusilla</i></b>		<b>V</b>		X	O		X	O		X	O		X	O		X	O		X	O		X	O	
Birds	Crimson Rosella	<i>Platycercus elegans</i>																								
Birds	Eastern Rosella	<i>Platycercus eximius</i>				X	O		X	O		X	O		X	O		X	O		X	O		X	O	
Birds	Australian King-Parrot	<i>Alisterus scapularis</i>				X	O		X	O								X	O							
Birds	Pallid Cuckoo	<i>Cuculus pallidus</i>													X	O		X	O							
Birds	Fan-tailed Cuckoo	<i>Cacomantis flabelliformis</i>				X	O											X	O							
Birds	Black-eared Cuckoo	<i>Chalcites osculans</i>																X	O		X	O				

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 1	Observation Type	Confidence Level	Site 2	Observation Type	Confidence Level	Site 3	Observation Type	Confidence Level	Site 4	Observation Type	Confidence Level	Site 5	Observation Type	Confidence Level	Site 6	Observation Type	Confidence Level	Site 7	Observation Type	Confidence Level
Birds	Horsfield's Bronze-Cuckoo	<i>Chalcites basalis</i>				X	O					X	O					X	O		X	O		X	O	
Birds	Shining Bronze Cuckoo	<i>Chrysococcyx lucidus</i>							X	O		X	O													
Birds	Eastern Koel	<i>Eudynamis orientalis</i>													X	O		X	OW							
Birds	Channel-billed Cuckoo	<i>Scythrops novaehollandiae</i>				X	OW		X	O		X	OW		X	O		X	O		X	O				
Birds	Eastern Barn Owl	<i>Tyto javanica</i>				X	O		X	OW					X	W					X	O				
Birds	Southern Boobook	<i>Ninox novaeseelandiae</i>																								
Birds	Tawny Frogmouth	<i>Podargus strigoides</i>				X	O		X	O		X	OW					X	O		X	O		X	O	
Birds	White-throated Nightjar	<i>Eurostopodus mystacalis</i>																								
Birds	Australian Owlet-nightjar	<i>Aegotheles cristatus</i>				X	O, W		X	O, W		X	OW		X	OW		X	OW		X	O		X	O	
Birds	Laughing Kookaburra	<i>Dacelo novaeguineae</i>				X	O		X	O		X	O		X	O		X	O		X	O		X	O	
Birds	Sacred Kingfisher	<i>Todiramphus sanctus</i>										X	O											X	W	
<b>Birds</b>	<b>Rainbow Bee-eater</b>	<b><i>Merops ornatus</i></b>			<b>M</b>													X	O							
<b>Birds</b>	<b>Brown Treecreeper (eastern subspecies)</b>	<b><i>Climacteris picumnus victoriae</i></b>		<b>V</b>														X	O							
Birds	Variigated Fairy-wren	<i>Malurus lamberti</i>							X	O																
Birds	Superb Fairy-wren	<i>Malurus cyaneus</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Birds	Spotted Pardalote	<i>Pardalotus punctatus</i>				X	O		X	W		X	O								X	O		X	O	
Birds	Striated Pardalote	<i>Pardalotus striatus</i>				X	O		X	O		X	O		X	O		X	O		X	O		X	O	
<b>Birds</b>	<b>Speckled Warbler</b>	<b><i>Chthonicola sagittata</i></b>		<b>V</b>		<b>X</b>	<b>O</b>		<b>X</b>	<b>O</b>		<b>X</b>	<b>O</b>					<b>X</b>	<b>O</b>		<b>X</b>	<b>O</b>		<b>X</b>	<b>O</b>	
Birds	Weebill	<i>Smicrornis brevirostris</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Birds	Western Gerygone	<i>Gerygone fusca</i>							X	O		X	OW		X	OW		X	O		X	O		X	O	
Birds	White-throated Gerygone	<i>Gerygone olivacea</i>										X	W					X	O							
Birds	Brown Thornbill	<i>Acanthiza pusilla</i>																								
Birds	Buff-rumped Thornbill	<i>Acanthiza reguloides</i>				X	O		X	O		X	O								X	O		X	O	
Birds	Yellow-rumped Thornbill	<i>Acanthiza chrysorrhoa</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Birds	Yellow Thornbill	<i>Acanthiza nana</i>				X	O		X	O		X	OW					X	O		X	O		X	O	
Birds	Striated Thornbill	<i>Acanthiza lineata</i>																								
Birds	Yellow-faced Honeyeater	<i>Caligavis chrysops</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Birds	Singing Honeyeater	<i>Lichenostomus virescens</i>																								
Birds	White-eared Honeyeater	<i>Nesoptilotis leucotis</i>																								
Birds	Fuscous Honeyeater	<i>Lichenostomus fuscus</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Birds	White-plumed Honeyeater	<i>Lichenostomus penicillatus</i>				X	O					X	O		X	O		X	O		X	O				
Birds	Noisy Miner	<i>Manorina melanocephala</i>							X	O		X	OW		X	O		X	O		X	O		X	O	
Birds	Blue-faced Honeyeater	<i>Entomyzon cyanotis</i>																								
<b>Birds</b>	<b>Black-chinned Honeyeater (eastern subspecies)</b>	<b><i>Melithreptus gularis gularis</i></b>		<b>V</b>																						
Birds	Brown-headed Honeyeater	<i>Melithreptus brevirostris</i>				X	O		X	O		X	O		X	O		X	O		X	O		X	O	
Birds	White-naped Honeyeater	<i>Melithreptus lunatus</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Birds	Noisy Friarbird	<i>Philemon corniculatus</i>				X	O		X	O		X	OW		X	O		X	O		X	O		X	O	
Birds	Striped Honeyeater	<i>Plectorhyncha lanceolata</i>				X	W		X	OW		X	O		X	O		X	O		X	O		X	O	
Birds	Spiny-cheeked Honeyeater	<i>Acanthagenys rufogularis</i>							X	O		X	O					X	OW		X	O				
Birds	Red Wattlebird	<i>Anthochaera carunculata</i>				X	O		X	O																
<b>Birds</b>	<b>Painted Honeyeater</b>	<b><i>Grantiella picta</i></b>		<b>V</b>	<b>V</b>	<b>X</b>	<b>W</b>																			
Birds	Eastern Spinebill	<i>Acanthorhynchus tenuirostris</i>																						X	O	
Birds	Scarlet Honeyeater	<i>Myzomela sanguinolenta</i>							X	W		X	W													
Birds	Eastern Yellow Robin	<i>Eopsaltria australis</i>				X	O		X	O														X	O	
Birds	Jacky Winter	<i>Microeca fascinans</i>																X	O		X	O		X	O	
Birds	Rose Robin	<i>Petroica rosea</i>				X	O																			

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 1	Observation Type	Confidence Level	Site 2	Observation Type	Confidence Level	Site 3	Observation Type	Confidence Level	Site 4	Observation Type	Confidence Level	Site 5	Observation Type	Confidence Level	Site 6	Observation Type	Confidence Level	Site 7	Observation Type	Confidence Level
Birds	Flame Robin	<i>Petroica phoenicea</i>		V		X	W																			
Birds	Scarlet Robin	<i>Petroica boodang</i>		V																		X	O		X	O
Birds	Red-capped Robin	<i>Petroica goodenovii</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Birds	Grey-crowned Babbler (eastern subspecies)	<i>Pomatostomus temporalis ssp temporalis</i>		V								X	OW		X	O		X	O		X	O		X	O	
Birds	Varied Sittella	<i>Daphoenositta chrysoptera</i>		V					X	O														X	O	
Birds	Golden Whistler	<i>Pachycephala pectoralis</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Birds	Rufous Whistler	<i>Pachycephala rufiventris</i>				X	O		X	OW		X	OW		X	O		X	O		X	O		X	O	
Birds	Grey Shrike-thrush	<i>Colluricincla harmonica</i>																								
Birds	Grey Fantail	<i>Rhipidura albiscapa</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Birds	Willie Wagtail	<i>Rhipidura leucophrys</i>				X	O		X	O		X	O		X	O		X	O		X	O		X	O	
Birds	Magpie-lark	<i>Grallina cyanoleuca</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Birds	Leaden Flycatcher	<i>Myiagra rubecula</i>			M	X	O											X	O							
Birds	Grey Butcherbird	<i>Cracticus torquatus</i>				X	O		X	O		X	W					X	O		X	O		X	O	
Birds	Pied Butcherbird	<i>Cracticus nigrogularis</i>				X	O		X	O		X	W		X	O		X	O		X	O		X	O	
Birds	Australian Magpie	<i>Cracticus tibicen</i>				X	O		X	O		X	O,Q		X	O		X	O		X	O		X	O	
Birds	Pied Currawong	<i>Strepera graculina</i>				X	O		X	OW		X	O		X	O		X	O		X	O		X	O	
Birds	Masked Woodswallow	<i>Artamus personatus</i>																X	O		X	O				
Birds	White-browed Woodswallow	<i>Artamus superciliosus</i>							X	O								X	O		X	O		X	O	
Birds	Dusky Woodswallow	<i>Artamus cyanopterus</i>		V														X	O		X	O				
Birds	Cicadabird	<i>Coracina tenuirostris</i>										X	O													
Birds	Black-faced Cuckoo-shrike	<i>Coracina novaehollandiae</i>				X	O		X	O		X	O		X	O		X	O		X	O		X	O	
Birds	Ground Cuckoo-shrike	<i>Coracina maxima</i>																								
Birds	White-winged Triller	<i>Lalage sueurii</i>				X	O					X	O					X	O		X	O		X	O	
Birds	Olive-backed Oriole	<i>Oriolus sagittatus</i>				X	O		X	O		X	O		X	O		X	O		X	O		X	O	
Birds	Little Raven	<i>Corvus mellori</i>																								
Birds	Australian Raven	<i>Corvus coronoides</i>				X	O		X	OW		X	O		X	O		X	O		X	O		X	O	
Birds	White-winged Chough	<i>Corcorax melanorhamphos</i>				X	O		X	O		X	Q, O		X	OW		X	O		X	O		X	O	
Birds	Common Starling	<i>Sturnus vulgaris</i>	X			X	O																			
Birds	Common Myna	<i>Sturnus tristis</i>	X																							
Birds	White-backed Swallow	<i>Cheramoeca leucosterna</i>																								
Birds	Welcome Swallow	<i>Hirundo neoxena</i>				X	O											X	O							
Birds	Fairy Martin	<i>Petrochelidon ariel</i>				X	O											X	O							
Birds	Tree Martin	<i>Petrochelidon nigricans</i>				X	O											X	O							
Birds	Silvereye	<i>Zosterops lateralis</i>				X	O																	X	O	
Birds	Australian Reed Warbler	<i>Acrocephalus australis</i>				X	OW		X	W																
Birds	Tawny Grassbird	<i>Megalurus timoriensis</i>										X	O													
Birds	Rufous Songlark	<i>Cincloramphus mathewsi</i>				X	O		X	O								X	O							
Birds	Brown Songlark	<i>Cincloramphus cruralis</i>							X	O								X	O							
Birds	Golden-headed Cisticola	<i>Cisticola exilis</i>																								
Birds	Horsfield's Bushlark	<i>Mirafra javanica</i>							X	O																
Birds	Mistletoebird	<i>Dicaeum hirundinaceum</i>				X	O		X	W		X	O		X	W		X	O		X	OW		X	O	
Birds	Richard's Pipit	<i>Anthus novaeseelandiae</i>																								
Birds	Red-browed Finch	<i>Neochmia temporalis</i>				X	O																			
Birds	Zebra Finch	<i>Taeniopygia guttata</i>																								
Birds	Double-barred Finch	<i>Taeniopygia bichenovii</i>				X	O		X	O		X	O					X	O		X	O		X	O	
Mammals	Short-beaked Echidna	<i>Tachyglossus aculeatus</i>				X	P, O		X	P, O		X	F, P					X	O		X	O				



MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 1	Observation Type	Confidence Level	Site 2	Observation Type	Confidence Level	Site 3	Observation Type	Confidence Level	Site 4	Observation Type	Confidence Level	Site 5	Observation Type	Confidence Level	Site 6	Observation Type	Confidence Level	Site 7	Observation Type	Confidence Level
Mammals	Yellow-footed Antechinus	<i>Antechinus flavipes</i>							X	O		X	T													
Mammals	Common Dunnart	<i>Sminthopsis murina</i>				X	O														X	O		X	O	
Mammals	Common Wombat	<i>Vombatus ursinus</i>																								
Mammals	Sugar Glider	<i>Petaurus breviceps</i>																X	F	PO						
<b>Mammals</b>	<b>Squirrel Glider</b>	<b><i>Petaurus norfolcensis</i></b>		V														X	O		X	O				
Mammals	Common Ringtail Possum	<i>Pseudocheirus peregrinus</i>																								
Mammals	Common Brushtail Possum	<i>Trichosurus vulpecula</i>				X	Q,O,X,H	D	X	O,Q		X	O		X	O		X	O		X	O		X	O	
Mammals	brushtail possum	<i>Trichosurus sp.</i>				X	H,P	PR																		
Mammals	Eastern Grey Kangaroo	<i>Macropus giganteus</i>				X	O,Y,P,X		X	O		X	O		X	O		X	O		X	O		X	O	
Mammals	Eastern Wallaroo	<i>Macropus robustus</i>				X	O																			
Mammals	Red-necked Wallaby	<i>Macropus rufogriseus</i>				X	O		X	Q,O		X	O		X	O		X	Q		X	O		X	O	
Mammals	Swamp Wallaby	<i>Wallabia bicolor</i>				X	X,Y																			
<b>Mammals</b>	<b>Grey-headed Flying-fox</b>	<b><i>Pteropus poliocephalus</i></b>		V	V													X	O							
<b>Mammals</b>	<b>Yellow-bellied Sheathtail-bat</b>	<b><i>Saccolaimus flaviventris</i></b>		V					X	U	D															
<b>Mammals</b>	<b>Eastern Freetail-bat</b>	<b><i>Mormopterus norfolkensis</i></b>		V																				X	U	PO
Mammals	Little Mastiff-bat	<i>Mormopterus planiceps</i>				X	U	D	X	U	D	X	U	D	X	U	D	X	U	D	X	U	D	X	U	D
Mammals	Eastern Free-tailed Bat	<i>Mormopterus ridei</i>				X	U	D	X	U	D				X	U	PO									
Mammals	White-striped Freetail-bat	<i>Austromus australis</i>				X	W,U	D	X	W,U	D	X	U	D	X	W,U	D	X	U	D	X	W		X	U	D
<b>Mammals</b>	<b>Little Bentwing-bat</b>	<b><i>Miniopterus australis</i></b>		V					X	U	D	X	U	PO				X	U	D	X	U	PO	X	U	PO
<b>Mammals</b>	<b>Eastern Bentwing-bat</b>	<b><i>Miniopterus schreibersii oceanensis</i></b>		V		X	U	D	X	U	D	X	U	D	X	U	D	X	U	D	X	U	PO	X	U	PO
Mammals	Lesser Long-eared Bat	<i>Nyctophilus geoffroyi</i>							X	T																
Mammals	Long-eared Bat	<i>Nyctophilus sp.</i>				X	U	PO	X	U	PO	X	U	PO	X	U	PO							X	U	PO
<b>Mammals</b>	<b>Large-eared Pied Bat</b>	<b><i>Chalinolobus dwyeri</i></b>		V	V	X	U	D	X	U	D	X	U	D	X	U	D							X	U	PR
Mammals	Gould's Wattled Bat	<i>Chalinolobus gouldii</i>				X	U	D	X	T,U	D	X	U	D	X	U	D	X	U	D	X	T		X	U	D
Mammals	Chocolate Wattled Bat	<i>Chalinolobus morio</i>				X	U	D	X	T,U	D	X	U	D	X	U	D	X	U	D	X	U	PO	X	U	D
<b>Mammals</b>	<b>Eastern False Pipistrelle</b>	<b><i>Falsistrellus tasmaniensis</i></b>		V		X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO
<b>Mammals</b>	<b>Southern Myotis</b>	<b><i>Myotis macropus</i></b>		V		X	U	PO	X	U	PO	X	U	PO	X	U	PO							X	U	PO
<b>Mammals</b>	<b>Greater Broad-nosed Bat</b>	<b><i>Scoteanax rueppellii</i></b>		V		X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO
Mammals	Inland Broad-nosed Bat	<i>Scotorepens balstoni</i>				X	U	D	X	U	D	X	U	D	X	U	D	X	U	D	X	U	PO	X	U	D
Mammals	Eastern Broad-nosed Bat	<i>Scotorepens orion</i>				X	U,T	D	X	U	D	X	U	D	X	U	D	X	U	D	X	U	D	X	U	D
Mammals	A Broad-nosed Bat	<i>Scotorepens sp.</i>				X	O																			
Mammals	Large Forest Bat	<i>Vespadelus darlingtoni</i>																								
Mammals	Eastern Forest Bat	<i>Vespadelus pumilus</i>				X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO
Mammals	Southern Forest Bat	<i>Vespadelus regulus</i>				X	U	D	X	U	D	X	U	D	X	U	D	X	U	PO	X	U	PO	X	U	PO
<b>Mammals</b>	<b>Eastern Cave Bat</b>	<b><i>Vespadelus troughtoni</i></b>		V		X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO	X	U	PO
Mammals	Little Forest Bat	<i>Vespadelus vulturnus</i>				X	T		X	T	D	X	T		X	U	PR	X	U	PO	X	T		X	U	PO
Mammals	House Mouse	<i>Mus musculus</i>	X						X	T														X	O	
Mammals	a rodent	<i>Family Muridae</i>				X	H	PR																		
Mammals	Dingo	<i>Canis lupus dingo</i>				X	O								X	O										
Mammals	Dog	<i>Canis lupus familiaris</i>	X			X	P					X	P	D	X	P								X	X	D
Mammals	Hybrid Dog	<i>Canis lupus/familiaris</i>	X									X	W		X	P										
Mammals	Fox	<i>Vulpes vulpes</i>	X			X	Q,O,H,P	D	X	P	PO	X	P	PO	X	P		X	O					X	P	PO
Mammals	Cat	<i>Felis catus</i>	X			X	O		X	H	PR	X	H	PR				X	H	D						
Mammals	Brown Hare	<i>Lepus capensis</i>	X									X	Q,O		X	O		X	O		X	O				
Mammals	Rabbit	<i>Oryctolagus cuniculus</i>	X			X	O		X	O		X	O,X	D	X	O		X	O		X	O		X	O	
Mammals	Horse	<i>Equus caballus</i>	X																							

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 1	Observation Type	Confidence Level	Site 2	Observation Type	Confidence Level	Site 3	Observation Type	Confidence Level	Site 4	Observation Type	Confidence Level	Site 5	Observation Type	Confidence Level	Site 6	Observation Type	Confidence Level	Site 7	Observation Type	Confidence Level
						X	O		X	Q		X	Q		X	O		X	O		X	O		X	O	
Mammals	Pig	<i>Sus scrofa</i>	X			X	O																			
Mammals	European Cattle	<i>Bos taurus</i>	X			X	O		X	Q		X	Q		X	O		X	O		X	O		X	O	

Sites 8-13

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 8	Observation Type	Confidence Level	Site 9	Observation Type	Confidence Level	Site 10	Observation Type	Confidence Level	Site 11	Observation Type	Confidence Level	Bat Roost Tree	Observation Type	Confidence Level	Site 12	Observation Type	Confidence Level	Site 13	Observation Type	Confidence Type
Amphibians	Common Eastern Froglet	<i>Crinia signifera</i>																								
Amphibians	Spotted Grass Frog	<i>Limnodynastes tasmaniensis</i>										X	W													
Amphibians	Dusky Toadlet	<i>Uperoleia fusca</i>																								
Amphibians	Smooth Toadlet	<i>Uperoleia laevigata</i>																								
Amphibians	Green Tree Frog	<i>Litoria caerulea</i>										X	O		X	O										
Amphibians	Eastern Dwarf Tree Frog	<i>Litoria fallax</i>																								
Amphibians	Broad-palmed Frog	<i>Litoria latopalmata</i>										X	OW		X	O										
Amphibians	Peron's Tree Frog	<i>Litoria peronii</i>										X	O		X	W										
Reptiles	Eastern Snake-necked Turtle	<i>Chelodina longicollis</i>										X	O		X	O										
Reptiles	Macquarie Turtle	<i>Emydura macquarii</i>																								
Reptiles	Eastern Stone Gecko	<i>Diplodactylus vittatus</i>																								
Reptiles	Robust Velvet Gecko	<i>Nebulifera robusta</i>													X	O										
Reptiles	Thick-tailed Gecko	<i>Underwoodisaurus milii</i>													X	O										
Reptiles	<b>Pink-tailed Legless Lizard</b>	<b><i>Aprasia parapulchella</i></b>		V	V																					
Reptiles	<b>Striped Legless Lizard</b>	<b><i>Delma impar</i></b>		V	V										X	O, H										
Reptiles	Two-clawed Worm-skink	<i>Anomalopus leuckartii</i>																								
Reptiles	Southern Rainbow-skink	<i>Carlia tetradactyla</i>										X	O		X	O										
Reptiles	Elegant Snake-eyed Skink	<i>Cryptoblepharus pulcher</i>																								
Reptiles	Robust Ctenotus	<i>Ctenotus robustus</i>				X	O					X	O		X	O										
Reptiles	Tree Skink	<i>Egernia striolata</i>										X	O		X	O										
Reptiles	Barred-sided Skink	<i>Concinnia tenuis</i>																								
Reptiles	Eastern Ranges Rock-skink	<i>Liopholis modesta</i>													X	O										
Reptiles	South-eastern Morethia Skink	<i>Morethia boulengeri</i>																								
Reptiles	Eastern Blue-tongue	<i>Tiliqua scincoides</i>				X	O																			
Reptiles	Eastern Water Dragon	<i>Intellagama lesueurii</i>																								
Reptiles	Eastern Bearded Dragon	<i>Pogona barbata</i>																								
Reptiles	Sand Goanna	<i>Varanus gouldii</i>																								
Reptiles	Lace Monitor	<i>Varanus varius</i>													X	O										
Reptiles	Brown-snouted Blind Snake	<i>Anilius wiedii</i>																								
Reptiles	Spotted Black Snake	<i>Pseudechis guttatus</i>																								
Birds	Stubble Quail	<i>Coturnix pectoralis</i>																								
Birds	Brown Quail	<i>Coturnix ypsilophora</i>																								
Birds	Plumed Whistling Duck	<i>Dendrocygna eytoni</i>										X	OW													
Birds	Black Swan	<i>Cygnus atratus</i>																								

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 8	Observation Type	Confidence Level	Site 9	Observation Type	Confidence Level	Site 10	Observation Type	Confidence Level	Site 11	Observation Type	Confidence Level	Bat Roost Tree	Observation Type	Confidence Level	Site 12	Observation Type	Confidence Level	Site 13	Observation Type	Confidence Level
Birds	Pink-eared Duck	<i>Malacorhynchus membranaceus</i>													X	O										
Birds	Australian Wood Duck	<i>Chenonetta jubata</i>										X	O		X	O										
Birds	Pacific Black Duck	<i>Anas superciliosa</i>										X	O		X	O										
Birds	Australasian Shoveler	<i>Anas rhynchotis</i>													X	O										
Birds	Grey Teal	<i>Anas gracilis</i>										X	O		X	O										
Birds	Chestnut Teal	<i>Anas castanea</i>																								
Birds	Hardhead Duck	<i>Aythya australis</i>																								
Birds	Musk Duck	<i>Biziura lobata</i>																								
Birds	Australasian Grebe	<i>Tachybaptus novaehollandiae</i>										X	O		X	O	X									
Birds	Hoary-headed Grebe	<i>Poliiocephalus poliocephalus</i>																								
Birds	Straw-necked Ibis	<i>Threskiornis spinicollis</i>																								
Birds	Nankeen Night Heron	<i>Nycticorax caledonicus</i>																								
<b>Birds</b>	<b>Cattle Egret</b>	<b><i>Ardea ibis</i></b>			<b>M</b>																			X	O	
Birds	White-necked Heron	<i>Ardea pacifica</i>																								
Birds	White-faced Heron	<i>Egretta novaehollandiae</i>							X	O												X	O	X	O	
Birds	Little Pied Cormorant	<i>Microcarbo melanoleucos</i>																								
Birds	Pied Cormorant	<i>Phalacrocorax varius</i>																								
Birds	Australasian Darter	<i>Anhinga novaehollandiae</i>																								
Birds	Nankeen Kestrel	<i>Falco cenchroides</i>				X	O					X	O		X	O						X	O			
Birds	Brown Falcon	<i>Falco berigora</i>																						X	O	
Birds	Black-shouldered Kite	<i>Elanus axillaris</i>																								
Birds	Black Kite	<i>Milvus migrans</i>																								
<b>Birds</b>	<b>Square-tailed Kite</b>	<b><i>Lophoictinia isura</i></b>		<b>V</b>																						
<b>Birds</b>	<b>White-bellied Sea Eagle</b>	<b><i>Haliaeetus leucogaster</i></b>		<b>V</b>	<b>M</b>																					
<b>Birds</b>	<b>Spotted Harrier</b>	<b><i>Circus assimilis</i></b>		<b>V</b>																						
Birds	Brown Goshawk	<i>Accipiter fasciatus</i>																								
Birds	Collared Sparrowhawk	<i>Accipiter cirrocephalus</i>													X	O										
Birds	Wedge-tailed Eagle	<i>Aquila audax</i>										X	O		X	O						X	O			
<b>Birds</b>	<b>Little Eagle</b>	<b><i>Hieraetus morphnoides</i></b>		<b>V</b>								<b>X</b>	<b>O</b>													
Birds	Purple Swamphen	<i>Porphyrio porphyrio</i>																								
Birds	Eurasian Coot	<i>Fulica atra</i>																								
Birds	Black-winged Stilt	<i>Himantopus himantopus</i>																								
Birds	Red-necked Avocet	<i>Recurvirostra novaehollandiae</i>																								
Birds	Banded Lapwing	<i>Vanellus tricolor</i>																								
Birds	Masked Lapwing	<i>Vanellus miles</i>																								
Birds	Black-fronted Dotterel	<i>Elsayornis melanops</i>																								
Birds	Common Bronzewing	<i>Phaps chalcoptera</i>													X	O										
Birds	Crested Pigeon	<i>Ocyphaps lophotes</i>							X	W		X	O											X	O	
Birds	Bar-shouldered Dove	<i>Geopelia humeralis</i>																								
<b>Birds</b>	<b>Glossy Black-Cockatoo</b>	<b><i>Calyptorhynchus lathami</i></b>		<b>V</b>																						
Birds	Yellow-tailed Black-Cockatoo	<i>Calyptorhynchus funereus</i>																								
Birds	Galah	<i>Eolophus roseicapillus</i>				X	O		X	O		X	O		X	O						X	O	X	O	
Birds	Little Corella	<i>Cacatua sanguinea</i>																								
Birds	Sulphur-crested Cockatoo	<i>Cacatua galerita</i>																						X	O	
Birds	Rainbow Lorikeet	<i>Trichoglossus haematodus</i>																								
Birds	Musk Lorikeet	<i>Glossopsitta concinna</i>													X	O										

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 8	Observation Type	Confidence Level	Site 9	Observation Type	Confidence Level	Site 10	Observation Type	Confidence Level	Site 11	Observation Type	Confidence Level	Bat Roost Tree	Observation Type	Confidence Level	Site 12	Observation Type	Confidence Level	Site 13	Observation Type	Confidence Level
Birds	Little Lorikeet	<i>Glossopsitta pusilla</i>		V											X	O										
Birds	Crimson Rosella	<i>Platycercus elegans</i>																								
Birds	Eastern Rosella	<i>Platycercus eximius</i>							X	O		X	O		X	O					X	O		X	O	
Birds	Australian King-Parrot	<i>Alisterus scapularis</i>																								
Birds	Pallid Cuckoo	<i>Cuculus pallidus</i>																								
Birds	Fan-tailed Cuckoo	<i>Cacomantis flabelliformis</i>													X	O										
Birds	Black-eared Cuckoo	<i>Chalcites osculans</i>																								
Birds	Horsfield's Bronze-Cuckoo	<i>Chalcites basalys</i>																								
Birds	Shining Bronze Cuckoo	<i>Chrysococcyx lucidus</i>																								
Birds	Eastern Koel	<i>Eudynamys orientalis</i>										X	O		X	O										
Birds	Channel-billed Cuckoo	<i>Scythrops novaehollandiae</i>													X	O										
Birds	Eastern Barn Owl	<i>Tyto javanica</i>																								
Birds	Southern Boobook	<i>Ninox novaeseelandiae</i>													X	O										
Birds	Tawny Frogmouth	<i>Podargus strigoides</i>													X	O										
Birds	White-throated Nightjar	<i>Eurostopodus mystacalis</i>																								
Birds	Australian Owlet-nightjar	<i>Aegotheles cristatus</i>													X	O										
Birds	Laughing Kookaburra	<i>Dacelo novaeguineae</i>													X	OW										
Birds	Sacred Kingfisher	<i>Todiramphus sanctus</i>																								
Birds	Rainbow Bee-eater	<i>Merops ornatus</i>			M										X	O										
Birds	Brown Treecreeper (eastern subspecies)	<i>Climacteris picumnus victoriae</i>		V											X	O										
Birds	Variegated Fairy-wren	<i>Malurus lamberti</i>																								
Birds	Superb Fairy-wren	<i>Malurus cyaneus</i>				X	O		X	O					X	O								X	O	
Birds	Spotted Pardalote	<i>Pardalotus punctatus</i>													X	O										
Birds	Striated Pardalote	<i>Pardalotus striatus</i>							X	O					X	O					X	O		X	O	
Birds	Speckled Warbler	<i>Chthonicola sagittata</i>		V											X	O								X	O	
Birds	Weebill	<i>Smicronis brevirostris</i>													X	O										
Birds	Western Gerygone	<i>Gerygone fusca</i>										X	O		X	O										
Birds	White-throated Gerygone	<i>Gerygone olivacea</i>													X	O										
Birds	Brown Thornbill	<i>Acanthiza pusilla</i>																								
Birds	Buff-rumped Thornbill	<i>Acanthiza reguloides</i>										X	O		X	O										
Birds	Yellow-rumped Thornbill	<i>Acanthiza chrysorrhoa</i>				X	O					X	O		X	O								X	O	
Birds	Yellow Thornbill	<i>Acanthiza nana</i>													X	O								X	O	
Birds	Striated Thornbill	<i>Acanthiza lineata</i>													X	O										
Birds	Yellow-faced Honeyeater	<i>Caligavis chrysops</i>													X	O										
Birds	Singing Honeyeater	<i>Lichenostomus virescens</i>																								
Birds	White-eared Honeyeater	<i>Nesoptilotis leucotis</i>																								
Birds	Fuscous Honeyeater	<i>Lichenostomus fuscus</i>													X	O										
Birds	White-plumed Honeyeater	<i>Lichenostomus penicillatus</i>													X	O								X	O	
Birds	Noisy Miner	<i>Manorina melanocephala</i>							X	O		X	O		X	O						X	O			
Birds	Blue-faced Honeyeater	<i>Entomyzon cyanotis</i>							X	O																
Birds	Black-chinned Honeyeater (eastern subspecies)	<i>Melithreptus gularis gularis</i>		V																						
Birds	Brown-headed Honeyeater	<i>Melithreptus brevirostris</i>													X	O										
Birds	White-naped Honeyeater	<i>Melithreptus lunatus</i>													X	O										
Birds	Noisy Friarbird	<i>Philemon corniculatus</i>													X	O					X	O				
Birds	Striped Honeyeater	<i>Plectorhyncha lanceolata</i>										X	O		X	O										
Birds	Spiny-cheeked Honeyeater	<i>Acanthagenys rufogularis</i>							X	O																

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 8	Observation Type	Confidence Level	Site 9	Observation Type	Confidence Level	Site 10	Observation Type	Confidence Level	Site 11	Observation Type	Confidence Level	Bat Roost Tree	Observation Type	Confidence Level	Site 12	Observation Type	Confidence Level	Site 13	Observation Type	Confidence Level
Birds	Red Wattlebird	<i>Anthochaera carunculata</i>																								
<b>Birds</b>	<b>Painted Honeyeater</b>	<b><i>Grantiella picta</i></b>		V	V																					
Birds	Eastern Spinebill	<i>Acanthorhynchus tenuirostris</i>																								
Birds	Scarlet Honeyeater	<i>Myzomela sanguinolenta</i>																								
Birds	Eastern Yellow Robin	<i>Eopsaltria australis</i>																								
Birds	Jacky Winter	<i>Microeca fascinans</i>													X	O										
Birds	Rose Robin	<i>Petroica rosea</i>																								
<b>Birds</b>	<b>Flame Robin</b>	<b><i>Petroica phoenicea</i></b>		V																						
<b>Birds</b>	<b>Scarlet Robin</b>	<b><i>Petroica boodang</i></b>		V											X	O										
Birds	Red-capped Robin	<i>Petroica goodenovii</i>													X	O										
<b>Birds</b>	<b>Grey-crowned Babbler (eastern subspecies)</b>	<b><i>Pomatostomus temporalis ssp temporalis</i></b>		V																						
<b>Birds</b>	<b>Varied Sittella</b>	<b><i>Daphoenositta chrysoptera</i></b>		V											X	O										
Birds	Golden Whistler	<i>Pachycephala pectoralis</i>							X	O					X	O								X	O	
Birds	Rufous Whistler	<i>Pachycephala rufiventris</i>							X	O					X	O								X	O	
Birds	Grey Shrike-thrush	<i>Colluricincla harmonica</i>																						X	O	
Birds	Grey Fantail	<i>Rhipidura albiscapa</i>										X	W		X	O										
Birds	Willie Wagtail	<i>Rhipidura leucophrys</i>				X	O		X	O					X	O								X	O	
Birds	Magpie-lark	<i>Grallina cyanoleuca</i>										X	O		X	O										
<b>Birds</b>	<b>Leaden Flycatcher</b>	<b><i>Myiagra rubecula</i></b>			M																					
Birds	Grey Butcherbird	<i>Cracticus torquatus</i>							X	O		X	O		X	O										
Birds	Pied Butcherbird	<i>Cracticus nigrogularis</i>							X	W		X	O		X	O					X	O		X	O	
Birds	Australian Magpie	<i>Cracticus tibicen</i>				X	O		X	O		X	O		X	O					X	O		X	O	
Birds	Pied Currawong	<i>Strepera graculina</i>													X	O					X	O		X	O	
Birds	Masked Woodswallow	<i>Artamus personatus</i>																								
Birds	White-browed Woodswallow	<i>Artamus superciliosus</i>				X	O																			
<b>Birds</b>	<b>Dusky Woodswallow</b>	<b><i>Artamus cyanopterus</i></b>		V																						
Birds	Cicadabird	<i>Coracina tenuirostris</i>																								
Birds	Black-faced Cuckoo-shrike	<i>Coracina novaehollandiae</i>										X	O		X	O										
Birds	Ground Cuckoo-shrike	<i>Coracina maxima</i>										X	O													
Birds	White-winged Triller	<i>Lalage sueurii</i>																								
Birds	Olive-backed Oriole	<i>Oriolus sagittatus</i>																								
Birds	Little Raven	<i>Corvus mellori</i>																								
Birds	Australian Raven	<i>Corvus coronoides</i>				X	O		X	O		X	OW		X	O					X	O		X	O	
Birds	White-winged Chough	<i>Corcorax melanorhamphos</i>										X	O		X	O										
Birds	Common Starling	<i>Sturnus vulgaris</i>	X						X	O											X	O				
Birds	Common Myna	<i>Sturnus tristis</i>	X						X	O																
Birds	White-backed Swallow	<i>Cheramoeca leucosterna</i>																								
Birds	Welcome Swallow	<i>Hirundo neoxena</i>				X	O																			
Birds	Fairy Martin	<i>Petrochelidon ariel</i>							X	E																
Birds	Tree Martin	<i>Petrochelidon nigricans</i>																								
Birds	Silvereye	<i>Zosterops lateralis</i>													X	O										
Birds	Australian Reed Warbler	<i>Acrocephalus australis</i>																								
Birds	Tawny Grassbird	<i>Megalurus timoriensis</i>																								
Birds	Rufous Songlark	<i>Cincloramphus mathewsi</i>																								
Birds	Brown Songlark	<i>Cincloramphus cruralis</i>																								
Birds	Golden-headed Cisticola	<i>Cisticola exilis</i>																								

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 8	Observation Type	Confidence Level	Site 9	Observation Type	Confidence Level	Site 10	Observation Type	Confidence Level	Site 11	Observation Type	Confidence Level	Bat Roost Tree	Observation Type	Confidence Level	Site 12	Observation Type	Confidence Level	Site 13	Observation Type	Confidence Level	
Birds	Horsfield's Bushlark	<i>Mirafrja javanica</i>																									
Birds	Mistletoebird	<i>Dicaeum hirundinaceum</i>							X	O		X	O		X	O											
Birds	Richard's Pipit	<i>Anthus novaeseelandiae</i>				X	O																				
Birds	Red-browed Finch	<i>Neochmia temporalis</i>																									
Birds	Zebra Finch	<i>Taeniopygia guttata</i>										X	O														
Birds	Double-barred Finch	<i>Taeniopygia bichenovii</i>																						X	O		
Mammals	Short-beaked Echidna	<i>Tachyglossus aculeatus</i>										X	P		X	O											
Mammals	Yellow-footed Antechinus	<i>Antechinus flavipes</i>																									
Mammals	Common Dunnart	<i>Sminthopsis murina</i>																									
Mammals	Common Wombat	<i>Vombatus ursinus</i>																						X	FB	PO	
Mammals	Sugar Glider	<i>Petaurus breviceps</i>										X	W														
<b>Mammals</b>	<b>Squirrel Glider</b>	<b><i>Petaurus norfolcensis</i></b>		V											X	O						X	M	PO			
Mammals	Common Ringtail Possum	<i>Pseudocheirus peregrinus</i>																									
Mammals	Common Brushtail Possum	<i>Trichosurus vulpecula</i>										X	O		X	O								X	X	PR	
Mammals	brushtail possum	<i>Trichosurus sp.</i>																									
Mammals	Eastern Grey Kangaroo	<i>Macropus giganteus</i>				X	O					X	O		X	O						X	O	X	O		
Mammals	Eastern Wallaroo	<i>Macropus robustus</i>																				X	O				
Mammals	Red-necked Wallaby	<i>Macropus rufogriseus</i>													X	O											
Mammals	Swamp Wallaby	<i>Wallabia bicolor</i>																									
<b>Mammals</b>	<b>Grey-headed Flying-fox</b>	<b><i>Pteropus poliocephalus</i></b>		V	V										X	W											
<b>Mammals</b>	<b>Yellow-bellied Sheath-tail-bat</b>	<b><i>Saccolaimus flaviventris</i></b>		V																							
<b>Mammals</b>	<b>Eastern Freetail-bat</b>	<b><i>Mormopterus norfolkensis</i></b>		V								X	U	PR													
Mammals	Little Mastiff-bat	<i>Mormopterus planiceps</i>										X	U	D	X	U	D	X	U	D							
Mammals	Eastern Free-tailed Bat	<i>Mormopterus ridei</i>													X	U	PO										
Mammals	White-striped Freetail-bat	<i>Austronomus australis</i>										X	U	D	X	U	D										
<b>Mammals</b>	<b>Little Bentwing-bat</b>	<b><i>Miniopterus australis</i></b>		V								X	U	D	X	U	D										
<b>Mammals</b>	<b>Eastern Bentwing-bat</b>	<b><i>Miniopterus schreibersii oceanensis</i></b>		V								X	U	D	X	U	D										
Mammals	Lesser Long-eared Bat	<i>Nyctophilus geoffroyi</i>																									
Mammals	Long-eared Bat	<i>Nyctophilus sp.</i>										X	U	PO	X	U	PO										
<b>Mammals</b>	<b>Large-eared Pied Bat</b>	<b><i>Chalinolobus dwyeri</i></b>		V	V							X	U	D	X	U	D										
Mammals	Gould's Wattled Bat	<i>Chalinolobus gouldii</i>										X	U	PO	X	U	D	X	U	PO							
Mammals	Chocolate Wattled Bat	<i>Chalinolobus morio</i>										X	U	D	X	U	D										
<b>Mammals</b>	<b>Eastern False Pipistrelle</b>	<b><i>Falsistrellus tasmaniensis</i></b>		V								X	U	PO													
<b>Mammals</b>	<b>Southern Myotis</b>	<b><i>Myotis macropus</i></b>		V								X	U/O	PO/D	X	U	PO										
<b>Mammals</b>	<b>Greater Broad-nosed Bat</b>	<b><i>Scoteanax rueppellii</i></b>		V								X	U	PO	X	U	PO										
Mammals	Inland Broad-nosed Bat	<i>Scotorepens balstoni</i>										X	U	PR	X	U	D	X	U	D							
Mammals	Eastern Broad-nosed Bat	<i>Scotorepens orion</i>										X	U	PO	X	U	PO										
Mammals	A Broad-nosed Bat	<i>Scotorepens sp.</i>																									
Mammals	Large Forest Bat	<i>Vespadelus darlingtoni</i>													X	U	PO										
Mammals	Eastern Forest Bat	<i>Vespadelus pumilus</i>										X	U	PO	X	U	D										
Mammals	Southern Forest Bat	<i>Vespadelus regulus</i>										X	U	PO	X	U	D										
<b>Mammals</b>	<b>Eastern Cave Bat</b>	<b><i>Vespadelus troungtoni</i></b>		V								X	U	PO	X	U	PO										
Mammals	Little Forest Bat	<i>Vespadelus vulturinus</i>										X	U	PO	X	U	PO										
Mammals	House Mouse	<i>Mus musculus</i>	X																								
Mammals	a rodent	Family Muridae																									
Mammals	Dingo	<i>Canis lupus dingo</i>																									

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 8	Observation Type	Confidence Level	Site 9	Observation Type	Confidence Level	Site 10	Observation Type	Confidence Level	Site 11	Observation Type	Confidence Level	Bat Roost Tree	Observation Type	Confidence Level	Site 12	Observation Type	Confidence Level	Site 13	Observation Type	Confidence Level
Mammals	Dog	<i>Canis lupus familiaris</i>	X												X	P					X	P				
Mammals	Hybrid Dog	<i>Canis lupus/familiaris</i>	X												X	P					X					
Mammals	Fox	<i>Vulpes vulpes</i>	X												X	O, P	D				X	O		X	P	
Mammals	Cat	<i>Felis catus</i>	X									X	O		X	O										
Mammals	Brown Hare	<i>Lepus capensis</i>	X																							
Mammals	Rabbit	<i>Oryctolagus cuniculus</i>	X									X	O		X	O, X	D				X	P, X, Y	D	X	FB	
Mammals	Horse	<i>Equus caballus</i>	X																		X	O				
Mammals	Pig	<i>Sus scrofa</i>	X												X	O										
Mammals	European Cattle	<i>Bos taurus</i>	X												X	X	D				X	O		X	O	

Sites 14 - Incidental

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 14	Observation Type	Confidence Type	Site 15	Observation Type	Confidence Type	Site 16	Observation Type	Confidence Type	Site 17	Observation Type	Confidence Type	Site 18	Observation Type	Confidence Type	Small rock escarpment Main Entrance	Observation Type	Confidence Type	Savoy Dam	Observation Type	Confidence Type	Incidental	Observation Type
Amphibians	Common Eastern Froglet	<i>Crinia signifera</i>							X	W		X	W		X	W											X	W
Amphibians	Spotted Grass Frog	<i>Limnodynastes tasmaniensis</i>													X	OW		X	OW								X	O
Amphibians	Dusky Toadlet	<i>Uperoleia fusca</i>							X	W																		
Amphibians	Smooth Toadlet	<i>Uperoleia laevisgata</i>																										
Amphibians	Green Tree Frog	<i>Litoria caerulea</i>				X	O											X	O								X	OW
Amphibians	Eastern Dwarf Tree Frog	<i>Litoria fallax</i>													X	OW												
Amphibians	Broad-palmed Frog	<i>Litoria latopalmeta</i>				X	O					X	O		X	W		X	W								X	O
Amphibians	Peron's Tree Frog	<i>Litoria peronii</i>													X	W		X	O									
Reptiles	Eastern Snake-necked Turtle	<i>Chelodina longicollis</i>				X	O																				X	O
Reptiles	Macquarie Turtle	<i>Emydura macquarii</i>																									X	O
Reptiles	Eastern Stone Gecko	<i>Diplodactylus vittatus</i>													X	O		X	O									
Reptiles	Robust Velvet Gecko	<i>Nebulifera robusta</i>																			X	O					X	O
Reptiles	Thick-tailed Gecko	<i>Underwoodisaurus milii</i>																										
<b>Reptiles</b>	<b>Pink-tailed Legless Lizard</b>	<b><i>Aprasia parapulchella</i></b>		V	V																							
<b>Reptiles</b>	<b>Striped Legless Lizard</b>	<b><i>Delma impar</i></b>		V	V	X	O																				X	O
Reptiles	Two-clawed Worm-skink	<i>Anomalopus leuckartii</i>																										
Reptiles	Southern Rainbow-skink	<i>Carlia tetradactyla</i>							X	O								X	O		X	O					X	O
Reptiles	Elegant Snake-eyed Skink	<i>Cryptoblepharus pulcher</i>													X	O												
Reptiles	Robust Ctenotus	<i>Ctenotus robustus</i>				X	O		X	O		X	O		X	O		X	O								X	O
Reptiles	Tree Skink	<i>Egernia striolata</i>													X	O		X	O		X	O					X	O
Reptiles	Barred-sided Skink	<i>Concinnia tenuis</i>																										
Reptiles	Eastern Ranges Rock-skink	<i>Liopholis modesta</i>																			X	O					X	O
Reptiles	South-eastern Morethia Skink	<i>Morethia boulengeri</i>																										
Reptiles	Eastern Blue-tongue	<i>Tiliqua scincoides</i>																										
Reptiles	Eastern Water Dragon	<i>Intellagama lesueurii</i>							X	O																		
Reptiles	Eastern Bearded Dragon	<i>Pogona barbata</i>							X	O																	X	O
Reptiles	Sand Goanna	<i>Varanus gouldii</i>																										

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 14	Observation Type	Confidence Type	Site 15	Observation Type	Confidence Type	Site 16	Observation Type	Confidence Type	Site 17	Observation Type	Confidence Type	Site 18	Observation Type	Confidence Type	Small rock escarpment Main Entrance	Observation Type	Confidence Type	Savoy Dam	Observation Type	Confidence Type	Incidental	Observation Type
Reptiles	Lace Monitor	<i>Varanus varius</i>													X	O											X	O, Q
Reptiles	Brown-snouted Blind Snake	<i>Anilius wiedii</i>																										
Reptiles	Spotted Black Snake	<i>Pseudechis guttatus</i>				X	O					X	O															
Birds	Stubble Quail	<i>Coturnix pectoralis</i>																						X	O		X	O
Birds	Brown Quail	<i>Coturnix ypsilophora</i>				X	O																					
Birds	Plumed Whistling Duck	<i>Dendrocygna eytoni</i>																										
Birds	Black Swan	<i>Cygnus atratus</i>										X	O															
Birds	Pink-eared Duck	<i>Malacorhynchus membranaceus</i>																										
Birds	Australian Wood Duck	<i>Chenonetta jubata</i>				X	O					X	OW					X	O					X	O		X	O
Birds	Pacific Black Duck	<i>Anas superciliosa</i>				X	O		X	O														X	O		X	O
Birds	Australasian Shoveler	<i>Anas rhynchotis</i>										X	O															
Birds	Grey Teal	<i>Anas gracilis</i>							X	O		X	O											X	O		X	O
Birds	Chestnut Teal	<i>Anas castanea</i>																										
Birds	Hardhead Duck	<i>Aythya australis</i>										X	O														X	O
Birds	Musk Duck	<i>Biziura lobata</i>										X	O														X	O
Birds	Australasian Grebe	<i>Tachybaptus novaehollandiae</i>							X	O														X	O		X	O
Birds	Hoary-headed Grebe	<i>Poliiocephalus poliocephalus</i>							X	O		X	O														X	O
Birds	Straw-necked Ibis	<i>Threskiornis spinicollis</i>																									X	O
Birds	Nankeen Night Heron	<i>Nycticorax caledonicus</i>																										
<b>Birds</b>	<b>Cattle Egret</b>	<b><i>Ardea ibis</i></b>			<b>M</b>																							
Birds	White-necked Heron	<i>Ardea pacifica</i>																						X	O		X	O
Birds	White-faced Heron	<i>Egretta novaehollandiae</i>							X	O		X	O					X	O					X	O		X	O
Birds	Little Pied Cormorant	<i>Microcarbo melanoleucos</i>										X	O															
Birds	Pied Cormorant	<i>Phalacrocorax varius</i>							X	O																		
Birds	Australasian Darter	<i>Anhinga novaehollandiae</i>													X	O												
Birds	Nankeen Kestrel	<i>Falco cenchroides</i>				X	O		X	O					X	O												
Birds	Brown Falcon	<i>Falco berigora</i>				X	O											X	O					X	O		X	O
Birds	Black-shouldered Kite	<i>Elanus axillaris</i>																									X	O
Birds	Black Kite	<i>Milvus migrans</i>																									X	O
<b>Birds</b>	<b>Square-tailed Kite</b>	<b><i>Lophoictinia isura</i></b>		<b>V</b>																								
<b>Birds</b>	<b>White-bellied Sea Eagle</b>	<b><i>Haliaeetus leucogaster</i></b>		<b>V</b>	<b>M</b>				X	O																		
<b>Birds</b>	<b>Spotted Harrier</b>	<b><i>Circus assimilis</i></b>		<b>V</b>														X	O								X	O
Birds	Brown Goshawk	<i>Accipiter fasciatus</i>																									X	O
Birds	Collared Sparrowhawk	<i>Accipiter cirrocephalus</i>																										
Birds	Wedge-tailed Eagle	<i>Aquila audax</i>							X	O		X	O					X	O					X	O		X	O
<b>Birds</b>	<b>Little Eagle</b>	<b><i>Hieraetus morphnoides</i></b>		<b>V</b>																							X	O
Birds	Purple Swamphen	<i>Porphyrio porphyrio</i>							X	O																		
Birds	Eurasian Coot	<i>Fulica atra</i>							X	O		X	O											X	O		X	O
Birds	Black-winged Stilt	<i>Himantopus himantopus</i>										X	O															
Birds	Red-necked Avocet	<i>Recurvirostra novaehollandiae</i>										X	O															
Birds	Banded Lapwing	<i>Vanellus tricolor</i>																									X	O
Birds	Masked Lapwing	<i>Vanellus miles</i>										X	OW		X	O		X	O					X	O		X	O
Birds	Black-fronted Dotterel	<i>Euseyonis melanops</i>							X	O		X	O															
Birds	Common Bronzewing	<i>Phaps chalcoptera</i>																										
Birds	Crested Pigeon	<i>Ocyphaps lophotes</i>				X	O		X	O								X	O									
Birds	Bar-shouldered Dove	<i>Geopelia humeralis</i>																										



MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 14	Observation Type	Confidence Type	Site 15	Observation Type	Confidence Type	Site 16	Observation Type	Confidence Type	Site 17	Observation Type	Confidence Type	Site 18	Observation Type	Confidence Type	Small rock escarpment Main Entrance	Observation Type	Confidence Type	Savoy Dam	Observation Type	Confidence Type	Incidental	Observation Type
<b>Birds</b>	<b>Glossy Black-Cockatoo</b>	<b><i>Calyptorhynchus lathami</i></b>		V								X	G															
Birds	Yellow-tailed Black-Cockatoo	<i>Calyptorhynchus funereus</i>													X	O												
Birds	Galah	<i>Eolophus roseicapillas</i>				X	O		X	O					X	O		X	O					X	O		X	O
Birds	Little Corella	<i>Cacatua sanguinea</i>																										
Birds	Sulphur-crested Cockatoo	<i>Cacatua galerita</i>							X	O		X	OW		X	O		X	O									
Birds	Rainbow Lorikeet	<i>Trichoglossus haematodus</i>										X	OW															
Birds	Musk Lorikeet	<i>Glossopsitta concinna</i>				X	O		X	OW																		
<b>Birds</b>	<b>Little Lorikeet</b>	<b><i>Glossopsitta pusilla</i></b>		V																								
Birds	Crimson Rosella	<i>Platycercus elegans</i>										X	O															
Birds	Eastern Rosella	<i>Platycercus eximius</i>				X	O		X	O		X	O		X	O		X	O					X	O		X	O
Birds	Australian King-Parrot	<i>Alisterus scapularis</i>				X	O					X	O														X	W
Birds	Pallid Cuckoo	<i>Cuculus pallidus</i>																									X	O
Birds	Fan-tailed Cuckoo	<i>Cacomantis flabelliformis</i>							X	O																		
Birds	Black-eared Cuckoo	<i>Chalcites osculans</i>																										
Birds	Horsfield's Bronze-Cuckoo	<i>Chalcites basalus</i>													X	O											X	O
Birds	Shining Bronze Cuckoo	<i>Chrysococcyx lucidus</i>																										
Birds	Eastern Koel	<i>Eudynamis orientalis</i>																X	O									
Birds	Channel-billed Cuckoo	<i>Scythrops novaehollandiae</i>													X	O		X	O									
Birds	Eastern Barn Owl	<i>Tyto javanica</i>							X	O																		
Birds	Southern Boobook	<i>Ninox novaeseelandiae</i>																										
Birds	Tawny Frogmouth	<i>Podargus strigoides</i>																X	O								X	O
Birds	White-throated Nightjar	<i>Eurostopodus mystacalis</i>																									X	OW
Birds	Australian Owlet-nightjar	<i>Aegotheles cristatus</i>				X	O					X	W														X	O
Birds	Laughing Kookaburra	<i>Dacelo novaeguineae</i>				X	O								X	OW		X	O									
Birds	Sacred Kingfisher	<i>Todiramphus sanctus</i>																X	W									
<b>Birds</b>	<b>Rainbow Bee-eater</b>	<b><i>Merops ornatus</i></b>			M										X	O												
<b>Birds</b>	<b>Brown Treecreeper (eastern subspecies)</b>	<b><i>Climacteris picumnus victoriae</i></b>		V																								
Birds	Variiegated Fairy-wren	<i>Malurus lamberti</i>																										
Birds	Superb Fairy-wren	<i>Malurus cyaneus</i>				X	O		X	O		X	O		X	O		X	O								X	O
Birds	Spotted Pardalote	<i>Pardalotus punctatus</i>							X	OW		X	OW		X	O												
Birds	Striated Pardalote	<i>Pardalotus striatus</i>							X	OW		X	OW		X	O		X	W									
<b>Birds</b>	<b>Speckled Warbler</b>	<b><i>Chthonicola sagittata</i></b>		V					X	O																		
Birds	Weebill	<i>Smicromis brevirostris</i>							X	OW		X	OW					X	O								X	W
Birds	Western Gerygone	<i>Gerygone fusca</i>							X	OW																		
Birds	White-throated Gerygone	<i>Gerygone olivacea</i>																										
Birds	Brown Thornbill	<i>Acanthiza pusilla</i>							X	OW																		
Birds	Buff-rumped Thornbill	<i>Acanthiza reguloides</i>							X	OW																		
Birds	Yellow-rumped Thornbill	<i>Acanthiza chrysorrhoa</i>				X	O		X	O								X	O								X	O
Birds	Yellow Thornbill	<i>Acanthiza nana</i>				X	O		X	O		X	O					X	O									
Birds	Striated Thornbill	<i>Acanthiza lineata</i>																										
Birds	Yellow-faced Honeyeater	<i>Caligavis chrysops</i>							X	OW		X	OW															
Birds	Singing Honeyeater	<i>Lichenostomus virescens</i>							X	O																		
Birds	White-eared Honeyeater	<i>Nesoptilotis leucotis</i>										X	OW															
Birds	Fuscous Honeyeater	<i>Lichenostomus fuscus</i>										X	OW															
Birds	White-plumed Honeyeater	<i>Lichenostomus penicillatus</i>													X	OW												
Birds	Noisy Miner	<i>Manorina melanocephala</i>							X	OW		X	OW		X	O		X	O									

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 14	Observation Type	Confidence Type	Site 15	Observation Type	Confidence Type	Site 16	Observation Type	Confidence Type	Site 17	Observation Type	Confidence Type	Site 18	Observation Type	Confidence Type	Small rock escarpment Main Entrance	Observation Type	Confidence Type	Savoy Dam	Observation Type	Confidence Type	Incidental	Observation Type
Birds	Blue-faced Honeyeater	<i>Entomyzon cyanotis</i>																										
<b>Birds</b>	<b>Black-chinned Honeyeater (eastern subspecies)</b>	<b><i>Melithreptus gularis gularis</i></b>		V																								
Birds	Brown-headed Honeyeater	<i>Melithreptus brevirostris</i>							X	OW		X	OW															
Birds	White-naped Honeyeater	<i>Melithreptus lunatus</i>							X	OW		X	OW															
Birds	Noisy Friarbird	<i>Philemon corniculatus</i>				X	O		X	OW		X	OW		X	O		X	O									
Birds	Striped Honeyeater	<i>Plectorhyncha lanceolata</i>				X	O											X	O									
Birds	Spiny-cheeked Honeyeater	<i>Acanthagenys rufogularis</i>				X	O		X	O					X	O		X	O									
Birds	Red Wattlebird	<i>Anthochaera carunculata</i>							X	O		X	OW															
<b>Birds</b>	<b>Painted Honeyeater</b>	<b><i>Grantiella picta</i></b>		V	V																							
Birds	Eastern Spinebill	<i>Acanthorhynchus tenuirostris</i>										X	OW															
Birds	Scarlet Honeyeater	<i>Myzomela sanguinolenta</i>																										
Birds	Eastern Yellow Robin	<i>Eopsaltria australis</i>																										
Birds	Jacky Winter	<i>Microeca fascians</i>																										
Birds	Rose Robin	<i>Petroica rosea</i>							X	O		X	O															
<b>Birds</b>	<b>Flame Robin</b>	<b><i>Petroica phoenicea</i></b>		V					X	O																		
<b>Birds</b>	<b>Scarlet Robin</b>	<b><i>Petroica boodang</i></b>		V																								
Birds	Red-capped Robin	<i>Petroica goodenovii</i>				X	O											X	O									
<b>Birds</b>	<b>Grey-crowned Babbler (eastern subspecies)</b>	<b><i>Pomatostomus temporalis ssp temporalis</i></b>		V					X	O								X	OW							X	O	
<b>Birds</b>	<b>Varied Sittella</b>	<b><i>Daphoenositta chrysoptera</i></b>		V					X	O																		
Birds	Golden Whistler	<i>Pachycephala pectoralis</i>							X	OW		X	OW					X	O									
Birds	Rufous Whistler	<i>Pachycephala rufiventris</i>				X	O		X	O					X	O		X	OW		X	O						
Birds	Grey Shrike-thrush	<i>Colluricincla harmonica</i>				X	O											X	O									
Birds	Grey Fantail	<i>Rhipidura albiscapa</i>				X	O		X	O		X	O															
Birds	Willie Wagtail	<i>Rhipidura leucophrys</i>							X	O					X	O		X	O					X	O	X	O	
Birds	Magpie-lark	<i>Grallina cyanoleuca</i>				X	O											X	O					X	O	X	E	
<b>Birds</b>	<b>Leaden Flycatcher</b>	<b><i>Myiagra rubecula</i></b>			M																							
Birds	Grey Butcherbird	<i>Cracticus torquatus</i>				X	O		X	OW		X	OW		X	O		X	O									
Birds	Pied Butcherbird	<i>Cracticus nigrogularis</i>				X	O		X	OW		X	OW		X	O		X	O					X	O	X	W	
Birds	Australian Magpie	<i>Cracticus tibicen</i>				X	O		X	OW		X	OW		X	O		X	O					X	O			
Birds	Pied Currawong	<i>Strepera graculina</i>				X	O		X	O		X	OW		X	OW		X	O									
Birds	Masked Woodswallow	<i>Artamus personatus</i>																										
Birds	White-browed Woodswallow	<i>Artamus superciliosus</i>																X	O									
<b>Birds</b>	<b>Dusky Woodswallow</b>	<b><i>Artamus cyanopterus</i></b>		V					X	O					X	O												
Birds	Cicadabird	<i>Coracina tenuirostris</i>																									X	W
Birds	Black-faced Cuckoo-shrike	<i>Coracina novaehollandiae</i>				X	O		X	O					X	O												
Birds	Ground Cuckoo-shrike	<i>Coracina maxima</i>																										
Birds	White-winged Triller	<i>Lalage sueurii</i>													X	O		X	O									
Birds	Olive-backed Oriole	<i>Oriolus sagittatus</i>							X	O					X	O												
Birds	Little Raven	<i>Corvus mellori</i>													X	O												
Birds	Australian Raven	<i>Corvus coronoides</i>				X	O		X	OW		X	OW		X	O		X	OW					X	O	X	Q	
Birds	White-winged Chough	<i>Corcorax melanorhamphos</i>				X	O		X	OW																		
Birds	Common Starling	<i>Sturnus vulgaris</i>	X			X	O																			X	O	
Birds	Common Myna	<i>Sturnus tristis</i>	X																							X	O	
Birds	White-backed Swallow	<i>Cheramoeca leucosterna</i>																								X	O	
Birds	Welcome Swallow	<i>Hirundo neoxena</i>				X	O		X	O		X	O		X	O												
Birds	Fairy Martin	<i>Petrochelidon ariel</i>																										

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 14	Observation Type	Confidence Type	Site 15	Observation Type	Confidence Type	Site 16	Observation Type	Confidence Type	Site 17	Observation Type	Confidence Type	Site 18	Observation Type	Confidence Type	Small rock escarpment Main Entrance	Observation Type	Confidence Type	Savoy Dam	Observation Type	Confidence Type	Incidental	Observation Type
Birds	Tree Martin	<i>Petrochelidon nigricans</i>				X	O					X	O					X	O								X	O
Birds	Silvereye	<i>Zosterops lateralis</i>							X	O		X	O															
Birds	Australian Reed Warbler	<i>Acrocephalus australis</i>																									X	OW
Birds	Tawny Grassbird	<i>Megalurus timoriensis</i>																										
Birds	Rufous Songlark	<i>Cincloramphus mathewsi</i>				X	O								X	O		X	O									
Birds	Brown Songlark	<i>Cincloramphus cruralis</i>																									X	OW
Birds	Golden-headed Cisticola	<i>Cisticola exilis</i>																						X	O		X	O
Birds	Horsfield's Bushlark	<i>Mirafrja javanica</i>																									X	O
Birds	Mistletoebird	<i>Dicaeum hirundinaceum</i>							X	OW					X	OW												
Birds	Richard's Pipit	<i>Anthus novaeseelandiae</i>				X	O																				X	O
Birds	Red-browed Finch	<i>Neochmia temporalis</i>													X	O												
Birds	Zebra Finch	<i>Taeniopygia guttata</i>																										
Birds	Double-barred Finch	<i>Taeniopygia bichenovii</i>				X	O											X	O									
Mammals	Short-beaked Echidna	<i>Tachyglossus aculeatus</i>																			X	P		X	O			
Mammals	Yellow-footed Antechinus	<i>Antechinus flavipes</i>																										
Mammals	Common Dunnart	<i>Sminthopsis murina</i>																										
Mammals	Common Wombat	<i>Vombatus ursinus</i>																									X	P
Mammals	Sugar Glider	<i>Petaurus breviceps</i>																										
<b>Mammals</b>	<b>Squirrel Glider</b>	<b><i>Petaurus norfolcensis</i></b>		<b>V</b>																								
Mammals	Common Ringtail Possum	<i>Pseudocheirus peregrinus</i>							X	E																		
Mammals	Common Brushtail Possum	<i>Trichosurus vulpecula</i>							X	O		X	O, H					X	O									
Mammals	brushtail possum	<i>Trichosurus sp.</i>																										
Mammals	Eastern Grey Kangaroo	<i>Macropus giganteus</i>				X	O		X	O		X	O		X	O		X	O					X	O			
Mammals	Eastern Wallaroo	<i>Macropus robustus</i>							X	O																		
Mammals	Red-necked Wallaby	<i>Macropus rufogriseus</i>							X	O		X	O		X	O												
Mammals	Swamp Wallaby	<i>Wallabia bicolor</i>							X	P	PR																X	X
<b>Mammals</b>	<b>Grey-headed Flying-fox</b>	<b><i>Pteropus poliocephalus</i></b>		<b>V</b>	<b>V</b>																							
<b>Mammals</b>	<b>Yellow-bellied Sheathtail-bat</b>	<b><i>Saccolaimus flaviventris</i></b>		<b>V</b>																								
<b>Mammals</b>	<b>Eastern Freetail-bat</b>	<b><i>Mormopterus norfolkensis</i></b>		<b>V</b>																								
Mammals	Little Mastiff-bat	<i>Mormopterus planiceps</i>																			X	U	D					
Mammals	Eastern Free-tailed Bat	<i>Mormopterus ridei</i>																										
Mammals	White-striped Freetail-bat	<i>Austronomus australis</i>																										
<b>Mammals</b>	<b>Little Bentwing-bat</b>	<b><i>Miniopterus australis</i></b>		<b>V</b>																								
<b>Mammals</b>	<b>Eastern Bentwing-bat</b>	<b><i>Miniopterus schreibersii oceanensis</i></b>		<b>V</b>																								
Mammals	Lesser Long-eared Bat	<i>Nyctophilus geoffroyi</i>																										
Mammals	Long-eared Bat	<i>Nyctophilus sp.</i>																										
<b>Mammals</b>	<b>Large-eared Pied Bat</b>	<b><i>Chalinolobus dwyeri</i></b>		<b>V</b>	<b>V</b>																							
Mammals	Gould's Wattled Bat	<i>Chalinolobus gouldii</i>																			X	U	D					
Mammals	Chocolate Wattled Bat	<i>Chalinolobus morio</i>																										
<b>Mammals</b>	<b>Eastern False Pipistrelle</b>	<b><i>Falsistrellus tasmaniensis</i></b>		<b>V</b>																								
<b>Mammals</b>	<b>Southern Myotis</b>	<b><i>Myotis macropus</i></b>		<b>V</b>																								
<b>Mammals</b>	<b>Greater Broad-nosed Bat</b>	<b><i>Scoteanax rueppellii</i></b>		<b>V</b>																								
Mammals	Inland Broad-nosed Bat	<i>Scotorepens balstoni</i>																			X	U	D					
Mammals	Eastern Broad-nosed Bat	<i>Scotorepens orion</i>																										
Mammals	A Broad-nosed Bat	<i>Scotorepens sp.</i>																										
Mammals	Large Forest Bat	<i>Vespadelus darlingtoni</i>																										

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Fauna Group	Common Name	Scientific Name	Introduced	NSW Status	Federal Status	Site 14	Observation Type	Confidence Type	Site 15	Observation Type	Confidence Type	Site 16	Observation Type	Confidence Type	Site 17	Observation Type	Confidence Type	Site 18	Observation Type	Confidence Type	Small rock escarpment Main Entrance	Observation Type	Confidence Type	Savoy Dam	Observation Type	Confidence Type	Incidental	Observation Type
Mammals	Eastern Forest Bat	<i>Vespadelus pumilus</i>																										
Mammals	Southern Forest Bat	<i>Vespadelus regulus</i>																										
<b>Mammals</b>	<b>Eastern Cave Bat</b>	<b><i>Vespadelus troughtoni</i></b>		<b>V</b>																								
Mammals	Little Forest Bat	<i>Vespadelus vulturnus</i>																										
Mammals	House Mouse	<i>Mus musculus</i>	X																									
Mammals	a rodent	<i>Family Muridae</i>																										
Mammals	Dingo	<i>Canis lupus dingo</i>																									X	O
Mammals	Dog	<i>Canis lupus familiaris</i>	X						X	P, F	D							X	P	D							X	P
Mammals	Hybrid Dog	<i>Canis lupus familiaris</i>	X																									
Mammals	Fox	<i>Vulpes vulpes</i>	X						X	P	PO	X	P	PO							X	F					X	Q, O
Mammals	Cat	<i>Felis catus</i>	X																								X	O
Mammals	Brown Hare	<i>Lepus capensis</i>	X			X	O											X	O								X	O
Mammals	Rabbit	<i>Oryctolagus cuniculus</i>	X			X	O		X	O					X	O		X	O		X	H	D	X	O, Y	D	X	O
Mammals	Horse	<i>Equus caballus</i>	X																									
Mammals	Pig	<i>Sus scrofa</i>	X																									
Mammals	European Cattle	<i>Bos taurus</i>	X			X	O					X	X	D				X	H	D							X	O, Q

Key

X: detected  
 O: observed  
 W: heard  
 U: ultrasonic call recorded (microbats)  
 H: hair sample  
 Q: captured on camera  
 T: trapped  
 XX: in a scat

D: Definite detection (for identification via hair or ultrasonic call)  
 Pr: Probable detection (for identification via hair or ultrasonic call)  
 Po: Possible detection (for identification via hair or ultrasonic call)  
 Bold type: listed threatened and/or protected migratory species  
 V: listed as vulnerable under the BC and/or EPBC Act  
 E: listed as endangered under the BC and/or EPBC Act  
 CE: listed as critically endangered under the BC and/or EPBC Act  
 M: listed as a migratory and/or marine species under the EPBC Act

## Appendix B Fauna Survey Site Descriptions

**Site Number:** 1

**Site Description:** Open eucalypt forest with a sparse shrub layer and mainly rocks and leaf litter for ground cover.

**Habitat Condition:** Evidence of heavy grazing and very dry conditions have left site in poor condition.

**Site Disturbance Level and Type:** Highly disturbed due to cattle grazing.

**Connectivity:** Connected to areas of woodland to the south through a narrow and broken corridor.

**Site Location:** -32.38643, 150.89076

**Patch Size:** 40 hectares

**Topography, Slope and Aspect:** Rocky hill with a moderately steep slope to the south-west.

**Soil Type:** Shallow, stoney brown clay loam.

**Canopy Height:** 10-15 m

**DBH Canopy Trees:** 0.4 to 1.2 m

**DBH Sub-canopy Trees:** 0.1 to 0.3 m

**Tree Hollow Density:** moderate

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Blakely's Red Gum - Narrow-leaved Ironbark - Rough-barked Apple shrubby woodland of the upper Hunter (PCT 1607).
- Blakely's Red Gum - Narrow-leaved Ironbark - Rough-barked Apple shrubby woodland of the upper Hunter – DNG (PCT 1607).
- Narrow-leaved Ironbark - Grey Box grassy woodland of the central and upper Hunter (PCT 1691).
- Dam.



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 2

**Site Description:** Dry open Box Gum woodland, with patches of dense regrowth of Bulloak and dense cover of leaf litter.

**Habitat Condition:** Poor very dry conditions with shrub layer with sparse foliage. The intermittent watercourse consists of an eroded gully.

**Site Disturbance Level and Type:** Some evidence of low disturbance through grazing.

**Connectivity:** Connected to the south and to the west, with cleared areas to the north and east.

**Site Location:** -32.39606, 150.89162

**Patch Size:** 55 hectares

**Topography, Slope and Aspect:** Gently sloping hillside to the east along drainage line.

**Soil Type:** Red-brown deep clay.

**Canopy Height:** 12-15 m

**DBH Canopy Trees:** 0.4 to 0.8 m      **DBH Sub-canopy Trees:** 0.1 to 0.2 m

**Tree Hollow Density:** low      **Fallen Log Density:** low

**Standing Dead Tree Density:** low      **Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Vegetation Community (Hunter Eco, 2019):**

- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter (PCT 1606).
- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter – DNG (PCT 1606).



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 3

**Site Description:** Sparse open box woodland with sparse shrub layer and ground cover on ridge top with Bulloak regeneration on slope and scattered Grey Box on lower slopes near creek lines and gullies.

**Habitat Condition:** Generally poor with evidence of grazing creating sparse ground cover with no regeneration of eucalypts.

**Site Disturbance Level and Type:** Heavily disturbed through cattle grazing.

**Connectivity:** Connected to the south to a much larger remnant offsite.

**Site Location:** -32.40436, 150.88962

**Patch Size:** 20 hectares

**Topography, Slope and Aspect:** Ridge top running north-west to south-east with a slope to a valley floor to the south-west.

**Soil Type:** Deep red-brown clay loam.

**Canopy Height:** 12 to 18 m

**DBH Canopy Trees:** 0.7 to 1.2 m

**DBH Sub-canopy Trees:** 0.2 to 0.5 m

**Tree Hollow Density:** moderate

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** moderate

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter (PCT 1606).
- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter – DNG (PCT 1606).
- Narrow-leaved Ironbark - Grey Box grassy woodland of the central and upper Hunter (PCT 1691).
- Narrow-leaved Ironbark - Grey Box grassy woodland of the central and upper Hunter – DNG (PCT 1691).
- Bull Oak grassy woodland of the central and upper Hunter (PCT 1692).



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 4

**Site Description:** Narrow band of mature White Box forming a very open woodland along a ridge top with very sparse shrub and ground cover.

**Habitat Condition:** Very open habitat mostly cleared and heavily grazed by cattle.

**Site Disturbance Level and Type:** Heavily disturbed through clearing and cattle grazing.

**Connectivity:** Connected to a large remnant to the north and to the rest of the narrow band to the south, which eventually reaches another large remnant in about 2 kilometres.

**Site Location:** -32.40188, 150.88299

**Patch Size:** 5 hectares

**Topography, Slope and Aspect:** Ridge top running north-south.

**Soil Type:** Shallow red-brown clay loam with some stones.

**Canopy Height:** 12-15 m

**DBH Canopy Trees:** 0.5 to 1.3 m

**DBH Sub-canopy Trees:** 0.2 to 0.3 m

**Tree Hollow Density:** moderate

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter (PCT 1606).
- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter – DNG (PCT 1606).
- Narrow-leaved Ironbark - Grey Box grassy woodland of the central and upper Hunter (PCT 1691).





## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 5

**Site Description:** Mature White Box/Grey Box open forest with areas of Bulloak regrowth and 2 dams (dry at the time of the survey). Shrub layer and ground cover very sparse.

**Habitat Condition:** Generally poor with evidence of heavy grazing.

**Site Disturbance Level and Type:** Very disturbed due to cattle grazing.

**Connectivity:** Partial tenuous connection with woodland remnants to the west and surrounded by open pasture on all other sides.

**Site Location:** -32.41405, 150.86669

**Patch Size:** 35 hectares

**Topography, Slope and Aspect:** Gentle slope to the north with a drainage line running to the north through the middle of the slope.

**Soil Type:** Deep red-brown clay loam.

**Canopy Height:** 15 to 18 m

**DBH Canopy Trees:** 0.4 to 1.3 m

**DBH Sub-canopy Trees:** 0.2 to 0.4 m

**Tree Hollow Density:** moderate

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Grey Box – Slaty Box shrub – grass woodland on sandstone slopes of the upper Hunter and Sydney Basin (PCT 1655).
- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter (PCT 1606).
- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter – DNG (PCT 1606).
- Narrow-leaved Ironbark - Grey Box grassy woodland of the central and upper Hunter (PCT 1691).
- Dam.



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 6

**Site Description:** Open White Box/Grey Box/Yellow Box forest with scattered patches of regenerating Bulloak and Cooba with sparse shrub layer and groundcover. Contains 2 dams with water at time of survey.

**Habitat Condition:** Poor due heavy grazing with much bare earth around dams.

**Site Disturbance Level and Type:** Heavily disturbed due to cattle grazing with little eucalypt regeneration.

**Connectivity:** Some weak connectivity to smaller remnants to the east and to the west.

**Site Location:** -32.41378, 150.84975

**Patch Size:** 80 hectares

**Topography, Slope and Aspect:** Low area with gentle slope to the north.

**Soil Type:** Deep red-brown clay loam.

**Canopy Height:** 12 to 15 m

**DBH Canopy Trees:** 0.3 to 0.7 m

**DBH Sub-canopy Trees:** 0.1 to 0.3 m

**Tree Hollow Density:** moderate

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** moderate

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant grass cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Fuzzy Box woodland on alluvial brown loam soils mainly in the NSW South Western Slopes Bioregion (PCT 201).
- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter (PCT 1606).
- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter – DNG (PCT 1606).
- Narrow-leaved Ironbark - Grey Box grassy woodland of the central and upper Hunter (PCT 1691).
- Yellow Box - Rough-barked Apple grassy woodland of the upper Hunter and Liverpool Plains (PCT 1693).
- Dam.



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 7

**Site Description:** Open White Box/Grey Box forest with patches of Bulloak and sparse shrub layer and mainly leaf litter as groundcover.

**Habitat Condition:** Evidence of grazing pressure and in generally poor condition with little eucalypt regeneration.

**Site Disturbance Level and Type:** Moderately disturbed with cattle grazing.

**Connectivity:** Continuous with very open woodland to the south-east and to a denser woodland patch to the north-east.

**Site Location:** -32.41698, 150.83850

**Patch Size:** 25 hectares

**Topography, Slope and Aspect:** Low ridge top that slopes gently to the west along 2 gullies.

**Soil Type:** Deep red-brown clay loam.

**Canopy Height:** 10 to 15 m

**DBH Canopy Trees:** 0.4 to 0.7 m

**DBH Sub-canopy Trees:** 0.2 to 0.3 m

**Tree Hollow Density:** low

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Grey Box – Slaty Box shrub – grass woodland on sandstone slopes of the upper Hunter and Sydney Basin (PCT 1655).
- Grey Box – Slaty Box shrub – grass woodland on sandstone slopes of the upper Hunter and Sydney Basin - DNG (PCT 1655).
- White Box -Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter – DNG (PCT 1606).
- Narrow-leaved Ironbark - Grey Box grassy woodland of the central and upper Hunter (PCT 1691).
- Bull Oak grassy woodland of the central Hunter Valley (PCT 1692).



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 8

**Site Description:** Dense narrow corridor of Swamp Oak regrowth along a creek-line.

**Habitat Condition:** Evidence of recent grazing and with thin cover of grass and much bare ground, condition poor.

**Site Disturbance Level and Type:** Recent grazing and mostly cleared around patch.

**Connectivity:** Connectivity poor with small isolated patches of regrowth along creek-line and with cleared grazing land around patch.

**Site Location:** -32.41063, 150.82563

**Patch Size:** 3 hectares

**Topography, Slope and Aspect:** Low area with gentle slopes along creek-line which drains towards the west.

**Soil Type:** Deep red-brown clay loam.

**Canopy Height:** 8-10 m

**DBH Canopy Trees:** 0.1 to 0.3 m

**DBH Sub-canopy Trees:** 0.03 to 0.1 m

**Tree Hollow Density:** nil

**Fallen Log Density:** low

**Standing Dead Tree Density:** nil

**Mistletoe Density:** nil

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Swamp Oak - Weeping Grass grassy riparian forest of the Hunter Valley (PCT 1731).
- Grey Box – Slaty Box shrub – grass woodland on sandstone slopes of the upper Hunter and Sydney Basin - DNG (PCT 1655).



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 9

**Site Description:** Areas along roads at the intersection have been planted with trees in three rows within fenced areas about 20 m wide along roadsides. The area of planting runs for about 100 to 200 m along three sections and for about 1 kilometre along the other road edge.

**Habitat Condition:** Trees are tall and thin because of close planting but there is some recruitment of local native shrub and grass species. Habitat value is low due to narrow area of planting.

**Site Disturbance Level and Type:** Very low level of disturbance after original planting as good fencing has excluded livestock.

**Connectivity:** Poor connectivity to the north with some patchy regenerating woodland along creek-line.

**Site Location:** -32.44012, 150.82915

**Patch Size:** 3 hectares

**Topography, Slope and Aspect:** Gently slopes from west to east along main planting area and to road junction for other 3 areas.

**Soil Type:** Red-brown deep clay-loam.

**Canopy Height:** 10-15 m

**DBH Canopy Trees:** 0.3 to 0.7

**DBH Sub-canopy Trees:** 0.1 to 0.2

**Tree Hollow Density:** nil

**Fallen Log Density:** nil

**Standing Dead Tree Density:** nil

**Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Planted trees.



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 10

**Site Description:** Sparse open box woodland along a gully with patches of Bulloak with sparse shrub layer and poor cover of grasses and forbes. A dam (dry at the time of the survey) is also in this site.

**Habitat Condition:** Very poor open woodland with little regeneration of eucalypts.

**Site Disturbance Level and Type:** Highly disturbed by grazing cattle.

**Connectivity:** Some connectivity to the north of the site to areas of regeneration Bulloak and scattered eucalypts.

**Site Location:** -32.43750, 150.85237

**Patch Size:** 10 hectares

**Topography, Slope and Aspect:** Moderately steep slope to the south along a gully.

**Soil Type:** Red-brown clays.

**Canopy Height:** 8-12 m

**DBH Canopy Trees:** 0.4 to 0.8 m

**DBH Sub-canopy Trees:** 0.1 to 0.2 m

**Tree Hollow Density:** low

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter (PCT 1606).
- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter – DNG (PCT 1606).



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 11

**Site Description:** Mature stand of White Box in an open forest on a moderately steep slope with a sparse understorey and shrub layer and moderate grass cover. There is a large dam with water at the bottom of the slope.

**Habitat Condition:** Fair condition although there is little regeneration of eucalypts.

**Site Disturbance Level and Type:** Moderate with cattle grazing.

**Connectivity:** Connects to very open woodland to the west and to a larger remnant to the east.

**Site Location:** -32.42325, 150.87886

**Patch Size:** 10 hectares

**Topography, Slope and Aspect:** Moderately steep slope to the west from a ridge top.

**Soil Type:** Red-brown clay loam.

**Canopy Height:** 12 to 16 m

**DBH Canopy Trees:** 0.7 to 1.3 m

**DBH Sub-canopy Trees:** 0.5 to 0.7 m

**Tree Hollow Density:** moderate

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter (PCT 1606).
- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter – DNG (PCT 1606).
- Dam.



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 12

**Site Description:** A long remnant of riparian woodland in mostly cleared area of pastureland with scattered Grey Box and Bulloak with some dense areas of Bulloak and Cooba regeneration.

**Habitat Condition:** Evidence of heavy grazing and very dry conditions have left site in poor condition.

**Site Disturbance Level and Type:** Highly disturbed due to cattle grazing.

**Connectivity:** Connected to larger patch of vegetation to the north at the head of the creek valley.

**Site Location:** -32.459599, 150.856216

**Patch Size:** 15 hectares

**Topography, Slope and Aspect:** Creek valley sloping to the south.

**Soil Type:** Shallow, stoney brown clay loam.

**Canopy Height:** 10-18 m

**DBH Canopy Trees:** 0.3 to 1.3 m

**DBH Sub-canopy Trees:** 0.2 to 0.4 m

**Tree Hollow Density:** low

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter (PCT 1606).
- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter – DNG (PCT 1606).
- Yellow Box - Rough-barked Apple grassy woodland of the upper Hunter and Liverpool Plains (PCT 1693).





## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 13

**Site Description:** A small remnant of riparian woodland along Saddlers Creek in mostly cleared area of pastureland with scattered Fuzzy Box and Swamp Oak with some dense areas of Swamp Oak regeneration.

**Habitat Condition:** Evidence of heavy grazing and very dry conditions have left site in poor condition.

**Site Disturbance Level and Type:** Highly disturbed due to cattle grazing.

**Connectivity:** Poorly connected via sparse cover of riparian vegetation along creeks passing through the site.

**Site Location:** -32.414284, 150.821075

**Patch Size:** 10 hectares

**Topography, Slope and Aspect:** Mainly level area along the banks of the creek.

**Soil Type:** Shallow, stoney brown clay loam.

**Canopy Height:** 10-12 m

**DBH Canopy Trees:** 0.3 to 0.6 m

**DBH Sub-canopy Trees:** 0.1 to 0.3 m

**Tree Hollow Density:** low

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** Very low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Fuzzy Box woodland on alluvial brown loam soils mainly in the NSW South Western Slopes Bioregion – DNG (PCT 201).
- Swamp Oak - Weeping Grass grassy riparian forest of the Hunter Valley (PCT 1731).



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 14

**Site Description:** Mostly cleared area of pastureland with scattered remnant of eucalypts and Bulloak with some dense areas of Bulloak regeneration.

**Habitat Condition:** Evidence of heavy grazing and very dry conditions have left site in poor condition.

**Site Disturbance Level and Type:** Highly disturbed due to cattle grazing.

**Connectivity:** Poorly connected via sparse cover of riparian vegetation along creeks passing through the site.

**Site Location:** -32.429379, 150.824432

**Patch Size:** 50 hectares

**Topography, Slope and Aspect:** Gently undulating with a slight slope to the north along the drainage lines passing through the site.

**Soil Type:** Shallow, stoney brown clay loam.

**Canopy Height:** 15-20 m

**DBH Canopy Trees:** 0.4 to 1.2 m

**DBH Sub-canopy Trees:** 0.2 to 0.4 m

**Tree Hollow Density:** High density of hollows in each tree but only moderate density over the whole site due to tree spacing.

**Fallen Log Density:** Moderate

**Standing Dead Tree Density:** Moderate

**Mistletoe Density:** Very low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Fuzzy Box woodland on alluvial brown loam soils mainly in the NSW South Western Slopes Bioregion (PCT 201).
- Fuzzy Box woodland on alluvial brown loam soils mainly in the NSW South Western Slopes Bioregion – DNG (PCT 201).
- Grey Box – Slaty Box shrub – grass woodland on sandstone slopes of the upper Hunter and Sydney Basin (PCT 1655)
- Grey Box – Slaty Box shrub – grass woodland on sandstone slopes of the upper Hunter and Sydney Basin – DNG (PCT 1655).



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 15

**Site Description:** Open eucalypt forest with a patchy shrub layer and some dense areas of sapling regrowth with a mixture of sparse grass cover and leaf litter for ground cover.

**Habitat Condition:** No evidence of recent grazing and much regeneration. Habitat in fair condition despite very dry conditions.

**Site Disturbance Level and Type:** Partial clearing for grazing in the past but little recent disturbance.

**Connectivity:** Connected to a much larger area of woodland to the south, broken only by a road and conveyer belt.

**Site Location:** -32.380155, 150.931008

**Patch Size:** 70 hectares

**Topography, Slope and Aspect:** Ridge top at western edge and slope to the east with several small gullies passing through site and joining at eastern edge.

**Soil Type:** Shallow, red-brown clay loam with some exposed rock.

**Canopy Height:** 15-20 m

**DBH Canopy Trees:** 0.2 to 0.9 m

**DBH Sub-canopy Trees:** 0.1 to 0.2 m

**Tree Hollow Density:** low

**Fallen Log Density:** moderate

**Standing Dead Tree Density:** moderate

**Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Narrow-leaved Ironbark - Grey Box - Spotted Gum shrub - grass woodland of the central and lower Hunter (PCT 1604).
- White Box -Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter - DNG (PCT 1606).



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 16

**Site Description:** Tall dense regrowth Spotted Gum forest with some dry open Box Gum woodland and small patches of shrubby cover.

**Habitat Condition:** Regrowth areas are structurally very simple with only tall trees and leaf litter. Patches of Box Gum woodland structurally more complex and in fair condition.

**Site Disturbance Level and Type:** Much of the site has been previously cleared for grazing but is now covered with regrowth forest with no recent evidence of disturbance.

**Connectivity:** Connected to the north-west through to the north-east to a very large area of woodland.

**Site Location:** -32.335047, 150.935035

**Patch Size:** 80 hectares

**Topography, Slope and Aspect:** Gently slopes towards large dam at the centre of the site.

**Soil Type:** Red-brown shallow stoney clay-loam.

**Canopy Height:** 15-20 m

**DBH Canopy Trees:** 0.2 to 0.6 m

**DBH Sub-canopy Trees:** 0.1 to 0.2 m

**Tree Hollow Density:** low

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Narrow-leaved Ironbark - Grey Box - Spotted Gum shrub - grass woodland of the central and lower Hunter (PCT 1604).
- Forest Red Gum grassy open forest on floodplains of the lower Hunter (PCT 1598).
- Dam.
- Infrastructure and old workings.



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 17

**Site Description:** Open eucalypt forest regeneration with a few old growth trees, with a patchy and mostly sparse shrub layer and some dense areas of sapling regrowth with a mixture of sparse grass cover and leaf litter for ground cover.

**Habitat Condition:** No evidence of recent grazing and much regeneration. Habitat in poor condition due to sparse cover and very dry conditions.

**Site Disturbance Level and Type:** Previously mostly cleared but little recent disturbance.

**Connectivity:** Connected to a much larger area of remnant woodland to the north and the west. Mine infrastructure to the east and south.

**Site Location:** -32.336447, 150.924972

**Patch Size:** 10 hectares

**Topography, Slope and Aspect:** Gentle slopes to the west on undulating land with some small flood water courses running from the southern edge to the north-west.

**Soil Type:** Shallow, stoney red-brown clay loam.

**Canopy Height:** 12-17 m

**DBH Canopy Trees:** 0.2 to 1.3 m

**DBH Sub-canopy Trees:** 0.1 to 0.2 m

**Tree Hollow Density:** low

**Fallen Log Density:** low

**Standing Dead Tree Density:** low

**Mistletoe Density:** low

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Narrow-leaved Ironbark - Grey Box - Spotted Gum shrub - grass woodland of the central and lower Hunter (PCT 1604).
- Infrastructure and old workings.



## MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

---

**Site Number:** 18

**Site Description:** Mixture of riparian remnant old growth trees and Swamp Oak and Bulloak regeneration along a creek running east to west through mostly cleared grazing area that has only been released from grazing for a few months.

**Habitat Condition:** Evidence of recent grazing and much regeneration. Habitat in poor condition due to sparse cover and very dry conditions.

**Site Disturbance Level and Type:** Previously mostly cleared with evidence of grazing pressure under very dry conditions.

**Connectivity:** Poorly connected except along watercourse where the riparian zone varies in thickness and quality and with some open sections with little woody cover.

**Site Location:** -32.405822, 150.847956

**Patch Size:** 3 hectares

**Topography, Slope and Aspect:** Gentle slopes to the west on mostly flat land with watercourse running to the west through the site.

**Soil Type:** Deep red-brown clay loam.

**Canopy Height:** 8-18 m

**DBH Canopy Trees:** 0.1 to 0.9 m

**DBH Sub-canopy Trees:** 0.1 to 0.2 m

**Tree Hollow Density:** low

**Fallen Log Density:** low

**Standing Dead Tree Density:** moderate

**Mistletoe Density:** nil

**Dominant Canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant canopy species present.

**Dominant Sub-canopy Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant sub-canopy species present.

**Dominant Shrub Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant shrub species present.

**Dominant Ground Cover Species:** Refer to the *Maxwell Project Baseline Flora Report* (Hunter Eco, 2019) for details on the dominant ground cover species present.

**Vegetation Community (Hunter Eco, 2019):**

- Swamp Oak - Weeping Grass grassy riparian forest of the Hunter Valley (PCT 1731).
- Yellow Box - Rough-barked Apple grassy woodland of the upper Hunter and Liverpool Plains – DNG (PCT 1693).
- White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter – DNG (PCT 1606).



# Appendix C Threatened Biodiversity Data Collection Species PCT Associations Sydney Basin – Hunter IBRA Sub-region

**Table C-1  
Threatened Biodiversity Data Collection Species PCT Associations Sydney Basin – Hunter IBRA  
Sub-region**

Scientific Name	Common Name	PCT	Notes
<i>Litoria aurea</i>	Green and Golden Bell Frog	1598	Species not present.
<i>Litoria aurea</i>	Green and Golden Bell Frog	1604	
<i>Litoria aurea</i>	Green and Golden Bell Frog	1606	
<i>Litoria aurea</i>	Green and Golden Bell Frog	1691	
<i>Litoria aurea</i>	Green and Golden Bell Frog	1692	
<i>Litoria aurea</i>	Green and Golden Bell Frog	1731	
<i>Litoria brevipalmata</i>	Green-thighed Frog	1598	No potential habitat in the study area. Species not present.
<i>Litoria brevipalmata</i>	Green-thighed Frog	1604	
<i>Aprasia parapulchella</i>	Pink-tailed Legless Lizard	None	Recorded in PCT 1606. Potential habitat within rocky areas mapped in PCT 1606 and 1606 DNG.
<i>Delma impar</i>	Striped Legless Lizard	1604	Recorded in PCT 1655 and 1655 DNG, PCT 1692, PCT 1693 DNG, PCT 1691 DNG, PCT1606 and PCT1606 DNG so these are also considered habitat.
<i>Delma impar</i>	Striped Legless Lizard	1655	
<i>Delma impar</i>	Striped Legless Lizard	1691	
<i>Delma impar</i>	Striped Legless Lizard	1692	
<i>Delma impar</i>	Striped Legless Lizard	1693	
<i>Hoplocephalus bitorquatus</i>	Pale-headed Snake	1604	Species not present.
<i>Hoplocephalus bitorquatus</i>	Pale-headed Snake	1606	
<i>Hoplocephalus bitorquatus</i>	Pale-headed Snake	1655	
<i>Hoplocephalus bitorquatus</i>	Pale-headed Snake	1691	
<i>Hoplocephalus bitorquatus</i>	Pale-headed Snake	1692	
<i>Lophoictinia isura</i>	Square-tailed Kite	201	These PCTs are considered appropriate potential foraging habitat. No breeding habitat is present.
<i>Lophoictinia isura</i>	Square-tailed Kite	1604	
<i>Lophoictinia isura</i>	Square-tailed Kite	1606	
<i>Lophoictinia isura</i>	Square-tailed Kite	1607	
<i>Lophoictinia isura</i>	Square-tailed Kite	1655	
<i>Lophoictinia isura</i>	Square-tailed Kite	1691	
<i>Lophoictinia isura</i>	Square-tailed Kite	1692	
<i>Lophoictinia isura</i>	Square-tailed Kite	1693	
<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle	201	These PCTs are considered appropriate potential foraging habitat and PCT 1604 (recorded in). No breeding habitat is present.
<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle	1598	
<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle	1607	
<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle	1691	
<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle	1692	
<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle	1731	
<i>Circus assimilis</i>	Spotted Harrier	1731	PCT 1606 DNG and 1691 are also considered potential habitat.
<i>Hieraaetus morphnoides</i>	Little Eagle	116	These PCTs are considered appropriate potential foraging habitat. No breeding habitat is present.
<i>Hieraaetus morphnoides</i>	Little Eagle	201	
<i>Hieraaetus morphnoides</i>	Little Eagle	1604	
<i>Hieraaetus morphnoides</i>	Little Eagle	1606	
<i>Hieraaetus morphnoides</i>	Little Eagle	1655	
<i>Hieraaetus morphnoides</i>	Little Eagle	1691	
<i>Hieraaetus morphnoides</i>	Little Eagle	1692	
<i>Hieraaetus morphnoides</i>	Little Eagle	1731	
<i>Burhinus grallarius</i>	Bush Stone-curlew	201	Species not present.
<i>Burhinus grallarius</i>	Bush Stone-curlew	1604	
<i>Burhinus grallarius</i>	Bush Stone-curlew	1606	
<i>Burhinus grallarius</i>	Bush Stone-curlew	1655	
<i>Burhinus grallarius</i>	Bush Stone-curlew	1691	
<i>Burhinus grallarius</i>	Bush Stone-curlew	1692	

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Scientific Name	Common Name	PCT	Notes
<i>Calyptorhynchus lathamii</i>	Glossy Black-Cockatoo	201	These PCTs are considered appropriate foraging habitat. No breeding habitat is present.
<i>Calyptorhynchus lathamii</i>	Glossy Black-Cockatoo	1604	
<i>Calyptorhynchus lathamii</i>	Glossy Black-Cockatoo	1606	
<i>Calyptorhynchus lathamii</i>	Glossy Black-Cockatoo	1655	
<i>Calyptorhynchus lathamii</i>	Glossy Black-Cockatoo	1691	
<i>Calyptorhynchus lathamii</i>	Glossy Black-Cockatoo	1692	
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo	201	These PCTs are considered appropriate potential foraging habitat. No breeding habitat is present.
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo	1604	
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo	1606	
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo	1655	
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo	1691	
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo	1692	
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo	1731	
<i>Glossopsitta pusilla</i>	Little Lorikeet	201	These PCTs are considered appropriate.
<i>Glossopsitta pusilla</i>	Little Lorikeet	1598	
<i>Glossopsitta pusilla</i>	Little Lorikeet	1604	
<i>Glossopsitta pusilla</i>	Little Lorikeet	1606	
<i>Glossopsitta pusilla</i>	Little Lorikeet	1607	
<i>Glossopsitta pusilla</i>	Little Lorikeet	1655	
<i>Glossopsitta pusilla</i>	Little Lorikeet	1691	
<i>Glossopsitta pusilla</i>	Little Lorikeet	1692	
<i>Glossopsitta pusilla</i>	Little Lorikeet	1693	
<i>Neophema pulchella</i>	Turquoise Parrot	201	These PCTs are considered appropriate.
<i>Neophema pulchella</i>	Turquoise Parrot	1604	
<i>Neophema pulchella</i>	Turquoise Parrot	1606	
<i>Neophema pulchella</i>	Turquoise Parrot	1607	
<i>Neophema pulchella</i>	Turquoise Parrot	1655	
<i>Neophema pulchella</i>	Turquoise Parrot	1691	
<i>Neophema pulchella</i>	Turquoise Parrot	1692	
<i>Neophema pulchella</i>	Turquoise Parrot	1693	
<i>Lathamus discolor</i>	Swift Parrot	201	
<i>Lathamus discolor</i>	Swift Parrot	1604	
<i>Lathamus discolor</i>	Swift Parrot	1606	
<i>Lathamus discolor</i>	Swift Parrot	1655	
<i>Lathamus discolor</i>	Swift Parrot	1691	
<i>Lathamus discolor</i>	Swift Parrot	1692	
<i>Tyto longimembris</i>	Eastern Grass Owl	1731	This species is considered unlikely to use habitat in the study area. Species not present.
<i>Tyto novaehollandiae</i>	Masked Owl	201	These PCTs are considered appropriate potential foraging habitat. No breeding habitat is present.
<i>Tyto novaehollandiae</i>	Masked Owl	1604	
<i>Tyto novaehollandiae</i>	Masked Owl	1606	
<i>Tyto novaehollandiae</i>	Masked Owl	1655	
<i>Tyto novaehollandiae</i>	Masked Owl	1691	
<i>Tyto novaehollandiae</i>	Masked Owl	1692	
<i>Ninox strenua</i>	Powerful Owl	1604	These PCTs are considered appropriate potential foraging habitat. No breeding habitat is present.
<i>Ninox strenua</i>	Powerful Owl	1606	
<i>Ninox strenua</i>	Powerful Owl	1655	
<i>Ninox strenua</i>	Powerful Owl	1691	
<i>Ninox strenua</i>	Powerful Owl	1692	
<i>Ninox connivens</i>	Barking Owl	201	
<i>Ninox connivens</i>	Barking Owl	1598	
<i>Ninox connivens</i>	Barking Owl	1604	
<i>Ninox connivens</i>	Barking Owl	1606	
<i>Ninox connivens</i>	Barking Owl	1607	
<i>Ninox connivens</i>	Barking Owl	1655	
<i>Ninox connivens</i>	Barking Owl	1691	
<i>Ninox connivens</i>	Barking Owl	1692	
<i>Ninox connivens</i>	Barking Owl	1693	
<i>Ninox connivens</i>	Barking Owl	1731	
<i>Climacteris picumnus victoriae</i>	Brown Treecreeper (eastern subspecies)	201	These PCTs are considered appropriate.
<i>Climacteris picumnus victoriae</i>	Brown Treecreeper (eastern subspecies)	1598	



MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Scientific Name	Common Name	PCT	Notes
<i>Climacteris picumnus victoriae</i>	Brown Treecreeper (eastern subspecies)	1604	
<i>Climacteris picumnus victoriae</i>	Brown Treecreeper (eastern subspecies)	1606	
<i>Climacteris picumnus victoriae</i>	Brown Treecreeper (eastern subspecies)	1607	
<i>Climacteris picumnus victoriae</i>	Brown Treecreeper (eastern subspecies)	1655	
<i>Climacteris picumnus victoriae</i>	Brown Treecreeper (eastern subspecies)	1691	
<i>Climacteris picumnus victoriae</i>	Brown Treecreeper (eastern subspecies)	1693	
<i>Chthonicola sagittata</i>	Speckled Warbler	201	These PCTs are considered appropriate.
<i>Chthonicola sagittata</i>	Speckled Warbler	1598	
<i>Chthonicola sagittata</i>	Speckled Warbler	1604	
<i>Chthonicola sagittata</i>	Speckled Warbler	1606	
<i>Chthonicola sagittata</i>	Speckled Warbler	1607	
<i>Chthonicola sagittata</i>	Speckled Warbler	1655	
<i>Chthonicola sagittata</i>	Speckled Warbler	1691	
<i>Chthonicola sagittata</i>	Speckled Warbler	1692	
<i>Chthonicola sagittata</i>	Speckled Warbler	1693	
<i>Melithreptus gularis gularis</i>	Black-chinned Honeyeater (eastern subspecies)	201	These PCTs are considered appropriate.
<i>Melithreptus gularis gularis</i>	Black-chinned Honeyeater (eastern subspecies)	1604	
<i>Melithreptus gularis gularis</i>	Black-chinned Honeyeater (eastern subspecies)	1606	
<i>Melithreptus gularis gularis</i>	Black-chinned Honeyeater (eastern subspecies)	1655	
<i>Melithreptus gularis gularis</i>	Black-chinned Honeyeater (eastern subspecies)	1691	
<i>Melithreptus gularis gularis</i>	Black-chinned Honeyeater (eastern subspecies)	1692	
<i>Anthochaera phrygia</i>	Regent Honeyeater	201	These PCTs are considered appropriate potential foraging habitat. No important habitat (as defined by OEH) is present.
<i>Anthochaera phrygia</i>	Regent Honeyeater	1604	
<i>Anthochaera phrygia</i>	Regent Honeyeater	1606	
<i>Anthochaera phrygia</i>	Regent Honeyeater	1607	
<i>Anthochaera phrygia</i>	Regent Honeyeater	1655	
<i>Anthochaera phrygia</i>	Regent Honeyeater	1691	
<i>Anthochaera phrygia</i>	Regent Honeyeater	1693	
<i>Grantiella picta</i>	Painted Honeyeater	116	These PCTs are considered appropriate.
<i>Grantiella picta</i>	Painted Honeyeater	201	
<i>Grantiella picta</i>	Painted Honeyeater	1604	
<i>Grantiella picta</i>	Painted Honeyeater	1606	
<i>Grantiella picta</i>	Painted Honeyeater	1607	
<i>Grantiella picta</i>	Painted Honeyeater	1655	
<i>Grantiella picta</i>	Painted Honeyeater	1691	
<i>Grantiella picta</i>	Painted Honeyeater	1692	
<i>Melanodryas cucullata cucullata</i>	Hooded Robin (south-eastern form)	201	These PCTs are considered appropriate.
<i>Melanodryas cucullata cucullata</i>	Hooded Robin (south-eastern form)	1598	
<i>Melanodryas cucullata cucullata</i>	Hooded Robin (south-eastern form)	1604	
<i>Melanodryas cucullata cucullata</i>	Hooded Robin (south-eastern form)	1606	
<i>Melanodryas cucullata cucullata</i>	Hooded Robin (south-eastern form)	1607	
<i>Melanodryas cucullata cucullata</i>	Hooded Robin (south-eastern form)	1655	
<i>Melanodryas cucullata cucullata</i>	Hooded Robin (south-eastern form)	1691	
<i>Melanodryas cucullata cucullata</i>	Hooded Robin (south-eastern form)	1692	
<i>Melanodryas cucullata cucullata</i>	Hooded Robin (south-eastern form)	1693	
<i>Petroica phoenicea</i>	Flame Robin	116	These PCTs are considered appropriate.
<i>Petroica phoenicea</i>	Flame Robin	1604	
<i>Petroica phoenicea</i>	Flame Robin	1606	
<i>Petroica phoenicea</i>	Flame Robin	1607	
<i>Petroica phoenicea</i>	Flame Robin	1655	
<i>Petroica phoenicea</i>	Flame Robin	1691	
<i>Petroica phoenicea</i>	Flame Robin	1692	
<i>Petroica boodang</i>	Scarlet Robin	116	These PCTs are considered appropriate.
<i>Petroica boodang</i>	Scarlet Robin	201	
<i>Petroica boodang</i>	Scarlet Robin	1598	

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Scientific Name	Common Name	PCT	Notes
<i>Petroica boodang</i>	Scarlet Robin	1604	
<i>Petroica boodang</i>	Scarlet Robin	1606	
<i>Petroica boodang</i>	Scarlet Robin	1607	
<i>Petroica boodang</i>	Scarlet Robin	1655	
<i>Petroica boodang</i>	Scarlet Robin	1691	
<i>Petroica boodang</i>	Scarlet Robin	1692	
<i>Petroica boodang</i>	Scarlet Robin	1693	
<i>Pomatostomus temporalis temporalis</i>	Grey-crowned Babbler (eastern subspecies)	201	PCT 1693 and PCT 1731 are also considered potential habitat.
<i>Pomatostomus temporalis temporalis</i>	Grey-crowned Babbler (eastern subspecies)	1604	
<i>Pomatostomus temporalis temporalis</i>	Grey-crowned Babbler (eastern subspecies)	1606	
<i>Pomatostomus temporalis temporalis</i>	Grey-crowned Babbler (eastern subspecies)	1655	
<i>Pomatostomus temporalis temporalis</i>	Grey-crowned Babbler (eastern subspecies)	1691	
<i>Pomatostomus temporalis temporalis</i>	Grey-crowned Babbler (eastern subspecies)	1692	
<i>Daphoenositta chrysoptera</i>	Varied Sittella	116	PCT 1692 and PCT 1598 are also considered potential habitat.
<i>Daphoenositta chrysoptera</i>	Varied Sittella	201	
<i>Daphoenositta chrysoptera</i>	Varied Sittella	1598	
<i>Daphoenositta chrysoptera</i>	Varied Sittella	1604	
<i>Daphoenositta chrysoptera</i>	Varied Sittella	1606	
<i>Daphoenositta chrysoptera</i>	Varied Sittella	1607	
<i>Daphoenositta chrysoptera</i>	Varied Sittella	1655	
<i>Daphoenositta chrysoptera</i>	Varied Sittella	1691	
<i>Daphoenositta chrysoptera</i>	Varied Sittella	1693	
<i>Daphoenositta chrysoptera</i>	Varied Sittella	1731	
<i>Artamus cyanopterus cyanopterus</i>	Dusky Woodswallow	201	PCT 1604 and PCT 1606 DNG are also considered potential habitat.
<i>Stagonopleura guttata</i>	Diamond Firetail	201	PCT 1691 is also considered potential habitat.
<i>Stagonopleura guttata</i>	Diamond Firetail	1604	
<i>Stagonopleura guttata</i>	Diamond Firetail	1606	
<i>Stagonopleura guttata</i>	Diamond Firetail	1655	
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	201	These PCTs are considered appropriate (in woodland and DNG form).
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	1598	
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	1604	
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	1606	
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	1607	
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	1655	
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	1691	
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	1692	
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	1693	
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	1731	
<i>Phascogale tapoatafa</i>	Brush-tailed Phascogale	201	These PCTs are considered appropriate.
<i>Phascogale tapoatafa</i>	Brush-tailed Phascogale	1604	
<i>Phascogale tapoatafa</i>	Brush-tailed Phascogale	1606	
<i>Phascogale tapoatafa</i>	Brush-tailed Phascogale	1691	
<i>Phascogale tapoatafa</i>	Brush-tailed Phascogale	1692	
<i>Phascogale tapoatafa</i>	Brush-tailed Phascogale	1731	
<i>Planigale maculata</i>	Common Planigale	1604	This species is considered unlikely to use habitat in the study area. Species not present.
<i>Planigale maculata</i>	Common Planigale	1606	
<i>Planigale maculata</i>	Common Planigale	1655	
<i>Planigale maculata</i>	Common Planigale	1691	
<i>Planigale maculata</i>	Common Planigale	1692	
<i>Phascolarctos cinereus</i>	Koala	201	These PCTs are considered appropriate.
<i>Phascolarctos cinereus</i>	Koala	1598	
<i>Phascolarctos cinereus</i>	Koala	1604	
<i>Phascolarctos cinereus</i>	Koala	1606	
<i>Phascolarctos cinereus</i>	Koala	1607	
<i>Phascolarctos cinereus</i>	Koala	1655	
<i>Phascolarctos cinereus</i>	Koala	1693	
<i>Cercartetus nanus</i>	Eastern Pygmy-possum	1604	This species is considered unlikely to use habitat in the study area. Species not present.
<i>Cercartetus nanus</i>	Eastern Pygmy-possum	1606	
<i>Cercartetus nanus</i>	Eastern Pygmy-possum	1655	

MAXWELL PROJECT BASELINE FAUNA SURVEY REPORT

Scientific Name	Common Name	PCT	Notes
<i>Cercartetus nanus</i>	Eastern Pygmy-possum	1691	
<i>Cercartetus nanus</i>	Eastern Pygmy-possum	1692	
<i>Petaurus australis</i>	Yellow-bellied Glider	1604	This species is considered unlikely to use habitat in the study area. Species not present.
<i>Petaurus australis</i>	Yellow-bellied Glider	1606	
<i>Petaurus norfolcensis</i>	Squirrel Glider	201	PCT 1598 and 1604 are also considered potential habitat.
<i>Petaurus norfolcensis</i>	Squirrel Glider	1606	
<i>Petaurus norfolcensis</i>	Squirrel Glider	1655	
<i>Petrogale penicillata</i>	Brush-tailed Rock-wallaby	201	This species is considered unlikely to use habitat in the study area. Species not present.
<i>Petrogale penicillata</i>	Brush-tailed Rock-wallaby	1604	
<i>Petrogale penicillata</i>	Brush-tailed Rock-wallaby	1655	
<i>Petrogale penicillata</i>	Brush-tailed Rock-wallaby	1691	
<i>Petrogale penicillata</i>	Brush-tailed Rock-wallaby	1692	
<i>Pteropus poliocephalus</i>	Grey-headed Flying-fox	1604	These PCTs are considered appropriate.
<i>Pteropus poliocephalus</i>	Grey-headed Flying-fox	1606	
<i>Pteropus poliocephalus</i>	Grey-headed Flying-fox	1655	
<i>Pteropus poliocephalus</i>	Grey-headed Flying-fox	1691	
<i>Pteropus poliocephalus</i>	Grey-headed Flying-fox	1692	
<i>Saccolaimus flaviventris</i>	Yellow-bellied Sheath-tail-bat	201	These PCTs are considered appropriate.
<i>Saccolaimus flaviventris</i>	Yellow-bellied Sheath-tail-bat	1604	
<i>Saccolaimus flaviventris</i>	Yellow-bellied Sheath-tail-bat	1606	
<i>Saccolaimus flaviventris</i>	Yellow-bellied Sheath-tail-bat	1655	
<i>Saccolaimus flaviventris</i>	Yellow-bellied Sheath-tail-bat	1691	
<i>Saccolaimus flaviventris</i>	Yellow-bellied Sheath-tail-bat	1692	
<i>Mormopterus norfolkensis</i>	Eastern Freetail-bat	1604	PCT 1598, 1606 and 1693 are also considered potential habitat.
<i>Mormopterus norfolkensis</i>	Eastern Freetail-bat	1691	
<i>Mormopterus norfolkensis</i>	Eastern Freetail-bat	1692	
<i>Miniopterus australis</i>	Little Bentwing-bat	1604	PCT 1598 and 1606 are also considered potential habitat.
<i>Miniopterus schreibersii oceanensis</i>	Eastern Bentwing-bat	201	PCT 1598, 1607 and 1693 are also considered potential habitat.
<i>Miniopterus schreibersii oceanensis</i>	Eastern Bentwing-bat	1604	
<i>Miniopterus schreibersii oceanensis</i>	Eastern Bentwing-bat	1606	
<i>Miniopterus schreibersii oceanensis</i>	Eastern Bentwing-bat	1655	
<i>Miniopterus schreibersii oceanensis</i>	Eastern Bentwing-bat	1691	
<i>Miniopterus schreibersii oceanensis</i>	Eastern Bentwing-bat	1692	
<i>Nyctophilus corbeni</i>	Corben's Long-eared Bat	201	PCT 1691 is also considered potential habitat.
<i>Nyctophilus corbeni</i>	Corben's Long-eared Bat	1606	
<i>Nyctophilus corbeni</i>	Corben's Long-eared Bat	1655	
<i>Chalinolobus dwyeri</i>	Large-eared Pied Bat	201	Adopted for use of the 'Species Credit' Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method (OEH, 2018).
<i>Chalinolobus dwyeri</i>	Large-eared Pied Bat	1604	
<i>Chalinolobus dwyeri</i>	Large-eared Pied Bat	1606	
<i>Chalinolobus dwyeri</i>	Large-eared Pied Bat	1655	
<i>Chalinolobus dwyeri</i>	Large-eared Pied Bat	1691	
<i>Chalinolobus dwyeri</i>	Large-eared Pied Bat	1692	
<i>Falsistrellus tasmaniensis</i>	Eastern False Pipistrelle	1604	PCT 1598 is also considered potential habitat.
<i>Falsistrellus tasmaniensis</i>	Eastern False Pipistrelle	1606	
<i>Falsistrellus tasmaniensis</i>	Eastern False Pipistrelle	1655	
<i>Falsistrellus tasmaniensis</i>	Eastern False Pipistrelle	1691	
<i>Falsistrellus tasmaniensis</i>	Eastern False Pipistrelle	1692	
<i>Myotis macropus</i>	Southern Myotis	1604	Adopted for use of the 'Species Credit' Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method (OEH, 2018).
<i>Myotis macropus</i>	Southern Myotis	1691	
<i>Myotis macropus</i>	Southern Myotis	1692	
<i>Scoteanax rueppellii</i>	Greater Broad-nosed Bat	1604	PCT 1598 is also considered potential habitat.
<i>Scoteanax rueppellii</i>	Greater Broad-nosed Bat	1606	
<i>Scoteanax rueppellii</i>	Greater Broad-nosed Bat	1655	
<i>Scoteanax rueppellii</i>	Greater Broad-nosed Bat	1691	
<i>Scoteanax rueppellii</i>	Greater Broad-nosed Bat	1692	
<i>Vespadelus troughtoni</i>	Eastern Cave Bat	1604	Adopted for use of the 'Species Credit' Threatened Bats and their Habitats: NSW Survey Guide for the Biodiversity Assessment Method (OEH, 2018).
<i>Vespadelus troughtoni</i>	Eastern Cave Bat	1606	
<i>Vespadelus troughtoni</i>	Eastern Cave Bat	1655	

ATTACHMENT B  
VEGETATION INTEGRITY DATA

**Table B-1 Proposed MEA Extensions**

plot	pct	area	patchsize	conditionclass	zone	easting	northing	bearing	compTree	compShrub	compGrass	compForbs	compFerns	compOther	strucTree	strucShrub	strucGrass	strucForbs	strucFerns	strucOther	funLargeTrees	funHollowtrees	funLitterCover	funLenFallenLogs	funTreeStem5to10	funTreeStem10to20	funTreeStem20to30	funTreeStem30to50	funTreeStem50to80	funTreeRegen	funHighThreatExotic
181121P5	1606	3.8	101	DNG	56	299224	6412435	350	0	2	4	15	0	3	0	3.1	55.1	1.8	0	0.3	0	0	8	0	0	0	0	0	0	0	2.1
210628P2	1606	3.8	101	DNG	56	298926	6412370	228	1	2	10	10	1	2	25	0.2	70.6	15.8	0.1	0.2	0	0	11	0	0	1	0	0	0	1	5.1
210628P3	1606	3.8	101	DNG	56	299203	6412041	131	1	1	7	4	1	3	0.1	0.1	75.2	5.3	0.1	0.3	0	0	56	0	0	0	0	0	0	1	6.1
180912P2	1692	0.5	101	Moderate	56	299017	6411944	110	2	3	3	11	1	1	93	0.3	0.3	1.1	0.1	0.1	0	0	52	0.0	1	1	0	1	0	1	0
210628P4	1692	0.5	101	Moderate	56	298993	6412060	204	2	5	14	9	1	4	90.1	2.5	6.3	1.9	0.1	0.4	0	0	52	3	1	1	1	0	0	1	0.2

**Table B-2 Proposed Water Treatment Facility Location**

plot	pct	area	patchsize	conditionclass	zone	easting	northing	bearing	compTree	compShrub	compGrass	compForbs	compFerns	compOther	strucTree	strucShrub	strucGrass	strucForbs	strucFerns	strucOther	funLargeTrees	funHollowtrees	funLitterCover	funLenFallenLogs	funTreeStem5to10	funTreeStem10to20	funTreeStem20to30	funTreeStem30to50	funTreeStem50to80	funTreeRegen	funHighThreatExotic
210628P1	1692	0.1	0.1	Low	56	305643	6417517	143	2	3	4	5	0	1	50.1	0.3	0.4	0.5	0	0.1	0	0	62	36	0	1	1	0	0	1	45.3

ATTACHMENT C  
EXPERT REPORT EXPECTED PRESENCE OF THREATENED TERRESTRIAL ORCHIDS (DIURIS  
TRICOLOR, PRASOPHYLLUM PETILUM, PTEROSTYLIS CHAETOPHORA) MAXWELL PROJECT

# Expert Report

## Expected Presence of Threatened Terrestrial Orchids (*Diuris tricolor*, *Prasophyllum petilum*, *Pterostylis chaetophora*)

### Maxwell Project



February 2021

Final Report

**Malabar Resources**

Thomas Mitchell Drive  
Muswellbrook NSW 2333

Dr Stephen Bell

Eastcoast Flora Survey  
PO Box 216  
Kotara Fair NSW 2289



## SUMMARY

Maxwell Ventures Management Pty Ltd, a wholly owned subsidiary of Malabar Resources, propose to establish an underground coal mine within EL5460 (Muswellbrook local government area, Hunter Valley, New South Wales), which will require limited ground disturbance across the Project Area. Never-the-less, proposed disturbances associated with planned development may potentially impact on threatened terrestrial orchids (*Diuris tricolor* [Pine Donkey Orchid], *Prasophyllum petilum* [Tarengo Leek Orchid] or *Pterostylis chaetophora* [Rusty Greenhood]) or their habitat. No records of the three target species exist for the Project Area, and after assessment I considered that 1.6 hectares (ha) of the proposed 320 ha (0.5%) ground disturbance area may provide habitat for two of these species (*Diuris tricolor*, *Prasophyllum petilum*). I do not consider that the third species (*Pterostylis chaetophora*) is likely to occur anywhere within the Project Area.

*Diuris tricolor* and *Prasophyllum petilum* occupy extensive geographical ranges outside of the Hunter region, however *Pterostylis chaetophora* is endemic here. Populations for *Diuris tricolor* exist close to the Project Area along Thomas Mitchell Drive, and *Prasophyllum petilum* has historically also been recorded at the eastern end of Thomas Mitchell Drive but that colony is presumed extinct. There are no validated populations of *Pterostylis chaetophora* west of North Rothbury (52 kilometres [km] to the south-east), and further inland records at Mangoola Coal (21 km north-west) and Wingen Maid Nature Reserve (55 km north) also remain unvalidated, and may represent a different species.

Following a site inspection of the Project Area in July 2020, I noted observable differences in the floristic composition of the habitats there to other areas where I have seen the three target orchids, and the predominance of former Box (*Eucalyptus 'albemol'* and *Eucalyptus moluccana*) landscapes within these largely cleared lands conflicts with my experience of orchid habitat elsewhere. Differences were supported by a review of ecological information prepared by Hunter Eco for the Project Area, the mapping of which I found to be accurate and acceptable. Numerical analysis of defined vegetation communities revealed clear differences to habitat from elsewhere in the region known to support *Pterostylis chaetophora*, but show one community similar to habitats typical of *Diuris tricolor* and *Prasophyllum petilum*. Additionally, the close proximity of known populations of *Diuris tricolor* (c. 400 metres [m]) and *Prasophyllum petilum* (1300 m) to the northern section of the Project Area suggests that potential habitat for these may also occur in a second community. I therefore considered that the vegetation units defined by Hunter Eco as Ironbark - Grey Box Grassy Woodland Derived Native Grassland (Unit 9a) and Grey Box – Spotted Gum – Narrow-leaved Ironbark Woodland (Unit 11) provide the only potential habitat for *Diuris tricolor* and *Prasophyllum petilum*, which together comprise 1.6 ha of the Project Area. Subsequently, Dr Colin Driscoll (Hunter Eco) was commissioned by Malabar Resources to conduct a targeted survey of this area during suitable survey conditions in 2020, and none of the target species were found. *Diuris tricolor* and *Prasophyllum* (syn. *Prasophyllum* sp. Wybong) are therefore unlikely to be present within the Project Area.

In support, a detailed assessment of environmental attributes across all Hunter populations of *Diuris tricolor* (n=983), *Prasophyllum petilum* (n=485) and *Pterostylis chaetophora* (n=128) found differences in geological units, soil landscapes, and (for *Pterostylis chaetophora*) annual rainfall when compared to the Project Area. Of these, soil landscapes were seen as particularly important given they best encapsulate the likely distribution of the necessary mycorrhizal fungi across the region, without which no orchids would germinate and prosper. As a surrogate for fungi distribution, an assessment of all terrestrial orchid observation records (common and threatened) for the upper Hunter Valley demonstrated a scarcity of records in and around the Project Area, suggesting that mycorrhizal fungi were therefore also scarce or localised. This lends further credence to my conclusion that no populations of *Diuris tricolor* or *Prasophyllum petilum* are expected across the bulk of the Project Area.



The combination of geology, soil landscape, floristics, rainfall, and general orchid distribution (as a surrogate for mycorrhizal fungi) suggest that the Project Area is unlikely to support extensive populations of *Diuris tricolor* or *Prasophyllum petilum* (syn. *Prasophyllum* sp. Wybong), and none of *Pterostylis chaetophora*. The areas where I considered the former two species may occur occupy <1% of the total disturbance footprint, but subsequent targeted searches in these areas did not record these species.

**Report produced by:**

Dr Stephen A.J. Bell  
Eastcoast Flora Survey  
PO Box 216 Kotara Fair NSW 2289 Australia

Telephone (02) 4953 6523  
Email: [sajbell@bigpond.com](mailto:sajbell@bigpond.com)

**This document should be cited as:**

Bell, S.A.J. (2021) *Expert Report - Expected Presence of Threatened Terrestrial Orchids (Diuris tricolor, Prasophyllum petilum, Pterostylis chaetophora): Maxwell Project*. Unpublished Final Report to Malabar Resources. February 2021. Eastcoast Flora Survey.

© Stephen Bell

This report is copyright. Apart from fair dealing for the purposes of private study, research, criticism or review as permitted under the Copyright Act 1968, no part may be reproduced, copied, transmitted in any form or by any means (electronic, mechanical or graphic) without the prior written permission of the author.

**Document cover:** View across part of the proposed surface development area (top), and target orchids (from left) *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis chaetophora*.

**Report produced for:**

Malabar Resources  
Thomas Mitchell Drive  
Muswellbrook NSW 2333

## Table of Contents

SUMMARY .....	i
1. Introduction .....	1
1.1 Background .....	1
1.2 Project Overview .....	1
1.3 Report Criteria & Structure .....	2
1.4 DPIE Approval to Prepare Expert Report .....	3
2. Criterion (a) - The Relevant Species .....	5
2.1 Legal Status .....	5
2.2 Distribution and Known Populations .....	5
2.2.1 <i>Diuris tricolor</i> .....	6
2.2.2 <i>Prasophyllum petilum</i> .....	7
2.2.3 <i>Pterostylis chaetophora</i> .....	8
2.3 Habitat .....	9
2.3.1 <i>Diuris tricolor</i> .....	9
2.3.2 <i>Prasophyllum petilum</i> .....	12
2.3.3 <i>Pterostylis chaetophora</i> .....	14
2.4 Ecology .....	14
2.4.1 Orchid Detectability .....	14
2.4.2 Mycorrhizal Fungi .....	17
2.4.2.1 <i>Diuris</i> .....	18
2.4.2.2 <i>Prasophyllum</i> .....	18
2.4.2.3 <i>Pterostylis</i> .....	18
2.4.3 Pollination and Seed Production .....	18
2.4.3.1 <i>Diuris</i> .....	19
2.4.3.2 <i>Prasophyllum</i> .....	19
2.4.3.3 <i>Pterostylis</i> .....	19
2.4.4 Reproduction and Dispersal .....	19
2.4.4.1 <i>Diuris</i> .....	20
2.4.4.2 <i>Prasophyllum</i> .....	20
2.4.4.3 <i>Pterostylis</i> .....	20
3. Criterion (b) – Justification for an Expert Report .....	21
3.1 Survey Effort .....	21

3.2 Impact of Drought .....	22
4. Criterion (c) – Likelihood of Species Presence in the Project Area .....	25
4.1 Project Area Attributes .....	25
4.1.1 Land-use History .....	25
4.1.2 Existing Orchid Records .....	25
4.1.3 Vegetation Communities .....	26
4.1.3.1 Plant Community Types .....	26
4.1.3.2 Comparative Numerical Analyses .....	28
4.2 Environmental Analysis .....	32
4.2.1 Project Area .....	34
4.2.2 Target Orchids Records .....	35
4.3 Orchid Diversity: a Surrogate for Mycorrhizal Fungi .....	36
4.4 Field Inspection of the Project Area .....	38
4.5 Concluding Opinion: Are the Target Species Present? .....	44
5. Criterion (d) – Size of Population or Habitat .....	46
6. Criterion (e) – Documents & Data Reviewed .....	49
7. Criterion (f) – Expert Credentials .....	50
8. Conclusion .....	54
9. References .....	56
Appendix 1 – Dr Stephen Bell DPIE Accreditation .....	67
Appendix 2 – Floristic Composition of Grassland Habitat (Bell 2012a) .....	69
Appendix 3 – Biophysical Attributes of the Project Area .....	72
Appendix 4 – Biophysical Attributes of Orchid Locations .....	75
Appendix 5 – Resume: Dr Stephen Bell .....	79
Appendix 6 – Endorsement: Dr Lachlan Copeland .....	91

## 1. Introduction


### 1.1 Background

1. I have been engaged by Resource Strategies on behalf of Maxwell Ventures Management Pty Ltd and Malabar Resources (Malabar) to undertake an expert review in relation to the potential occurrence of three threatened orchids (*Diuris tricolor* [Pine Donkey Orchid], *Prasophyllum petilum* [Tarengo Leek Orchid] and *Pterostylis chaetophora* [Rusty Greenhood]). This review will be incorporated into a Biodiversity Development Assessment Report (BDAR) prepared by Hunter Eco (2019), which addresses the proposed Maxwell Project (the Project). This expert review is as required and in accordance with section 5.3 of the New South Wales (NSW) Government's Biodiversity Assessment Method (BAM) (Department of Planning, Industry and Environment [DPIE] 2020). It aims to determine the habitat suitability of the proposed development lands for the subject orchids.
2. As part of my brief, I have been asked to examine the potential for *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis chaetophora* to occur within lands designated for ground disturbance and/or possible ponding following ground subsidence. These lands occupy 320 hectares (ha), of which approximately 226 ha comprises native vegetation (Hunter Eco 2019). My assessment is required as drought conditions within the Project Area in recent years have impinged on comprehensive surveys for all three orchid species. On 3 July 2020, I undertook an inspection of the Project Area to examine habitat quality and disturbance levels, expanded upon in **Section 4**.

### 1.2 Project Overview

3. Maxwell Ventures (Management) Pty Ltd, a wholly owned subsidiary of Malabar, has consent to develop an underground coal mining operation, known as the Maxwell Project (the Project), located wholly in the Muswellbrook local government area (LGA). It is proposed to undertake underground mining of coal within Exploration Licence 5460, using a combination of both existing and new infrastructure to support underground mining and coal handling activities. Four seams within the Whittingham Coal Measures would be mined using underground bord and pillar (with partial pillar extraction) in the Whynot Seam, and underground longwall extraction in the Woodlands Hill, Arrowfield and Bowfield Seams.
4. As part of proposed works, Malabar intend to undertake a range of actions which may impact on the existing natural environment across five defined areas (the Project Area):
  - expansion of the existing Product Stockpile Extension area to accommodate mined coal;
  - construction of a Surface Development Area to accommodate underground entrance surface facilities and a transport and services corridor;
  - Edderton Rd Realignment, to allow for potential ground subsidence damage;
  - Ancillary Disturbance Area (Ponding) 1, where ground subsidence may result in water ponding; and
  - Ancillary Disturbance Area (Ponding) 2, where ground subsidence may result in water ponding.

5. The focus of my report is on the 320 ha of proposed Project lands lying west of Lake Liddell and Plashett Reservoir, east of Denman and south-east of Muswellbrook (Figure 1 and 2). Note that the full extent of proposed disturbance areas continues through existing post-mine landforms, but only those lands outside of these areas (i.e. that component constituting the ‘controlled action’ under the Commonwealth EPBC Act) are addressed in this report.




**Figure 1 Location of the Project Area (circled) within the context of the Hunter Valley.**

6. To assist in later discussions, this area (the Project Area) has been designated into five geographically separate parcels of land (see Table 1).

### 1.3 Report Criteria & Structure

7. As detailed in the BAM (DPIE 2020), an expert report is required to address the following criteria (Box 3 in section 5.3), and these form the basis of the structure of this report:
- identify the species or population (see Section 2);
  - justify the use of an expert report (see Section 3);
  - justify the likelihood of occurrence of the species or population and prepare a species polygon as per subsection 5.2.5 (see Section 4);



**Figure 2 The Project Area, showing layout of proposed disturbance areas detailed in Table 1. Lands within existing post-mine landscapes are not addressed in this report.**

**Table 1 Land parcels comprising the Project Area.**

Land Parcel	Details	Size (ha)
Product Stockpile Extension	adjacent to the existing stockpile area, c. 500 metres (m) south of Thomas Mitchell Drive	5.2
Surface Development Area	immediately south of Mt Arthur Coal Mine, c. 2.6 kilometres (km) north-west of Plashett Reservoir	302.9
Edderton Rd Realignment	west of existing Edderton Road (south of Saddlers Creek), c. 1.1 km west of its junction with the Golden Highway	10.2
Ancillary Disturbance Area (Ponding) 1	tributary of Saddlers Creek, c. 2.3 km east of Edderton Road	0.5
Ancillary Disturbance Area (Ponding) 2	tributary of Saddlers Creek, c. 0.3 km west of Edderton Road and 2.2 km north of the Golden Highway – Edderton Road intersection	1.5
<b>Total</b>		<b>320.3</b>

- d. estimate the area of habitat (if the species is assessed by area) on the subject land (see **Section 5**), or
- e. estimate the maximum number of mature individuals (as identified in the Credit Calculator) for the subject land. Where the expert report is required because the species is assumed to be present, provide evidence such as a reference site, for this estimation (option d. above undertaken);
- f. include the information considered in making this determination (see **Section 6**); and
- g. state the expert's credentials (see **Section 7**).

#### **1.4 DPIE Approval to Prepare Expert Report**

8. I have been approved to prepare this expert report for all three target species by DPIE, see [Appendix 1](#) and the list of approved biodiversity experts available at <https://www.environment.nsw.gov.au/topics/animals-and-plants/biodiversity/biodiversity-offsets-scheme/experts>.



## 2. Criterion (a) - The Relevant Species

### 2.1 Legal Status

9. *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis chaetophora* are threatened species included in relevant State, Territory and Commonwealth legislation (Table 2). *Diuris tricolor* is listed both as vulnerable in NSW and as an endangered population in the Muswellbrook LGA under the *Biodiversity Conservation Act 2016* (BC Act), while *Prasophyllum petilum* is listed as endangered in NSW (BC Act) and the Australian Capital Territory (ACT) (*Nature Conservation Act 2014*), but under the synonym *Prasophyllum* sp. Wybong (C.Phelps ORG 5269) as critically endangered nationally (*Environment Protection and Biodiversity Conservation Act 1999* [EPBC Act]). *Pterostylis chaetophora* is listed as vulnerable only on the BC Act.

**Table 2 Legal status of target orchids within relevant States and Territories.**

Legislation	<i>Diuris tricolor</i>	<i>Prasophyllum petilum</i>	<i>Pterostylis chaetophora</i>
Commonwealth <i>Environment Protection and Biodiversity Conservation Act 1999</i>	-	Crit. End.	-
NSW <i>Biodiversity Conservation Act 2016</i>	Vul. / EPop.	End.	Vul.
ACT <i>Nature Conservation Act 2014</i>	-	End.	-
VIC <i>Flora and Fauna Guarantee Act 1988</i>	End.	-	-
QLD <i>Nature Conservation Act 1992</i>	LC	-	-

Key: Crit. End. = Critically Endangered; End. = Endangered; EPop. = Endangered Population; LC = Least Concern; Vul. = Vulnerable.

Note: *Prasophyllum petilum* is currently listed nationally on the EPBC Act as Endangered (as *Prasophyllum petilum*) and also as Critically Endangered (as *Prasophyllum* sp. Wybong (C.Phelps ORG 5269), in synonymy), because the Australian Plant Census has not yet accepted this synonym.

10. In recent years, there has been some taxonomic confusion over the identity of *Prasophyllum* plants growing in the upper Hunter (Wybong) area. Following an informal review by NSW orchid taxonomists over the past decade, these plants were placed in synonymy with the more widespread *Prasophyllum petilum* (see PlantNet; <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Prasophyllum~petilum>), a finding also supported by other orchid experts elsewhere in Australia (e.g. Backhouse *et al.* 2019). As a consequence, *Prasophyllum* sp. Wybong (C.Phelps ORG 5269) is now an accepted synonym of *Prasophyllum petilum*, but remains listed as critically endangered under that phrase name on the EPBC Act.

### 2.2 Distribution and Known Populations

11. *Diuris tricolor* and *Prasophyllum petilum* are present and co-occur in the upper Hunter Valley region of NSW, but the two species also occupy considerably wider geographical ranges throughout eastern Australia. In contrast, *Pterostylis chaetophora* is endemic to the Hunter region and occupies a smaller and more coastal distribution. However, historical records of *Pterostylis chaetophora* from the Muswellbrook (2005) and Scone (1998) areas are potentially

indicative also of a more inland extent, but these have not yet been verified. [Figure 3](#) shows images of all three species growing *in situ* in the Hunter region.

- Notes on the distribution of all three species in the following discussion are based on as-held database records from DPIE (extracted under licence ASH20009), augmented with notes and records from published and unpublished reports in the literature (cited accordingly). Records from licence ASH20009 are included with other publicly available data and cited collectively as 'NSW BioNet'.



**Figure 3** Target orchids (from left): *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis chaetophora*.

### 2.2.1 *Diuris tricolor*


- Diuris tricolor* (Pine Donkey Orchid) is a widespread terrestrial orchid, occurring on the western slopes and plains and tablelands of NSW, and also in the Moreton and Darling Downs districts of Queensland (Stanley & Ross 1989; Jones 1993). Populations of *Diuris tricolor* in the upper Hunter Valley between Denman and Muswellbrook essentially form the eastern extent of an east-west trending meta-population extending along the Goulburn River valley to Mudgee ([Figure 4](#)). Records exist for this species at approximately 20 km intervals along this 200 km extent, suggesting that some exchange of genetic material may be occurring with more westerly stands. The Type material of *Diuris tricolor* was collected at Mudgee in the late 1800s.
- The nearest records of *Diuris tricolor* to the Project Area are an observation in 2009 from land off Saddlers Creek (c. 600 m south of the proposed Ancillary Disturbance Area 1; NSW BioNet; also noted in Cumberland Ecology 2015 and relocated by Hunter Eco 2020), and several observations from 2004-2015 within the Mt Arthur complex along Thomas Mitchell Drive (c. 400 m north of the proposed Product Stockpile Extension area; NSW BioNet; Hunter Eco 2013). The 2009 observation recorded two plants in that year, and follow up surveys found none in November 2010 (outside of peak flowering) but 'at least 30' in 2011 (Cumberland Ecology 2015). Searches since that time have failed to locate any individuals, although several years of drought during this period may have reduced flowering and increased grazing pressure. Along Thomas Mitchell Drive, one observation of *Diuris tricolor* in 2004, three in 2009, one in 2012 and 70 in 2015 (within the Drayton Wildlife Refuge) attest to a sizeable population in that

vicinity. No plants have been observed since 2015 (Hunter Eco 2019), detection again likely to have been negatively impacted by drought.

15. A single, small disjunct population of *Diuris tricolor* was also recorded at North Rothbury in 2016 (noted in Bell 2017a), and represents the most easterly population known within NSW. No individuals have been observed at that location since 2016 due to dry conditions. North Rothbury lies more than 50 km to the south-east of the Project Area.
16. Elsewhere in NSW, *Diuris tricolor* is extensive across the north, central and south western slopes, and extends into south-eastern Queensland. A single record from the Hume region of Victoria suggests that the species is very rare in that state, and indeed Backhouse *et al.* (2016) indicate that it is known from just three plants.

### 2.2.2 *Prasophyllum petilum*

17. *Prasophyllum petilum* (Tarengo Leek Orchid) occupies a smaller distributional range than *Diuris tricolor*, with most records from the ACT but with outliers in the Kandos, Denman, Premer and Inverell districts on the tablelands and western slopes of NSW. Until recently, Hunter Valley plants were considered a distinct but un-named taxon, *Prasophyllum* sp. 'Wybong' (C.Phelps ORG 5269), but are now placed in synonymy with *P. petilum* by NSW taxonomic authorities. In support, Backhouse *et al.* (2019) do not include *Prasophyllum* sp. 'Wybong' in their comprehensive list of Australian orchid taxa, despite the inclusion of three other un-named taxa with close affinities to *P. petilum*. The Type material of *P. petilum* was collected from Hall (in the ACT) in 1988.
18. The nearest known occurrence of *Prasophyllum petilum* to the Project Area is a site near the eastern end of Thomas Mitchell Drive (NSW BioNet). This observation (made of 10 plants in 1999) was the first of the species for the Hunter Region, and advice from the original observers has confirmed the identity of those plants against current taxonomy (L. Copeland pers. comm.). Surveys subsequent to the initial discovery revealed no plants in 2000, five plants in 2001, no plants in 2002, nine plants in 2003, no plants in 2005 and one plant in 2005 (B. Holzinger pers. comm.). Occasional searches of the site since 2005 have failed to detect any plants, suggesting that the population may now be extinct (B. Holzinger pers. comm.).
19. All other Hunter Valley observations of *Prasophyllum petilum* are from Mangoola Coal (Wybong), approximately 25 km to the north-west, and no other observations are evident from other parts of the Hunter Valley (NSW BioNet; Australasian Virtual Herbarium [AVH]). Plants have been recorded at Mangoola in most years since 2009, although numbers were very low during the drought years of 2017-2019 (Bell 2020a).
20. Outside of the Hunter catchment, the next nearest populations of *Prasophyllum petilum* occur near Kandos, some 140 km to the south-west, and Premer 190 km to the north-west (Figure 4). Hunter Valley populations of *Prasophyllum petilum* are consequently isolated from all others, and opportunities for genetic exchange are minimal. Note that Jeanes (2015a) considered similar populations of *Prasophyllum* in Victoria to represent a different taxon (now described as *P. argillaceum*: Jones & Rouse 2018), implying that *Prasophyllum petilum* is endemic to NSW, a view also supported by Backhouse *et al.* (2019).



**Figure 4** Distribution of *Diuris tricolor* (+) (left), *Prasophyllum petilum* (x) (right) and *Pterostylis chaetophora* (□) (right) across eastern Australia, shown relative to the Project Area (circled, 50 km radius). Data from AVH and NSW BioNet, extracted July 2020. Note that near-coastal records of *Diuris tricolor* in south-eastern NSW are erroneous (historical collections with poor positional accuracy), and that Victorian records of *Prasophyllum petilum* represent a different taxon (now described as *P. argillaceum*: Jones & Rouse 2018).

### 2.2.3 *Pterostylis chaetophora*

21. *Pterostylis chaetophora* (Rusty Greenhood) is endemic to the broader Hunter Region, extending from southern Taree to Newcastle and Cessnock (Figure 4). The species was previously also thought to occur in Queensland, but those populations have been redetermined as a distinct species with close affinities to *Pterostylis chaetophora* (Backhouse *et al.* 2019). Additionally, two early collections in the AVH from Sydney, and an observation record from

Glenhaven (1949), are thought to be in error or are now locally extinct. Both Hosking and James (1998) and Hunter (2008) report populations of *Pterostylis chaetophora* from Warrabah National Park near Tamworth, but these are considered to be misidentifications as no recent observations have been made there (L. Copeland pers. comm.). The Type material for *Pterostylis chaetophora* was grown from a specimen collected at Neath, near Cessnock in the mid-Hunter Valley in the early 1980s (Clements 1989).

22. NSW BioNet includes 33 distinct observation records, with most occurring in the lower Hunter Valley (east from Cessnock). For the upper Hunter Valley, two collections lodged at the Australian National Herbarium (Canberra) have been identified as *Pterostylis chaetophora*, but there are no other records. The first of these collections was made by John Hoskings in 1998 from Wingen Maid Nature Reserve (c. 50 km north-north-west of the Project Area), and the second in 2005 by Dougall Herd from what is now a biodiversity offset for Mangoola Coal (c. 20 km west-north-west of the Project Area). It is unknown if these populations remain extant or if they may represent the closely related *Pterostylis praetermissa* (L. Copeland pers. comm.). Surveys planned for the 2020 flowering season aim to clarify the identification of orchids at both locations.
23. Consequently, the nearest known confirmed records of *Pterostylis chaetophora* to the Project Area is at Rothbury (c. 53 km south-east of the Project Area). This population has been monitored annually for the last three years, showing heavy grazing of tubers by White-winged Choughs (*Corcorax melanorhamphos*) (Bell 2020b). Columbey National Park, situated near Clarendtown, currently supports the most extensive populations of *Pterostylis chaetophora* (Bell & Hillier 2020), and lies 80 km to the east-south-east of the Project Area.

## 2.3 Habitat

### 2.3.1 *Diuris tricolor*


24. A range of habitats have been documented for *Diuris tricolor* throughout its range (Table 3), although few studies provide sufficient detail on co-occurring ground layer species. This makes it difficult to identify potential habitat at the local scale, although not insurmountable. Most texts document favoured habitat as grassy *Callitris* woodlands, although in Queensland it is 'eucalypt open forest'. In a study of remnant vegetation stands in the South Western Slopes of NSW, Burrows (1999) recorded *Diuris tricolor* at several sites, but all within *Callitris glaucophylla* dominated vegetation. J. Hunter (2010) located the species in areas associated with ironbark and Bullock in the Pilliga area, while Cunningham *et al.* (2011) noted habitat as *Eucalyptus populnea* for the Tottenham area (west of Dubbo). Clearly, a diversity of habitats support *Diuris tricolor* across the state, and this information alone cannot be used to generalise occupied habitat at specific locations.
25. First-hand experience of *Diuris tricolor* in the Hunter Valley is therefore influential in determining the suitability of an area to support this species. Field evidence and unpublished data from subpopulations of *Diuris tricolor* at Mangoola (near Denman) and Muswellbrook (Bulga Coal's Condran offset property) suggest that it occurs most commonly within grassy woodlands and grasslands derived from former Ironbark (*Eucalyptus crebra*) or Dawson's Box (*Eucalyptus dawsonii*) woodlands, and with minimal amounts in Box (*Eucalyptus 'albemol'*, a purported hybrid between *E. albens* and *E. moluccana*) woodlands. Note that in the central and upper Hunter, populations of *Eucalyptus 'albemol'* have historically and contemporarily been

referred to as either or both of the two supposed parent species, but ongoing taxonomic work suggests this entity to be a distinct species with no evidence of hybridisation (S. Bell & C. Driscoll unpubl. data). At the currently known eastern limit of distribution (North Rothbury), *Diuris tricolor* was recorded in 2016 in open forest of *Eucalyptus crebra*, *Corymbia maculata* and *Eucalyptus fibrosa*, although no individuals have been seen at that location since (Bell 2017a).

**Table 3 Habitat documented for *Diuris tricolor*.**

Habitat	Location	Source
<b>Hunter Valley</b>		
native grassland in areas not subject to intensive grazing	Hunter Valley generally	Umwelt 2011a, 2011b, 2013
derived grasslands of <i>Aristida/Cymbopogon</i> ; <i>Bothriochloa/Carthamus/Danthonia</i> ; <i>Dichanthium/Sporobolus/Chloris</i> woodlands of <i>Eucalyptus crebra</i> , <i>E. dawsonii</i> or <i>Allocasuarina luehmannii</i>	Wybong	Bell 2012a
disturbed grassland, often adjoining woodland and forest	Wybong	Umwelt 2012
grassland/open woodland	Wybong	Herd & Herd 2005
grassland	Wybong	Abel Ecology 2005
disturbed grassland	Wybong	Umwelt 2006
grassy woodland within Hunter Floodplain Red Gum Woodland (Threatened Ecological Community [TEC])	Muswellbrook LGA	Cumberland Ecology 2014
<i>Eucalyptus crebra</i> , <i>Corymbia maculata</i> , <i>E. fibrosa</i> grassy forest	North Rothbury	Bell 2017a
ungrazed grasslands on soils of low fertility	Ulan	Ecovision Consulting 2008
derived grassland within <i>Eucalyptus crebra</i>	Condran offset (Muswellbrook)	Bell <i>et al.</i> 2020
<b>Outside of the Hunter Valley</b>		
grassy <i>Callitris glaucophylla</i> woodlands	general	Jones 1993; Burrows 1999; Bishop 2000; Cameron <i>et al.</i> 2014; Jeanes 2015b
<i>Callitris glaucophylla</i> , <i>Eucalyptus populnea</i> , <i>Eucalyptus intertexta</i> , Ironbark and <i>Acacia</i> shrubland. The understorey is often grassy with herbaceous plants such as <i>Bulbine</i> species	general	URS 2009; Jacobs 2017
open grassy woodland often associated with Box-Gum Woodland [TEC]	general	Cumberland Ecology 2014
eucalypt open forest	Queensland	Stanley & Ross 1989
<i>Eucalyptus sideroxylon</i> , <i>E. crebra</i> and <i>Allocasuarina luehmannii</i>	Pilliga (NSW)	J. Hunter 2010
<i>Eucalyptus populnea</i>	west of Dubbo (NSW)	Cunningham <i>et al.</i> 2011

26. Using the Mangoola Coal population of *Diuris tricolor* as a case study, an assessment of vegetation communities supporting 975 GPS-recorded *Diuris tricolor* locations (n=5120 individual orchids) was undertaken (unpubl.). This involved intersecting in GIS point locations against a pre-1750 vegetation map prepared for the site in 2013 (unpubl. data). In that study, twelve vegetation communities were mapped on the basis of extensive field reconnaissance, where remnant paddock trees and landscape position were used to extrapolate across highly cleared lands. Of those twelve communities, *Diuris tricolor* was found to occur in five, with 61% of all records (n=599) and 60% of all individuals (n=3089) occurring within present or former Ironbark (*Eucalyptus crebra*) Woodland (Figure 5).



**Figure 5** Number of *Diuris tricolor* individuals (n=5120) by pre-1750 vegetation type within the 1492 ha Mangoola Coal study area (unpubl. data). Note that *Diuris tricolor* records almost all located within derived native grasslands (DNGs) from these vegetation types.

27. The next most favoured habitat at Mangoola was Dawson's Box (*Eucalyptus dawsonii*) Woodland with 29% of records (n=280) and 33% of individuals (n=1684), and these two communities combined supported >90% of all *Diuris tricolor* records and individuals. Three other communities comprised relatively minor *Diuris tricolor* habitat: Box (*Eucalyptus 'albemol'*) Woodland (7% of records), Rough-barked Apple (*Angophora floribunda*) Woodland (2% of records), and Red Gum (*Eucalyptus blakelyi*) Woodland (1% of records). In the absence of other similar habitat data from elsewhere in the Hunter Valley, the Mangoola study area provides a strong landscape-scale indication that *Diuris tricolor* is most closely associated with *Eucalyptus crebra* or *Eucalyptus dawsonii* grassy woodlands.
28. At a finer scale, a floristic analysis of derived grasslands undertaken at Mangoola Coal between 2009 and 2011 (comprising 168 plots sampled over 2000 ha) found that *Diuris tricolor* occurred within three of seventeen grassland types, in descending order of importance (Bell 2012a, Bell submitted 1):
- *Aristida/ Cymbopogon* Grassland (Unit 2);
  - *Bothriochloa biloba/ Carthamus/ Danthonia* Grassland (Unit 4); and
  - *Dichanthium/ Sporobolus/ Chloris* Grassland (Unit 1a).
29. In that study, *Diuris tricolor* was also sparingly present in three woodland communities, those characterised by *Eucalyptus crebra*, *Eucalyptus dawsonii* or *Allocasuarina luehmannii*.

Combined, the three derived grassland habitats defined encompassed a significantly large proportion of the grasslands included in that study (84% of 1069 ha). Detailed floristic compositions for each of these key grassland communities are replicated in [Appendix 2](#). Knowledge gained from the Mangoola floristic analysis of grassland types has been incorporated into my assessments of suitable *Diuris tricolor* habitat at the Maxwell Project discussed later in this report.

### 2.3.2 *Prasophyllum petilum*

30. Information on the habitat of *Prasophyllum petilum* throughout its range is brief but documents variable associations ([Table 4](#)). When describing the species, Jones (1991) reported the known habitat at that time (the Type locality only, in the ACT) as being “moist grassy patches in sparse woodland developed on fertile soils”, while Bishop (2000) describes it as remnant *Themeda* grassland on silty clay loams. The national recovery plan for this species (Department of Environment, Climate Change and Water [DECCW] 2010) provides more detail on floristic associations at the five known sites for which it was written, mostly on the Southern and Central Tablelands of NSW.
31. Notes associated with collections included in AVH indicate that most southern records of *Prasophyllum petilum* occur in grasslands dominated by *Themeda australis*, *Bothriochloa* spp. and *Danthonia* spp, with associated forbs of *Bulbine* sp., *Dichopogon* sp., *Wurmbea* sp., *Swainsona* sp., *Pimelea curviflora*, *Chrysocephalum* sp., *Ajuga australis*, *Craspedia* sp., *Stackhousia monogyna*, *Eryngium* sp., *Burchardia* sp., *Arthropodium* sp., and *Juncus* sp. Northern records occur in grassland of *Aristida* sp., *Themeda australis* and *Stackhousia monogyna*. With the exception of populations on the North Western Slopes, these habitats at collection locations are very different to those where *Prasophyllum petilum* occurs in the Hunter Valley. In this region plants occur most commonly in grasslands derived from former Ironbark (*Eucalyptus crebra*), Dawson’s Box (*Eucalyptus dawsonii*) and Box (*Eucalyptus ‘albemol’*) woodlands, co-occurring with species such as *Cymbopogon refractus*, *Aristida ramosa*, *Dichanthium sericeum* and *Chloris ventricosa*.
32. Once again, using Mangoola Coal as a case study, an assessment of the vegetation communities supporting 759 GPS-recorded *Prasophyllum petilum* locations (n=4073 individual orchids) was undertaken. This involved intersecting in GIS each point location against a pre-1750 vegetation map prepared for the site in 2013 (unpubl. data). In that study, twelve vegetation communities were mapped on the basis of extensive field reconnaissance, where remnant paddock trees and landscape position were used to extrapolate across highly cleared lands as required. Of those twelve communities, *Prasophyllum petilum* was also found to occur in five (the same as for *Diuris tricolor*), with 58% of all records (n=442) and 59% of all individuals (n=2413) occurring within present or former Ironbark (*Eucalyptus crebra*) Woodland ([Figure 6](#)).
33. As for *Diuris tricolor*, the next most favoured habitat was Dawson’s Box (*Eucalyptus dawsonii*) Woodland with 22% of records (n=164) and 20% of individuals (n=831), and these two communities combined supported 80% of all *Prasophyllum petilum* records and individuals. Box (*Eucalyptus ‘albemol’*) Woodland supported 17% of records (n=130) and 19% of individuals (n=764). Two other communities comprised minor *Prasophyllum petilum* habitat: Red Gum (*Eucalyptus blakelyi*) Woodland, and Rough-barked Apple (*Angophora floribunda*) Woodland (<4% combined). In the absence of other similar habitat data from elsewhere in the Hunter Valley, the Mangoola study area provides a strong landscape-scale indication that *Prasophyllum petilum* is most closely associated with *Eucalyptus crebra* grassy woodlands.




**Table 4 Habitat documented for *Prasophyllum petilum*.**

Habitat	Location	Source
Hunter Valley		
derived grasslands of <i>Aristida/Cymbopogon</i> ; <i>Bothriochloa/Carthamus/Danthonia</i> ; <i>Dichanthium/Sporobolus/Chloris</i> ; woodlands of <i>Eucalyptus crebra</i> , <i>E. dawsonii</i> or <i>Allocasuarina luehmannii</i>	Wybong	Bell 2012a
open eucalypt woodland and grassland	general	Wildthing 2011
Outside of the Hunter Valley		
remnant <i>Themeda</i> grassland on silty clay loams	general	Bishop 2000
shrubby and grassy habitats in dry to wet soil, in open eucalypt woodland and grassland	general	Umwelt 2013; Eco Logical Australia 2015
Grassy Box Woodlands with fertile to moderately fertile soils on undulating terrain	general	FloraSearch 2014
wet grassy woodlands on fertile ground	Southern Tablelands region (NSW & ACT)	Rouse 2002
moist grassy patches in sparse woodland developed on fertile soils	Type location (ACT)	Jones 1991
grassy woodland of <i>Eucalyptus pauciflora</i> and <i>E. aggregata</i> , with a sparse shrub layer of <i>Hakea microcarpa</i> , <i>Acacia dealbata</i> and <i>Leptospermum brevipes</i> and a ground layer of <i>Poa sieberiana</i> , <i>Themeda australis</i> and <i>Schoenus apogon</i>	Captains Flat cemetery (NSW)	DECCW 2010
grassy woodland of <i>Eucalyptus blakelyi</i> and <i>E. melliodora</i> , over <i>Poa sieberiana</i> and <i>Themeda australis</i>	Hall cemetery (ACT)	DECCW 2010
grassy woodland of <i>Eucalyptus blakelyi</i> and <i>E. melliodora</i> , over <i>Themeda australis</i> and <i>Sorghum leiocladum</i>	Ilford cemetery (NSW)	DECCW 2010
natural grassland of <i>Bothriochloa macra</i> , <i>Pentapogon quadrifidus</i> , <i>Austrodanthonia</i> spp., <i>Themeda australis</i> , <i>Schoenus apogon</i> , <i>Drasera peltata</i> , <i>Sebaea ovata</i> and <i>Haloragis heterophylla</i> on a treeless grassy plain	Tarengo TSR (NSW)	DECCW 2010
a treeless frost hollow, surrounded by <i>Eucalyptus pauciflora</i>	Steves TSR (NSW)	DECCW 2010

34. Floristic analysis of derived grasslands undertaken at Mangoola Coal between 2009 and 2011 (comprising 168 plots sampled over 2000 ha) found *Prasophyllum petilum* occurring in three of seventeen grassland types, in descending order of importance (Bell 2012a; Bell submitted 1):

- *Aristida/ Cymbopogon* Grassland (Unit 2);
- *Bothriochloa biloba/ Carthamus/ Danthonia* Grassland (Unit 4); and
- *Dichanthium/ Sporobolus/ Chloris* Grassland (Unit 1a).



**Figure 6** Number of *Prasophyllum petilum* individuals (n=4073) by pre-1750 vegetation type within the 1492 ha Mangoola Coal study area (unpubl. data). Note that *Prasophyllum petilum* records almost all located within DNGs from these vegetation types.

35. *Prasophyllum petilum* was also occasionally present in three woodland communities, those characterised by *Eucalyptus crebra*, *Eucalyptus dawsonii* or *Allocasuarina luehmannii*. Combined, the three derived grassland habitats defined encompassed a significantly large proportion of the grasslands included in that study (84% of 1069 ha). Knowledge gained from the Mangoola floristic analysis of grassland types (see [Appendix 2](#)) has been incorporated into my assessments of suitable orchid habitat at the Maxwell Project discussed later in this report.

### 2.3.3 *Pterostylis chaetophora*


36. There is little information available on habitat for the Hunter endemic *Pterostylis chaetophora*, although recent monitoring work associated with the NSW Government's *Saving our Species* initiative has collated some information (Bell 2020b). For Columbe National Park, *Pterostylis chaetophora* was found to predominantly occur in vegetation described as Floodplain Redgum-Box Forest and Lower Hunter Spotted Gum – Ironbark Forest (Bell & Hillier 2020: [Figure 7](#)). Paget (2008) provided brief notes on habitat for several other populations of the species, which has been added to and expanded upon in Bell and Hillier (2020). [Table 5](#) summarises habitat for *Pterostylis chaetophora* from all known locations.

## 2.4 Ecology

### 2.4.1 Orchid Detectability

37. The unpredictability of flowering in orchids from year-to-year is a widely recognised trait in this group of plants (e.g. Gillman & Dodd 1998; Kindlmann & Balounova 2001; McCormick & Jacquemyn 2014), and this is commonly governed by weather (e.g. Wells *et al.* 1998; Kindlmann & Balounova 2001; Pfeifer *et al.* 2006). Many species fail to emerge during dry phases, and although viable populations may persist underground, their presence above ground often leads to an incorrect assumption that no plants are present. In general terms, low rainfall in the three months leading up to flowering place individual orchids under stress, meaning that flowering may be postponed for that season for all but the most robust individuals. Because of this trait,

terrestrial orchids have been described as ‘time-travellers’ (Brundrett 2016), encapsulating the uncertainty in determining their presence in any given area.




**Figure 7** Number of *Pterostylis chaetophora* individuals (n=544) by vegetation type within the 720 ha Columbey National Park study area (after Bell & Hillier 2020).

**Table 5** Habitat documented for *Pterostylis chaetophora*.

Habitat	Location	Source
forest of <i>Eucalyptus fibrosa</i> , <i>Corymbia maculata</i> and <i>Eucalyptus umbra</i>	Columbey National Park, Pindimar, Raymond Terrace, Beresfield	Bell & Hillier 2020
forest of <i>Eucalyptus moluccana</i> and/or <i>Eucalyptus amplifolia</i>	Columbey National Park	Bell & Hillier 2020
shrubby forest of <i>Corymbia maculata</i> and <i>Eucalyptus fibrosa</i> , or <i>Eucalyptus tereticornis</i> , <i>Eucalyptus punctata</i> and <i>Angophora floribunda</i> in dry drainage lines	Kurri	Bell & Hillier 2020
forest of <i>Eucalyptus tereticornis</i> , <i>Angophora floribunda</i> and <i>Eucalyptus crebra</i>	North Rothbury	Bell (2020b)
<i>Eucalyptus propinqua</i> and <i>Eucalyptus microcorys</i> shrubby forest	Purfleet	Paget 2008
grasslands derived from cleared <i>Eucalyptus propinqua</i> , <i>Eucalyptus acmenoides</i> , <i>Eucalyptus microcorys</i> , <i>Eucalyptus placita</i> and <i>Eucalyptus siderophloia</i> forest	Burrell Creek	Paget 2008
forests of <i>Eucalyptus placita</i> , <i>Eucalyptus siderophloia</i> , <i>Eucalyptus paniculata</i> and <i>Corymbia maculata</i> , or <i>Eucalyptus amplifolia</i> and <i>Eucalyptus moluccana</i>	Twelve Mile Creek	Paget 2008

38. Part of the difficulty of detection experienced during drought years is the added stress placed on emerging orchids by herbivores searching for palatable foods: orchids may well emerge every year in some species but they may be quickly consumed by grazing mammals, birds or invertebrates. For example, Duncan and Moloney (2018) found that for the threatened *Diuris fragrantissima* good rainfall increased the probability of flowers setting seed and decreased the probability that plants would be browsed.
39. As a rule of thumb, dry winters in the Hunter Valley generally result in below average flowering in terrestrial orchids, and this has been shown for both *Diuris tricolor* and *Prasophyllum petilum* (Bell 2019a, b). The relationship between rainfall and flowering in these species has been highlighted over a ten-year translocation project undertaken at Mangoola Coal (Bell 2019a, b; Bell 2020; also reported annually in reports to Mangoola Coal). Over the course of nine years of monitoring, the July-to-August pre-flowering rainfall transitioned from three years of near- or below-average rainfall, to three years of above-average rainfall, and three years of well-below average rainfall. Dry years have been reflected in low rates of detection within recipient plots, while wetter years have shown an increase in detection (Figure 8). There are of course other factors contributing to the extent of orchid detection observed (expanded upon in Bell 2020d), but for these two species there is a clear trend associated with winter rainfall. Of the nine recipient plots, all displayed lower detection rates in the drought years of 2017-2019, following three seasons of above average winter falls. A similar downward trend was observed for the five recipient plots (n=440) established within mine rehabilitation, monitored over 3-4 years since 2015, and four control plots of naturally occurring orchids monitored since 2016 (data not presented here).



**Figure 8** Rainfall received (July to August) and orchid detection (n=2,592) during the course of monitoring across nine recipient plots within derived grassland at Mangoola Coal, over a period of four to nine years (after Bell 2019a).

40. Surveying for orchids requires repeat visits to an area to be confident in the presence and magnitude of any residing orchid populations. This is particularly the case for those species where flowering can be protracted over several weeks, although most do experience a 'peak' period (e.g. Yare *et al.* 2020). Vizer (2013) found peak flowering in *Diuris tricolor* and

*Prasophyllum petilum* at Mangoola to occur from mid- to late-September, but that less than 20% of plants would be flowering on any particular day at this time. This implies that a 'one-off' survey, even if conducted on the day of peak flowering, would likely overlook more than 80% of individuals in the population. Capsule production was also found during his study to occur in less than 3% of plants for both species, with herbivory identified as an important limiting factor in seed production. However, peak flowering time likely varies year-to-year, and will depend on available soil moisture and other site conditions. For example, during the 2015 flowering season at Mangoola, *Diuris tricolor* was found to peak in the second week of September, but *Prasophyllum petilum* flowering was still increasing in the first week of October (Bell 2020d).

41. For *Prasophyllum petilum*, Wilson *et al.* (2016) analysed annual monitoring data over a 25-year period from the largest known population on the southern tablelands of NSW, and identified the incidence of frost (nights  $\leq -4^{\circ}\text{C}$ ) as being instrumental in preventing flowering in any one season. Frost damage to emerging plant parts prior to reaching flowering stage prevents detection during monitoring surveys, influencing orchid counts. Warm winters are consequently of benefit to the orchids in that population, although it is unknown if the same applies to the Hunter Valley population where frosts are rarer.
42. Detectability in *Pterostylis chaetophora* appears less reliant on rainfall, although monitoring over three years has shown a weak (but untested) trend in this regard (Bell 2020b). The more coastal distribution of this species suggests that water stress through inadequate rainfall is less likely to influence orchid emergence and persistence, although heavy grazing of tubers by White-winged Choughs has occurred during recent drought conditions at North Rothbury. The geographical range of many members of subgenus *Oligochaetochilus* extends into drier inland habitats, demonstrating some level of adaptation to dry and harsh environments (Kuitert & Findlater-Smith 2017).

#### 2.4.2 Mycorrhizal Fungi

43. Orchid presence in any area is dependent on the availability of co-occurring mycorrhizal fungi present within the soil, and often different fungi are required by different orchid species (Waterman & Bidartondo 2008; McCormick *et al.* 2018), yet co-existence of several orchid species is still possible (Waud *et al.* 2016). Mycorrhizal fungi are necessary for nutrient transfer between soil and orchid, and they are also essential in aiding the germination of orchid seeds (Brundrett 2006). The minute size of orchid seeds means they carry no nutrients and hence associating with the correct fungi immediately after dispersal is imperative. Weston *et al.* (2005) noted a high degree of specificity between a particular species of orchid and their associated species of mycorrhiza, but that there are also commonalities between and within genera. Brundrett (2006) stresses the need for fungi and pollinators with orchids in his three key dimensions of orchid presence: differing levels of interactions between the three dimensions determine the specificity and availability or suitability of habitat.
44. Without intensive survey for the relevant mycorrhizal fungi present in an area, there is no way of knowing whether or not a specific site is capable of supporting an orchid population of any species. Research has shown that mycorrhizal fungi may be widespread in the landscape and occur in a range of habitats, but that it may also be patchy and is not necessarily reflected in observable orchid populations (e.g. Brundrett *et al.* 2003; McCormick & Jacquemyn 2014; Voyron *et al.* 2016). The assumption, therefore, that all potential orchid habitat based on biophysical characteristics (e.g. soil type, elevation, rainfall, co-occurring plant species) can be

equally occupied by a specific orchid species is misleading: without fungi, seed will not germinate and plants will not prosper. Investigation into whether specific abiotic variables, such as rainfall or geological substrate, govern the geographical distribution of mycorrhizal fungi has been identified as a priority for research (Jacquemyn *et al.* 2017), although this work is still in its infancy.

#### 2.4.2.1 *Diuris tricolor*

45. For *Diuris* sp., the *Tulasnella* genus (family Tulasnellaceae) is the most important mycorrhizal fungi (Weston *et al.* 2005; Smith *et al.* 2010). At Mangoola Coal, seed-baiting techniques were used by Vizer (2013) in an attempt to map the distribution of mycorrhizal fungi, and he found that the distribution of *Diuris tricolor* was actually more restricted than the relevant fungi. This implies that (for the Mangoola area) there may be extensive suitable habitat, complete with mycorrhizal fungi, within a wider area than is currently known to support the species. Similarly, Tierney *et al.* (2017) observed differences in seed germination within soil sampled from different subpopulations of *Diuris platycheila*, attributable either to differing concentrations of fungal hyphae within each soil sample or that some samples supported superior strains of fungi.

#### 2.4.2.2 *Prasophyllum petilum*

46. Weston *et al.* (2005) suggest that the genus *Ceratobasidium* (family Ceratobasidiaceae) is the likely mycorrhizal fungi for *Prasophyllum* sp., supported partially by work on two threatened species in Victoria by McQualter *et al.* (2007) where *Ceratobasidium cornigerum* (and a *Rhizoctonia* sp. were isolated). Within the Hunter Valley, mycorrhizal seed-baiting for *Prasophyllum petilum* was not successful in the study of Vizer (2013), which is not unusual for this genus, and further research in this species is required.

#### 2.4.2.3 *Pterostylis chaetophora*

47. Like *Prasophyllum* sp., the genus *Ceratobasidium* is thought to be the likely mycorrhizal fungi required for *Pterostylis* sp. (Weston *et al.* 2005). Jusaitis and Sorensen (1993) isolated mycorrhizal fungi from *Pterostylis arenicola* (closely related to *Pterostylis chaetophora*), but did not identify the fungi. Few other studies within this genus are evident in the literature.

### 2.4.3 Pollination and Seed Production

48. Pollination in *Diuris*, *Prasophyllum* and *Pterostylis* sp. (and most other orchids) is enacted by insects (Adams & Lawson 1993; Weston *et al.* 2005; Hawkeswood 2006). Many orchids rely on mimicry to deceive unsuspecting insects (Adams & Lawson 1993; Schiestl 2005), either by the development of nectar-bearing flowers that appear identical to those of co-occurring species in their habitat (food mimicry), or by individual flowers resembling (often with the aid of pheromones) the females of certain insects (sexual mimicry). Other species offer a nectarivorous reward, which when combined with the appropriate scent attract pollinating insects.
49. Once pollination has been enacted, the development of seed capsules progresses over the following weeks, although plants still remain vulnerable to grazing and desiccation during this time. Some studies have shown that, despite relatively high levels of historical and current-day fragmentation of landscapes, pollinators are still able to forage amongst orchids to produce seed (e.g. Brundrett 2019). For many nectarless orchids, this is reliant on the presence of co-occurring food plants to supply the necessary food, and for species with highly specific pollinators this is particularly important (Phillips *et al.* 2015; Brundrett 2019). However, fragmentation and loss of pollinator habitat is widely recognised as detrimental to most orchid

populations (Wraith & Pickering 2018; Phillips *et al.* 2020). Based on observations made at orchid translocation sites at Mangoola Coal over several years (e.g. Bell 2016), capsule development is unhindered in *Diuris tricolor* and *Prasophyllum petilum* despite close proximity to an active coal mine.

#### 2.4.3.1 *Diuris tricolor*

50. Most *Diuris* sp. mimic co-occurring species of pea (Fabaceae) to attract pollinators (e.g. Backhouse *et al.* 2016; Scaccabarozzi *et al.* 2018), and for *Diuris tricolor* in the Hunter Valley this is likely to be *Templetonia stenophylla* or *Daviesia genistifolia* (pers. obs.; Vizer 2013). Weston *et al.* (2005) indicate that the pollinators of *Diuris* sp. are likely to be various colletid bees from the *Trichocolletes* and *Leioproctus* genera, although Walker (1997) suggested opportunistic native halictid bees were potential pollinators of *Diuris basaltica*. Indsto *et al.* (2006) found male bees of the species *Trichocolletes venustus* to be the primary pollinators of *Diuris maculata* in Sydney.

#### 2.4.3.2 *Prasophyllum petilum*

51. *Prasophyllum* sp. employ a different strategy to attract pollinators, using nectar and scent to entice various insects to pollinate and be rewarded with food. The likely pollinators of *Prasophyllum* sp. are thought to be colletid and halictid bees, ichneumonid, tiphiid, scoliid and sphecid wasps, syrphid flies, and beetles (Weston *et al.* 2005). For *Prasophyllum odoratum*, Bernhardt & Burns-Balogh (1986) found that polytrophic flies (family Syrphidae) and opportunistic male bees in the genus *Leioproctus* (family Colletidae) were the principle pollinators.

#### 2.4.3.3 *Pterostylis chaetophora*

52. For *Pterostylis* sp., pollination through sexual deception has been demonstrated for several species (e.g. Bernhardt 1995; Phillips *et al.* 2014; Thalwitzer *et al.* 2018; Reiter *et al.* 2019), and the greenish to rusty colouration of flowers in this genus suggest fungus gnats (order Diptera) as the primary pollinators (Vogel 1973). Kuiter and Findlater-Smith (2017) provide support for this in their overview of pollinators of the Victorian members of *Pterostylis* sp., concluding that fungus gnats (families Mycetophilidae and Sciaridae) almost always enact pollination in these orchids. In their study, Reiter *et al.* (2019) found *Pterostylis boormanii* and *Pterostylis basaltica*, both closely related to *Pterostylis chaetophora* within section and subgenus *Oligochaetochilus* (Janes & Duretto 2010), were pollinated by male fungus gnats (*Xenoplasyra conformis*) of the family Keroplatidae. However, Kuiter and Findlater-Smith (2017) found the ten Victorian members of subgenus *Oligochaetochilus* were all pollinated by one of five species of fungus gnat from a different genus, *Orfelia* (family Mycetophilidae). Both investigations suggest that multiple pollinating fungus gnats may be responsible, perhaps related to geographical location and/or specific habitat requirements, and such a strategy potentially allows a wider distribution in this subgenus.

### 2.4.4 Reproduction and Dispersal

53. Reproduction in orchids can occur sexually or asexually, through the production of seed or via tuber multiplication ('daughter tuberosids') or through annual replacement. Following sexual reproduction, dispersal limitation is an important factor determining orchid distribution (McCormick & Jacquemyn 2014). Orchid seeds are very small ('dustlike'), produced in exceptionally large numbers within capsules, are very light (Arditti & Ghani 2000; Shefferson *et al.* 2020) and, in some situations, are capable of dispersion by wind over vast distances (Phillips

*et al.* 2020). In most cases, however, the low stature of terrestrial orchids and the structure of co-occurring vegetation means that seed falls close to the parent plant (Backhouse & Cameron 2005; Jersáková & Malinová 2007). Some species rely on streams for seed dispersal, and others may benefit through fauna movements (Weston *et al.* 2005), but this is rare.

#### 2.4.4.1 *Diuris tricolor*

54. As a genus, *Diuris* sp. replace their tubers annually and a small number are also capable of spreading vegetatively through daughter tuberoids, although this is rare (Jones 1993). Backhouse *et al.* (2016) hold a differing view, stating that most species can multiply vegetatively but several rely solely on seed dispersal. *Diuris tricolor* is most commonly encountered as individual plants or loose groups, consistent with spread through seed dispersal (pers. obs.).

#### 2.4.4.2 *Prasophyllum petilum*

55. With few exceptions, nearly all species of *Prasophyllum* replace their tubers annually and do not colonise adjacent habitat through daughter tuberoids. As a consequence, most species rely on seed production and dispersal within favoured habitat (Jones 1993), and most occur as scattered individuals or in loose groups. Within the Hunter Valley, *Prasophyllum petilum* follows this typical pattern in population growth habit (pers. obs.).

#### 2.4.4.3 *Pterostylis chaetophora*

56. In *Pterostylis* sp., some species are solitary while others form extensive colonies through daughter tuberoids on the ends of long stolons (Jones 1993; Backhouse *et al.* 2016). All species produce replacement tubers at the conclusion of the flowering season, emerging as leaves following good rainfall during Autumn. Members of subgenus *Oligochaetophilus* (including *Pterostylis chaetophora*) are rarely colonising species, generally replacing their tubers annually rather than spreading through daughter tuberoids (Juisaitis & Sorensen 1993). This implies that seed dispersal is the primary means of landscape dispersal for these species. Indeed, in my experience nearly all populations of *Pterostylis chaetophora* occur individually or in small, loose groups (Bell & Hillier 2020).



### 3. Criterion (b) – Justification for an Expert Report

57. A BDAR for the Project has been prepared by Hunter Eco (2019). This report details the extent of previous and new survey of vegetation, flora and fauna across the wider Maxwell lands including the Project Area. I have relied on this document to provide the most up-to-date information on the biodiversity of the locality, and used it to assist my assessment of the potential presence within the Project Area of the three target orchid species. However, I have used my own experience with these orchids elsewhere, and observations made during a one-day field inspection of the Project Area (detailed in **Section 4**), to inform my assessment.
58. During recent surveys completed for the Project, no threatened flora species were recorded by Hunter Eco (2019). This included sites where *Diuris tricolor* had previously been observed by other workers, but drought conditions would have impacted on flower emergence and persistence during recent years (see **Section 3.2**). Since initial preparation of this report, Hunter Eco (2020) completed additional surveys in and around the Project Area. *Diuris tricolor* was recorded near original records of the known population, however no threatened flora species were recorded within the Project Area.

#### 3.1 Survey Effort

59. Hunter Eco (2019) details (in Table 3 of Attachment A – *Baseline Flora Report*) the extent to which the Project Area and the immediately surrounding lands have been subject to flora and vegetation surveys, particularly in relation to the Mt Arthur Coal Mine and the former proposed Drayton South open cut mine (now the proposed Maxwell Project). Investigations shown there as occurring within the current Project Area include:
  - Hansen Bailey (2007)
  - Cumberland Ecology (2009)
  - Cumberland Ecology (2012)
  - Cumberland Ecology (2015)
60. In addition, flora surveys have been carried out across the Project Area and surrounds by Hunter Eco in 2017, 2018, 2019 and 2020, and other survey periods are noted in Cumberland Ecology (2015). [Table 6](#) summarises the timing of field survey for each of these investigations, relative to expected flowering periods (i.e. detectability) of the three target orchids. One of the target orchids (*Diuris tricolor*) was initially recorded in 2009 near to the Project Area by Cumberland Ecology (now contained within a fenced off area; referred to henceforth as the ‘*Diuris* Conservation Area’), where two individuals were located. Repeat surveys in the area in November 2010 did not locate any plants (not surprising as November falls outside of the flowering period), but 30 plants were recorded in ‘Spring’ (presumably September or October) 2011. Cumberland Ecology (2015) described the habitat at this location as Hunter Floodplain Red Gum Woodland, although on my inspection there on 3 July 2020 I saw woodland dominated only by *Eucalyptus conica* and *Eucalyptus moluccana*, with patches of *Angophora floribunda* and *Eucalyptus melliodora*.
61. From [Table 6](#) it is evident that few of these survey periods correspond with peak flowering times in the three target orchids. Cumberland Ecology expended three days of effort (incorporating vegetation mapping, habitat assessment, plot sampling and threatened flora searches) on 30 September to 2 October 2009, and one day (targeted orchid and *Acacia*

*pendula* surveys) on 23 September 2011 (Cumberland Ecology 2015). Hunter Eco undertook targeted orchid surveys over two days on 28 September and 17 October 2018, 10 days of floristic plot survey throughout September and October 2018, and targeted orchid surveys over two days on 28 and 29 September 2020.

**Table 6 Field survey effort (grey) and peak orchid flowering periods (black).** EC = Ecotone; HB = Hansen Bailey; CE = Cumberland Ecology; HE = Hunter Eco. Survey periods shown include both targeted searches and plot- or transect-based floristic surveys where it may be expected that orchids would be detected if present.


	J	F	M	A	M	J	J	A	S	O	N	D
EC 2000		█										
HB 2006		█							█			
HB 2007		█										
CE 2009					█					█	█	
CE 2010											█	
CE 2011			█			█		█	█			
CE 2013			█									
CE 2015		█										
HE 2017												█
HE 2018									█	█	█	█
HE 2019		█							█	█	█	█
HE 2020									█			
<i>D. tricolor</i>										█		
<i>P. petilum</i>										█		
<i>P.chaetophora</i>										█		

62. Throughout all orchid searches and floristic plot surveys, and from the data available to me, only two terrestrial orchids have been located across the Project Area or surrounds: *Diuris tricolor* by Cumberland Ecology in 2009 and 2011 and by Hunter Eco in 2020, and the common and unlisted *Pterostylis bicolor* by Hunter Eco in 2018 (Hunter Eco 2019), both recorded from the Diuris Conservation Area (C. Driscoll pers. comm.). A low diversity of terrestrial orchids is not unusual for this part of the Hunter Valley; surveys by Kleinfelder across the Bayswater Power Station area immediately adjacent to the east have reported only a *Microtis* sp. across 71 floristic plots (although mostly under drought conditions, discussed in Bell 2020c). Further discussion on the importance of this lack of terrestrial orchids is presented in **Section 4.3**.


### 3.2 Impact of Drought

63. As noted earlier, terrestrial orchids are sensitive to environmental conditions, and in the Hunter Valley prevailing rainfall patterns are influential in their detection during surveys. The past three years in the central Hunter Valley have been particularly dry. [Figure 9](#) shows that, apart from the very wet months of March 2017, March 2019, and September 2019, below average rainfall has been received at Muswellbrook (c. 10 km to the north) from 2017. Such dry conditions will place all plants under severe water stress, and for emergent geophytes like Spring-flowering orchids, dry Autumn-Winter periods over successive years commonly result in little or no emergence, and increased grazing pressure.

64. The NSW Department of Primary Industries (DPI) Combined Drought Indicator (CDI) provides a further dataset that can help to explain how orchids and other plants respond to climatic conditions. The CDI categorises months into one of six phases of drought on the basis of three indices (rainfall index, plant growth index, soil water index). The Wynn Parish (which includes the Project Area) formally experienced high levels of drought over most of January 2015 to March 2020 (Figure 10). Intense Drought was experienced between April 2018 and October 2018 (encompassing the 2018 orchid flowering period), and again from December 2018 to December 2019. The 2019 orchid flowering season (September-October) and the months leading up to it were also drought affected. Clearly, any targeted orchid surveys undertaken during the 2015 to 2019 flowering seasons would be unlikely to detect representative populations of these species if present.



**Figure 9** Monthly and average rainfall for Muswellbrook (station # 61374; c. 25 km from the Project Area) over the past three years. (Source: Bureau of Meteorology, <http://www.bom.gov.au/climate/data/>).



**Figure 10** Drought indices for the Wynn Parish, January 2013 to July 2020, showing Drought conditions prevailing from October 2014 to July 2017 and December 2018 to December 2019, and Intense Drought from April 2018 to December 2018 and December 2019 to January 2020. (Source: DPI Seasonal Conditions Information Portal, <https://edis.dpi.nsw.gov.au/statistics>).

65. Most terrestrial orchid species will not emerge to flower during stressful periods, or if leaves are produced at this time then inflorescences may not develop. Given the drought conditions experienced from October 2014 to the present, and in particular during the Autumn and Winter periods leading up to and including September-October orchid flowering from 2015 to 2019, there is clear justification for the preparation of this expert report. Apart from water stress, pressure from herbivory during drought periods escalates considerably (Duncan *et al.* 2005), not only from vertebrate grazers such as macropods and rabbits, but also invertebrates including grasshoppers and caterpillars (Light & MacConnail 2011; Vizer 2013). Bird species too are known to selectively feed on orchid species, with White-winged Choughs for example extracting orchids out of the ground to consume tubers (Duncan *et al.* 2005; Faast & Facelli 2009; Bell 2020b). Any vegetation present during dry times will be the focus of herbivore browsing, meaning a reduction in the time orchids will be present above ground and hence reduce detection rates during survey. Desiccation through heat and wind in periods of drought will also lessen above-ground periods of flowering orchids.
66. Persistent dry conditions and drought, including below-average rainfall during the crucial Winter period, over at least the last three flowering seasons (2017 to 2019) justify the need for an Expert Report to determine the likely presence of *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis chaetophora* within the Project Area.

## 4. Criterion (c) – Likelihood of Species Presence in the Project Area

67. To assist in determining the likely presence or absence of populations of *Diuris tricolor*, *Prasophyllum petilum* or *Pterostylis chaetophora* within the Project Area, I have examined aspects of previous land-use history, known existing records for all three species within the region, the floristic composition, geology, soil landscapes and soil qualities of habitats within the Project Area and other areas known to support the species. I have also incorporated my own observations made on a single-day site inspection in early July 2020.

### 4.1 Project Area Attributes

#### 4.1.1 Land-use History

68. As with much of Australia and NSW, the original ecosystems of Hunter Valley have been heavily impacted upon by European colonisation (Bradshaw 2012). The mid-Hunter Valley was one of the first to be opened up for European occupation, spanning out from the nearby settlement of Jerrys Plains in the early 1800s (Burley 1962; C. Hunter 2010). Much of the land was used for the grazing of cattle and sheep, and where necessary the original woodlands were thinned of trees to increase pasture growth (Perry 1955). As a consequence, current day vegetation is the result of 200 years of agricultural occupation, firstly by sheep and then cattle. The Project Area is now almost entirely comprised of derived grasslands, regrowth native and exotic woody vegetation, or planted species (Hunter Eco 2019).
69. Although cattle have historically grazed the Project Area for likely several decades (Hunter Eco 2019), they have been removed or rotated to other areas during recent drought conditions. It is unknown whether or not cattle (or sheep) were present across the Project Area during flora survey times indicated in Table 6 above. All of the derived grasslands which now predominate throughout the Project Area potentially provide potential habitat for orchids including *Diuris tricolor* and *Prasophyllum petilum*, while remnant woodlands may support *Diuris tricolor* or *Pterostylis chaetophora*, but this is further explored in the following sections.

#### 4.1.2 Existing Orchid Records

70. No populations of *Diuris tricolor*, *Prasophyllum petilum* or *Pterostylis chaetophora* have been recorded within the Project Area (Hunter Eco 2019), although one population of *Diuris tricolor* (c. 30 plants) located by Cumberland Ecology in 2009 and 2011 (and again by Hunter Eco in 2020) lies approximately 600 m from the disturbance footprint (south of the proposed Ancillary Disturbance Area 1), and another (of nearly 200 plants) occurs 400 m north of the Product Stockpile Extension area (NSW BioNet). A historical record of *Prasophyllum petilum* (c. 10 plants) occurs for a site along Thomas Mitchell Drive c. 1300 m east of the Product Stockpile Extension area. As noted in Section 3.2, drought conditions coincided with the preparation of documents relating to the proposed development in the area, and as a consequence there is a low likelihood that these species, if present, would have been detected.
71. Evidence of survey effort is shown in Figure 9 (Attachment A) of Hunter Eco (2019), where the extent of data collection made across 1708 Rapid Data Points (RDPs) (used for mapping purposes) and 109 full floristic plots (used for classification) is illustrated across the wider biodiversity survey area. Some RDP and 23 full floristic plots fall within the Project Area, and a portion of floristic plots were likely surveyed in September-October 2018. However, the Intense

Drought occurring at that time (see **Section 3.2**) would have curtailed detection of any terrestrial orchids that may have been present. Examination of surrounding records of the three target orchids may shed some light on the expected presence of *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis chaetophora* within the Project Area (further examined in **Section 4.2**).

#### 4.1.3 Vegetation Communities

72. Understanding the floristic patterns and vegetation communities in the Project Area is important in gaining an impression of how suitable the lands are to support one or more of the target orchid species. Fortunately, Hunter Eco (2019) has undertaken a numerical analysis of floristic plot data (n=109) to assist in classifying the native vegetation present, and provides detailed community profiles in Appendix 4 of the *Baseline Flora Report*. I have reviewed this classification and the profiles, and am satisfied that it represents a thorough treatise of vegetation community diversity present within the Project Area. This classification, together with other factors and my own notes made during a one-day site inspection (see **Section 4.3**), have been used to formulate an opinion on the likelihood of *Diuris tricolor*, *Prasophyllum petilum* and/or *Pterostylis chaetophora* being present within the Project Area (presented in **Section 4.5**).

##### 4.1.3.1 Plant Community Types

73. Across the broader biodiversity investigation area, Hunter Eco (2019) defined 11 Plant Community Types (PCTs) from their 18 draft field communities, all of which also occur elsewhere in the Hunter Valley. Thirteen (13) communities and eight PCTs (201, 1604, 1606, 1607, 1655, 1691, 1692 and 1731) have been mapped within the Project Area (**Table 7**). DNGs, which provide substantial habitat for *Diuris tricolor* and *Prasophyllum petilum* in other areas, are represented here by five PCTs (201, 1606, 1607, 1655 and 1691). Collectively, 136 ha of DNG (or 80% of all communities shown in **Table 7**) are present within the Project Area. Approximately three-quarters of the total proposed disturbance area is comprised of native grasslands derived from White Box – Ironbark – Red Gum Shrubby Forest (Unit 2a), followed by Ironbark - Grey Box Grassy Woodland (Unit 9; 5.7%) and White Box - Ironbark - Red Gum Shrubby Forest (Unit 2; 5.6%). All other communities each occupy <3% of the total Project Area.
74. Examining the floristic composition and condition of defined vegetation communities presented in Hunter Eco (2019) reveals the characteristic species which dominate each community. These data can then be used to compare against similar data from other areas of the Hunter Valley where *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis chaetophora* are known to occur (see **Section 4.2**). In this regard, **Table 8** summarises the key floristic elements of the 13 vegetation communities within the Project Area. To compile this table, I extracted those plant species contributing to the upper 50% of diversity, as shown in the community profiles in Attachment A of Hunter Eco (2019).
75. For the most widespread community within the Project Area (White Box – Ironbark – Red Gum Shrubby Forest DNG; Unit 2a; 74% of all communities shown in **Table 7**), grasslands dominated by *Aristida ramosa*, *Rytidosperma caespitosum*, *Eragrostis alveiformis*, *Dichanthium sericeum*, *Eriochloa pseudoacrotricha*, *Bothriochloa decipiens*, *Erodium crinitum* and *Chloris divaricata*, with high abundance of exotic species (particularly *Carthamus lanatus* and *Senecio madagascariensis*), characterises these areas (Hunter Eco 2019). In my experience, these relatively diverse grasslands growing on clay-based soils derived from former Box-Gum

landscapes tend not to support extensive populations of *Diuris tricolor*, *Prasophyllum petilum* or *Pterostylis chaetophora* (or few other terrestrial orchids), however I will reserve this view and further analyse available floristic and environmental data.

**Table 7 Vegetation Communities and Plant Community Types identified for the Project Area (from Hunter Eco 2019).** Excludes post-mine mapping. Condition classes within PCTs not shown.

Community	PCT	Extent (ha, %)
1: Red Gum - Ironbark - Apple Shrubby Woodland	1607: Blakely's Red Gum – Narrow-leaved Ironbark – Rough-barked Apple shrubby woodland of the upper Hunter	0.4 (0.2%)
1a: Red Gum - Ironbark - Apple Shrubby Woodland (DNG)	1607: Blakely's Red Gum - Narrow-leaved Ironbark - Rough-barked Apple shrubby woodland of the upper Hunter - DNG	4.9 (2.9%)
2: White Box - Ironbark - Red Gum Shrubby Forest	1606: White Box -Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter	9.6 (5.6%)
2a: White Box - Ironbark - Red Gum Shrubby Forest (DNG)	1606: White Box -Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter - DNG	125.6 (73.6%)
3: Slaty Box Shrubby Woodland	1655: Grey Box – Slaty Box shrub – grass woodland on sandstone slopes of the upper Hunter Valley and Sydney Basin	1.2 (0.7%)
3a: Slaty Box Shrubby Woodland (DNG)	1655: Grey Box – Slaty Box shrub – grass woodland on sandstone slopes of the upper Hunter Valley and Sydney Basin - DNG	2.4 (1.4%)
4: Swamp Oak Forest	1731: Swamp Oak – Weeping Grass grassy riparian forest of the Hunter Valley	0.2 (0.1%)
6: Bull Oak Grassy Woodland	1692: Bull Oak grassy woodland of the central Hunter Valley	2.7 (1.6%)
8. Fuzzy Box Woodland	201: Fuzzy Box Woodland on alluvial brown loam soils mainly in the NSW South Western Slopes Bioregion	0.5 (0.3%)
8a: Fuzzy Box Woodland (DNG)	201: Fuzzy Box Woodland on alluvial brown loam soils mainly in the NSW South Western Slopes Bioregion - DNG	2.8 (1.6%)
9: Ironbark - Grey Box Grassy Woodland	1691: Narrow-leaved Ironbark - Grey Box grassy woodland of the central and upper Hunter	9.6 (5.7%)
9a: Ironbark - Grey Box Grassy Woodland (DNG)	1691: Narrow-leaved Ironbark - Grey Box grassy woodland of the central and upper Hunter - DNG	0.3 (0.2%)
11: Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland	1604: Narrow-leaved Ironbark - Grey Box - Spotted Gum shrub - grass woodland of the central and lower Hunter	1.3 (0.8%)
Plantation, Dams etc	n/a	9.1 (5.3%)
<b>Total</b>		<b>170.6 (100%)</b>

#### 4.1.3.2 Comparative Numerical Analyses

76. To further explore floristic compositions between the Project Area and known orchid habitat elsewhere (Mangoola Coal for *Diuris tricolor* and *Prasophyllum petilum*, and Columbey National Park for *Pterostylis chaetophora*), I undertook numerical analysis using *PRIMER* software (Clarke & Gorley 2006). Both Mangoola and Columbey have had considerable directed survey undertaken for the target orchids, and both have also been the subject of numerical analysis using the same software and procedures as used by Hunter Eco (2019) to classify the vegetation within the Maxwell biodiversity investigation area. It is logical, therefore, to make use of this knowledge base to inform decisions on the suitability of habitat for the three target species.
77. Rather than using raw data for these analyses (which utilised different survey methods: 0.04 ha plots for the Project Area and Columbey but 0.01 ha plots at Mangoola, and surveyed during different seasons), I compared floristic compositions of the available community profiles derived for the Project Area with those profiles shown to support one or more of the target orchids. This included Units 1, 1a, 2, 2a, 3, 3a, 4, 6, 8, 9, 9a and 11 for the Project Area (see [Table 8](#)), Units 1a and 2 for Mangoola Coal (Bell submitted 1), and Units 4, 6, 7 and 8 for Columbey (Bell 2009) ([Table 9](#)).
78. All data was analysed on a presence-absence basis, meaning that dominant species held equal weight to rare species. Reducing data from cover abundance to presence-absence also lessens the impact of comparing surveys across different years and observers, where species commonly abundant during wet years will persist only in low numbers during drought. While not ideal, this process allowed me to determine if flora diversity (rather than abundance) present in the Project Area (collected during the drought year of 2018) correlated well with that for Mangoola where *Diuris tricolor* and *Prasophyllum petilum* were known to occur (collected between 2009 and 2011, c. average rainfall years), and with Columbey where *Pterostylis chaetophora* is present (collected in 2008 and 2009, above average rainfall years). Analyses with weed species retained and removed from the dataset were undertaken, but this made little difference to the overall result.
79. Because habitats of the three target orchids at the specified locations differ (predominantly DNGs for *Diuris tricolor* and *Prasophyllum petilum*, forest or woodland for *Pterostylis chaetophora*), I created sub-sets of data to reduce 'noise' in the analyses. This resulted in two separate analyses as shown in [Table 10](#). Note that Unit 8a (Fuzzy Box Woodland DNG) for the Project Area was not included as insufficient data prevented development of a floristic list by Hunter Eco (2019). Unit 4 (Swamp Oak Forest) from the Project Area was similarly excluded due to low diversity (six species), and based on past experience the very low likelihood of this habitat supporting terrestrial orchids.

For both analyses, I used the SIMPROF routine in combination with the CLUSTER module in *PRIMER* to identify statistically significant splits in the dataset ( $p < 0.01$ ). This provided cluster diagrams where sites supporting similar floristic combinations were grouped and linked to their most similar neighbours. I also ran the MDS (non-metric Multi-Dimensional Scaling) routine with a minimum stress level of 0.01 and 25 restarts to produce ordination plots of the same data. Grouping of similar sample plots (communities or vegetation zones) can be better appreciated across this two-dimensional ordination space than in a cluster diagram, hence the latter is not shown. Note that for the Mangoola and Columbey datasets, my original analyses truncated SIMPER routines at 90% contributions, but these were re-run to 95% to match that done by Hunter Eco (2019) for the Project Area.



**Table 8 Dominant species within defined vegetation communities (from Hunter Eco 2019).** \* = exotic species. Individual % contributions shown only for species contributing >5% of total diversity for that community.

Community	Dominant Species (upper 50%, in decreasing order of importance)
1: Red Gum - Ironbark - Apple Shrubby Woodland	<i>Eucalyptus blakelyi</i> (25.82), <i>Notelaea microcarpa</i> (10.47), * <i>Galenia pubescens</i> , <i>Brachychiton populneus</i> , * <i>Lycium ferocissimum</i>
1a: Red Gum - Ironbark - Apple Shrubby Woodland (DNG)	<i>Aristida ramosa</i> (15.17), * <i>Sida rhombifolia</i> (11.05), * <i>Galenia pubescens</i> (11.05), * <i>Senecio madagascariensis</i> (5.52), * <i>Petrohragia nanteuilii</i> , <i>Commelina cyanea</i>
2: White Box - Ironbark - Red Gum Shrubby Forest	<i>Eucalyptus albens</i> (25.75), <i>Dichondra repens</i> (5.46), <i>Glycine clandestina</i> (5.25), <i>Brunoniella australis</i> , <i>Acacia salicina</i> , <i>Eremophila debilis</i>
2a: White Box - Ironbark - Red Gum Shrubby Forest (DNG)	* <i>Carthamus lanatus</i> (10.73), <i>Aristida ramosa</i> (6.39), <i>Rytidosperma caespitosum</i> (6.07), * <i>Senecio madagascariensis</i> (5.48), <i>Glycine clandestina</i> , <i>Eragrostis alveiformis</i> , <i>Dichanthium sericeum</i> , <i>Maireana microphylla</i> , <i>Erodium crinitum</i> , <i>Chloris divaricata</i>
3: Slaty Box Shrubby Woodland	<i>Eucalyptus dawsonii</i> (44.84), <i>Aristida ramosa</i> (5.63)
3a: Slaty Box Shrubby Woodland (DNG)	<i>Chloris divaricata</i> (11.96), <i>Dichanthium sericeum</i> (10.31), * <i>Carthamus lanatus</i> (6.99), * <i>Medicago sp.</i> (5.55), <i>Rytidosperma caespitosum</i> , <i>Vittadinia pterochaeta</i> , * <i>Lepidium bonariense</i> , <i>Glycine clandestina</i>
4: Swamp Oak Forest	<i>Casuarina glauca</i> (54.55)
6: Bull Oak Grassy Woodland	<i>Allocasuarina luehmannii</i> (60.10)
8: Fuzzy Box Woodland	<i>Eucalyptus conica</i> (26.67), <i>Aristida ramosa</i> (24.44)
8a: Fuzzy Box Woodland (DNG)	no analysis of data
9: Ironbark - Grey Box Grassy Woodland	<i>Eucalyptus moluccana</i> (28.58), <i>Eremophila debilis</i> (6.09), <i>Brunoniella australis</i> (5.98), <i>Allocasuarina luehmannii</i> (5.95), <i>Dichondra repens</i>
9a: Ironbark - Grey Box Grassy Woodland (DNG)	<i>Aristida ramosa</i> (16.84), <i>Chrysocephalum semipapposum</i> (5.30), <i>Cymbopogon refractus</i> (5.10), <i>Brunoniella australis</i> , <i>Linum marginale</i> , <i>Wahlenbergia communis</i> , <i>Eragrostis leptostachya</i> , * <i>Gomphocarpus fruticosus</i> , * <i>Carthamus lanatus</i>
11: Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland	<i>Aristida ramosa</i> (14.27), <i>Lomandra filiformis</i> (9.39), <i>Eucalyptus moluccana</i> (6.41), <i>Eremophila debilis</i> (5.67), <i>Brunoniella australis</i> (5.67), <i>Lomandra multiflora</i> (5.67), <i>Glycine clandestina</i> (5.67)

**Table 9 Vegetation communities previously defined for Mangoola (Bell submitted 1) and Columbey (Bell 2009) shown to support populations of one or more of the target orchid species.**

Location	Unit		Target Orchid
Mangoola	1a	Dichanthium/ Sporobolus/ Chloris Grassland	<i>D. tricolor</i> , <i>P. petilum</i>
	2	Aristida/ Cymbopogon Grassland	<i>D. tricolor</i> , <i>P. petilum</i>
Columbey	4	Floodplain Redgum – Box Forest	<i>P. chaetophora</i>
	6	Seaham Ironbark Forest	<i>P. chaetophora</i>
	7	Seaham Spotted Gum – Ironbark Forest	<i>P. chaetophora</i>
	8	Lower Hunter Spotted Gum – Ironbark Forest	<i>P. chaetophora</i>


**Table 10 Included datasets in numerical analyses.**

Analysis No.	Habitat	Target Orchids	Dataset
1	Derived Native Grasslands	<i>Diuris tricolor</i> , <i>Prasophyllum petilum</i>	Project Area (Units 1a, 2a, 3a, 9a) Mangoola (Units 1a, 2)
2	Forests & Woodlands	<i>Pterostylis chaetophora</i>	Project Area (Units 2, 3, 6, 9, 11) Columbey (Units 4, 6, 7, 8)


80. For Analysis 1 (Derived Native Grasslands), three significant groups were evident in the dataset, with Unit 9a from the Project Area closely aligning with Units 1a and 2 from Mangoola (Figure 11). The other three DNG types fell distant from Mangoola data and evidently support different floristic compositions. The stress level of 0 shown in Figure 11 is an indication of the relative ease in which all data could be accommodated within two-dimensions. In general, a stress level of <0.2 is considered acceptable in these sorts of analyses, but increases in line with complexities associated with dataset size, multiple observers and seasons. From this grassland analysis, only Unit 9a (Ironbark - Grey Box Grassy Woodland DNG) from the Project Area could be considered as providing potential habitat for *Diuris tricolor* and *Prasophyllum petilum*.
81. For Analysis 2 (Forests & Woodlands), it was clearly evident that the forest and woodland vegetation present within the Project Area is floristically different to that known to support *Pterostylis chaetophora* at Columbey (Figure 12). This is not surprising given the wide geographical separation between these two localities (c. 100 km), however both areas support vegetation characterised by *Eucalyptus moluccana*, *Eucalyptus crebra* and *Corymbia maculata*, and provide superficially appropriate habitat. Based on this dataset, it is unlikely that any habitat within the Project Area is suitable for *Pterostylis chaetophora*.
82. One conclusion to reach from these analyses is that the floristic composition of nearly all vegetation communities defined for the Project Area differ sufficiently from the DNGs at Mangoola known to support populations of both *Diuris tricolor* and *Prasophyllum petilum*, and also from the forest and woodlands at Columbey known to support *Pterostylis chaetophora*. The one exception to this is the close alignment of Unit 9a (Ironbark - Grey Box Grassy Woodland DNG) from the Project Area with the Mangoola grasslands, suggestive of potential habitat for *Diuris tricolor* and *Prasophyllum petilum*. Acknowledging the different years of data collection between these two locations (2009-11 vs 2018), the different observers collecting

the data and the different methods of data collection (0.04 ha plots vs 0.01 ha plots), I am confident that within these datasets there are sufficient differences between all other grasslands and grassy woodlands present in the Project Area and those at Mangoola Coal. This is supported in part by my own field observations of the Project Area in July 2020 (see **Section 4.5**).

83. Notwithstanding this result, the presence of a known population of c. 200 *Diuris tricolor* occurring within Mt Arthur Coal lands in Grey Box – Spotted Gum – Narrow-leaved Ironbark Woodland (Unit 11 of Hunter Eco 2019) approximately 400 m north of the Product Stockpile Extension area, suggests that this species may also occur in this vegetation type within that Project Area. I do not have access to any floristic data from this area to allow more in-depth comparisons, but given its close proximity it is prudent to assume suitable habitat occurs there.



**Figure 11** nMDS ordination of floristic compositions comprising Derived Native Grasslands from the Project Area (MX) in relation to that favouring *Diuris tricolor* and *Prasophyllum petilum* at Mangoola Coal (MG). Dotted ellipses show significant groups ( $p < 0.01$ ) defined at a similarity level of 40%. MX\_1a = Red Gum - Ironbark - Apple Shrubby Woodland (DNG); MX\_2a = White Box - Ironbark - Red Gum Shrubby Forest DNG; MX\_3a = Slaty Box Shrubby Woodland DNG; MX\_9a = Ironbark - Grey Box Grassy Woodland DNG; MG\_1a = Dichanthium/ Sporobolus/ Chloris Grassland; MG\_2 = Aristida/ Cymbopogon Grassland. Data for the Project Area from Hunter Eco (2019), that from Mangoola from Bell (submitted 1).



**Figure 12** nMDS ordination of floristic compositions comprising forest and woodlands from the Project Area in relation to favoured *Pterostylis chaetophora* habitat at Columbey. Dotted ellipses show significant groups ( $p < 0.01$ ) defined at a similarity level of 40%. Individual community codes not shown, as these are irrelevant to the outcome. Data for the Project Area from Hunter Eco (2019), that from Columbey from Bell (2009).

84. Additionally, following the same reasoning, the historical population of *Prasophyllum petilum* (recorded 1999-2005; c. 10 plants) at the eastern end of Thomas Mitchell Drive suggests a conservative assumption that the Product Stockpile Extension area may also support this species. Data contained within NSW BioNet linked to this population show it to be woodland of *Corymbia maculata*, *Eucalyptus moluccana* and *Eucalyptus blakelyi*, similar to that which occurs within the Product Stockpile Extension area (pers. obs.).

## 4.2 Environmental Analysis

85. An assessment of selected environmental attributes (geology, dominant lithology, soil landscape, soil type, soil fertility, soil hydrology, annual rainfall, mean annual temperature) was made of the Project Area with the aim of profiling the broad non-floristic characteristics that may be important in defining habitat for the three target orchids. A number of resources were used to undertake this (Table 11), all performed in GIS against Project Area boundaries. Other commonly used modelling parameters (e.g. slope, aspect, solar radiation, roughness indices, moisture etc) were not explored in any detail, although they have reportedly proven informative for some species (e.g. Janes 2010). However, these attributes are unlikely to be instructive in the open and gently undulating landscapes of the Hunter Valley floor, and in addition they may lead assessment away from the central aim of identifying habitat for orchids and their mycorrhizal fungi. Understanding the biophysical properties of the Project Area is important, though, to allow comparisons with point data from other areas where *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis chaetophora* occurs (see Section 4.2.2).

**Table 11 Digital resources utilised to categorise biophysical attributes of the Project Area, and compare these with point data for the three target orchids.**

Attribute	Source
Geological Unit	Colquhoun G.P., Hughes K.S., Deyssing L., Ballard J.C., Folkes C.B, Phillips G., Troedson A.L. & Fitzherbert J.A. (2020) <i>New South Wales Seamless Geology dataset, version 2</i> [Digital Dataset]. Geological Survey of New South Wales, Department of Regional NSW, Maitland. Available at <a href="https://search.geoscience.nsw.gov.au/product/9232">https://search.geoscience.nsw.gov.au/product/9232</a>
Dominant Lithology	Colquhoun G.P., Hughes K.S., Deyssing L., Ballard J.C., Folkes C.B, Phillips G., Troedson A.L. & Fitzherbert J.A. (2020) <i>New South Wales Seamless Geology dataset, version 2</i> [Digital Dataset]. Geological Survey of New South Wales, Department of Regional NSW, Maitland. Available at <a href="https://search.geoscience.nsw.gov.au/product/9232">https://search.geoscience.nsw.gov.au/product/9232</a>
Soil Landscapes	Office of Environment and Heritage (2019) <i>Soil Landscapes of Central and Eastern NSW - v2</i> , NSW Office of Environment and Heritage, Sydney. Available at <a href="https://datasets.seed.nsw.gov.au/dataset/published-soil-landscapes-of-central-and-eastern-nsw37d37">https://datasets.seed.nsw.gov.au/dataset/published-soil-landscapes-of-central-and-eastern-nsw37d37</a>
Great Soil Group	Office of Environment and Heritage (2017) <i>Great Soil Group (GSG) Soil Type map of NSW</i> , NSW Office of Environment and Heritage, Sydney. Available at <a href="https://datasets.seed.nsw.gov.au/dataset/great-soil-group-gsg-soil-type-map-of-nsw1cf19">https://datasets.seed.nsw.gov.au/dataset/great-soil-group-gsg-soil-type-map-of-nsw1cf19</a>
Australian Soil Class	Office of Environment and Heritage (2017) <i>Australian Soil Classification (ASC) Soil Type map of NSW</i> , NSW Office of Environment and Heritage, Sydney. Available at <a href="https://datasets.seed.nsw.gov.au/dataset/australian-soil-classification-asc-soil-type-map-of-nsw10">https://datasets.seed.nsw.gov.au/dataset/australian-soil-classification-asc-soil-type-map-of-nsw10</a>
Soil Fertility	Office of Environment and Heritage (2017) <i>Estimated Inherent Soil Fertility of NSW</i> , NSW Office of Environment and Heritage, Sydney. Available at <a href="https://datasets.seed.nsw.gov.au/dataset/estimated-inherent-soil-fertility-of-nsw793e">https://datasets.seed.nsw.gov.au/dataset/estimated-inherent-soil-fertility-of-nsw793e</a>
Hydrologic Soils	Office of Environment and Heritage (2017) <i>Hydrologic Groups of Soils in NSW</i> , NSW Office of Environment and Heritage, Sydney. Available at <a href="https://datasets.seed.nsw.gov.au/dataset/hydrologic-groups-of-soils-in-nsw7f9e8">https://datasets.seed.nsw.gov.au/dataset/hydrologic-groups-of-soils-in-nsw7f9e8</a>
Annual Rainfall	GeoScience Australia, raster layer based on monthly mean precipitation 1976-2005, with grid resolution of 30m. Available at: <a href="https://datasets.seed.nsw.gov.au/dataset/anuclim-annual-mean-rainfall-raster-layer">https://datasets.seed.nsw.gov.au/dataset/anuclim-annual-mean-rainfall-raster-layer</a>
Annual Average Temperature	GeoScience Australia, raster layer based on monthly mean temperatures 1976-2005, with grid resolution of 30m. Available at: <a href="https://datasets.seed.nsw.gov.au/dataset/anuclim-annual-mean-temperature-raster-layer">https://datasets.seed.nsw.gov.au/dataset/anuclim-annual-mean-temperature-raster-layer</a>

**Table 12 Summary of biophysical attributes of the Project Area.** See [Appendix 3](#) for further detail.

Project Area	Biophysical Summary
Product Stockpile Extension (5.2 ha)	Part Cessnock Sandstone and part Branxton Formation geology; equally of sandstone and conglomerate lithology; entirely of the Roxburgh soil landscape, supporting Yellow Podzolic (Kurosol) soils of Moderately Low fertility with slow infiltration rates. Annual rainfall is 671 mm (StDev=0.67, n=4) and mean temperature is 16.6° (StDev=0.01, n=4).
Surface Development Area (302.9 ha)	Predominantly of Jerrys Plains Subgroup and Rowan Formation geology, with Mulbring Siltstone and Branxton Formation also well represented; mostly of sandstone lithology, but with siltstone, conglomerate and quartzite also present; largely of the Brays Hill and Liddell soil landscapes, supporting Grey, Brown or Red Clay (Vertosol) soils and Soloths (Kurosols), of Moderate or Moderately Low fertility with very slow infiltration rates. Annual rainfall is 658 mm (StDev=13.4, n=20) and mean temperature is 16.7° (StDev=0.16, n=20).
Edderton Rd Realignment (10.2 ha)	Entirely Jerrys Plains Subgroup geology; of sandstone lithology; on the Bayswater and Brays Hill soil landscapes, supporting Solodic (Sodosols) and Grey, Brown or Red Clay (Vertosol) soils, of Moderately Low and Moderate fertility with very slow infiltration rates. Annual rainfall is 623 mm (StDev=6.64, n=11) and mean temperature is 17.1° (StDev=0.10, n=11).
Ancillary Disturbance Area (Ponding) 1 (0.5 ha)	Entirely Jerrys Plains Subgroup geology; of sandstone lithology; on the Brays Hill soil landscape, supporting Grey, Brown or Red Clay (Vertosol) soils, of Moderate fertility with very slow infiltration rates. Annual rainfall is 628 mm (StDev=0.78, n=2) and mean temperature is 17.1° (StDev=0.01, n=2).
Ancillary Disturbance Area (Ponding) 2 (1.5 ha)	Entirely Jerrys Plains Subgroup geology; of sandstone lithology; on the Bayswater soil landscape, supporting Solodic (Sodosol) soils, of Moderately Low fertility with very slow infiltration rates. Annual rainfall is 619 mm (StDev=0.01, n=2) and mean temperature is 17.2° (StDev=0.01, n=2).

86. The *NSW Flora: Ecological Niche Finder* online resource portal, arising out of the work of Gallagher (2016) and accessible at (<http://nswnichefinder.net/index.php>), was also considered as a possible tool to better understand the ecological requirements of the target species. The Niche Finder provides broad modelling capabilities for NSW plant taxa across a number of environmental domains, however because the target orchids are classified as sensitive species (location data withheld) this resource was not available.

#### 4.2.1 Project Area

87. [Table 12](#) summarises the environmental characteristics of the Project Area, categorised into the five separate disturbance areas (see also [Appendix 3](#) for graphed details). Nine geological units comprise the Project Area, however sediments of the Jerrys Plain Subgroup are the most common (46%). The Surface Development Area is the most complex area with seven units, the Product Stockpile Extension area has two units, and all others one. Soils derived from sandstone

lithology are dominant (>70%) across most areas, with siltstones and conglomerate present in the Surface Development Area and Product Stockpile Extension area, and a small area of quartzite. Four soil landscapes have been mapped for the area, the two most dominant being Brays Hill and Liddell. These mirror the distribution of the Great Soil Groups, where Grey, Brown and Red Clays dominate over Soloths, Solodic Soils and Yellow Podzolics. Under the Australian Soil Classification scheme, three soil types (Kurosols, Vertosols and Sodosols) occur across the area, and are of either moderately low or moderate fertility.

88. In terms of water infiltration (hydrology), most soils comprise Group D (very slow infiltration rates due to high clay content, high water tables and/or shallow soils over impervious material), although the Product Stockpile Extension area is Group C (slow infiltration rates due to impeded drainage and/or moderately fine textures). For climatic data, average values were computed from GeoScience Australia raster layers at 39 field inspection point locations across the Project Area (see **Section 4.4**), distributed as: Product Stockpile Extension area, n=4; Surface Development Area, n=20; Edderton Rd Realignment, n=11; Ancillary Disturbance Area (Ponding) 1, n=2; Ancillary Disturbance Area (Ponding) 2, n=2. Annual rainfall was consequently found to lie between 619 and 671 mm/year across the Project Area, while mean temperature was around 17°.


#### 4.2.2 Target Orchids Records

89. A compilation of all records for the three target orchids within the Hunter Local Land Services (Hunter LLS) area was interrogated against a range of environmental variables within GIS to determine ecological niches. This process aimed to identify, based on available records, the key environmental factors present within known populations of each species. Data included:
  - as-held NSW BioNet records of target orchids obtained under DPIE licence ASH20009;
  - currently undatabased but geospatially accurate records from various projects, including recent surveys for *Pterostylis chaetophora* collected under the NSW Government's *Saving our Species* program; and
  - other miscellaneous records gleaned from unpublished reports or databases.

Specimen collection data from the AVH was not incorporated as these data have been de-natured and therefore location precision is unreliable.

90. All data was checked for duplication across multiple data sources and positional accuracy, with records >100 m accuracy discarded from analysis. To reduce the influence of high-density point records resulting from targeted surveys over comparatively small search areas, the entire Hunter LLS region was overlain by a 100 m x 100 m grid (i.e. 1 ha squares) and orchid data reduced to presence-absence at this resolution. Centroids of all cells returning a 'true' result were adopted as surrogates for position and used to analyse environmental niches. Duplicates resulting from multiple high-density records within each 100 m grid were reduced to a single record.
91. Cleaning and simplifying orchid location data in this way resulted in an analysis dataset of 983 point locations (centroids) for *Diuris tricolor*, 485 point locations (centroids) for *Prasophyllum petilum* and 128 point locations (centroids) for *Pterostylis chaetophora*. These datasets were individually assessed against those environmental variables shown in [Table 11](#) above. The relative proportion of known orchid occurrences compared to representative locations within the Project Area (n=39) were graphed to illustrate similarities and differences (see [Appendix 4](#)), and a summary is provided in [Figure 13](#). For *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis*

*chaetophora*, there was little correlation between geological unit and soil landscape at locations known to support these species when compared to the Project Area. Additionally, there was also a low correlation in the Australian soil class and annual rainfall for *Pterostylis chaetophora*, and a weak match for annual temperature in this species. Overall, these results suggest there to be little apparent differences in soil type, fertility and hydrology for *Diuris tricolor* and *Prasophyllum petilum*, but some differences for *Pterostylis chaetophora*. It also implies that the Project Area may be too dry (mean 1005 mm/yr vs 640 mm/yr) and perhaps slightly too cool (17.2° vs 16.9°) for *Pterostylis chaetophora*, although this is only minor. The Project Area therefore potentially provides suitable soils and climate for *Diuris tricolor* and *Prasophyllum petilum*, but contingent on the availability of suitable mycorrhizal fungi in the soil and pollinators for ongoing persistence.



**Figure 13 Summary of percentage commonality of selected environmental attributes between known target orchid locations (n=983 *Diuris tricolor*; n=485 *Prasophyllum petilum*; n=128 *Pterostylis chaetophora*) and the Project Area (n=39).**

92. The markedly different soil landscapes are perhaps the most informative environmental attribute that might inform suitable habitat for orchids. By definition, soil landscapes attempt to combine elements of soil structure, chemistry and composition with features of the landscapes in which they were formed. This means that related variables such as vegetation, hydrology and topography (which all influence the biology of a soil) are captured within a single map unit type. For orchids this may be particularly telling, as the environments in which the necessary soil mycorrhiza form and persist are likely to be similar within the same soil landscapes, although there is no firm evidence for this to date. Following this logic, the fact that only 5% of 983 locations of *Diuris tricolor* (and <1% of 485 *Prasophyllum petilum* and 128 *Pterostylis chaetophora* locations) shared the same soil landscapes with the Project Area implies that soil biology between the two datasets is likely to differ significantly.


### 4.3 Orchid Diversity: A Surrogate for Mycorrhizal Fungi

93. The concept of locally high orchid diversity ('hotspots') is well known, and is often reported in the literature (e.g. Seaton 2007; Nurfadilah *et al.* 2013). However, orchid hotspots where



multiple orchid species co-occur are dependent on a diversity of mycorrhizal fungi being present within the soil, and some landscapes evidently support more of these than others (McCormick *et al.* 2018). Several studies have linked the patchy nature of orchid distribution to a similarly patchy distribution of mycorrhizal fungi in the soils (e.g. Voyron *et al.* 2016). In the upper Hunter region, there is a clear orchid hotspot in the Mangoola Coal area (including the target species *Diuris tricolor* and *Prasophyllum petilum*), where 15 species have been recorded (Umwelt 2006; pers. obs.). Elsewhere, other hotspots occur at Barrington Tops (Heinrich & Dowling 2000; Zoete 2000); the Hunter Economic Zone (3300 ha; 23 species) near Cessnock (Bell 2004a); and in Columbey National Park (720 ha; 16 species) near Clarencetown (including the target species *Pterostylis chaetophora*; Bell 2019d). More localised but lower levels of diversity can also occur, such as to the west of Lake Liddell where 5 species occur on the 50 ha Condran biodiversity offset property (Bell & Murray 2013), and at North Rothbury (1500 ha) where 7 species occur (Bell & Driscoll 2005). Along Thomas Mitchell Drive, 8 species were present at the time of the 1999 find of *Prasophyllum petilum*, although these are not databased (B. Holzinger pers. comm.).


94. To test the hypothesis that parts of the upper Hunter may be poorly endowed with mycorrhizal fungi, I examined the distribution of all terrestrial orchid records in the Hunter region, and specifically within the area around the Project Area. This was effectively using orchid presence as a surrogate for mycorrhizal fungi presence. I extracted from the NSW BioNet database all observation records of terrestrial orchid species within the Hunter LLS area, incorporating over 180 taxa. The de-natured locations of this dataset were not an issue for this procedure, as the level of resolution was to remain broad.
95. As indicated in [Figure 14](#) and [Figure 15](#), there is a distinct scarcity of terrestrial orchids of any kind within the Singleton to Muswellbrook region, irrespective of the extensive amount of field surveys that have been conducted over many years in relation to the development (coal mine) industry. Aside from *Diuris tricolor*, the few species recorded on Permian-aged sediments within a 15 km radius of the Project Area (*Calochilus spp.* and *Pterostylis curta* on Mt Arthur, *Pterostylis spp.* and *Diuris punctata* along Thomas Mitchell Drive, *Pterostylis nutans* and *Pterostylis bicolor* north of Lake Liddell, and *Micortis unifolia* near Ravensworth) are suggestive of a paucity of mycorrhizal fungi in these landscapes.
96. Given the level of survey that has been expended in and around the former Drayton/ Maxwell Project Area, the Mt Arthur Coal mine, the Bayswater and Liddell Power Stations, the Mount Owen mine and Ravensworth Operations, it may be expected that more terrestrial orchids would be present than is indicated through observed records. For example, the ecological assessment of the Mount Owen Continued Operations Project recorded only two species: *Acianthus fornicatus* and a *Pterostylis spp.* (Umwelt 2014), while for the Ravensworth Operations Project no orchids were recorded (Umwelt 2010). Closer to the Project Area (and while under drought conditions), the Bayswater and Liddell Power Stations upgrade project (adjacent to the east) recorded only a single *Microtis* species (Kleinfelder 2017), while at Mt Arthur (adjoining to the north and west), no orchids were recorded (Hunter Eco 2013). The low number of terrestrial orchid records in these landscapes is likely largely the result of very little mycorrhizal fungi within the soils there, and where they do occur, they are limited in extent and/or diversity. Elsewhere, some forms of mycorrhizal fungi (including those associated with orchids) have been shown to be affected by long-term agricultural grazing (e.g. Su & Guo 2007; Ba *et al.* 2012; Oja *et al.* 2017), but it is unknown if this may apply to Hunter landscapes.



**Figure 14** Point locations of all terrestrial orchids (excluding *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis chaetophora*) for the Hunter Local Land Services area. The Project Area is shown within a 15 km radius circle. Note aggregations of records indicative of diversity ‘hotspots’ for orchids, and the scarcity of orchid records within the main Hunter Valley region. Data sourced from NSW BioNet, extracted 13 July 2020.

#### 4.4 Field Inspection of the Project Area

97. I inspected all parcels of land that comprise the Project Area on 3 July 2020, in the company of Dr Colin Driscoll (Hunter Eco, on behalf of Resource Strategies). The extreme north of the main surface development area (immediately north of the old quarry) could not be accessed, however I understand that nearly all of this comprises rehabilitated land and its potential to support orchid populations is lessened considerably.
98. Field inspections involved traversing each area in a vehicle, with periodic stops to inspect the ground vegetation and its condition. Notes were made at 39 geo-referenced locations (c. 150-400 m apart) on habitat and key species present (the same locations used for the environmental analysis discussed above), and representative photographs were also taken periodically. A subjective assessment of the likelihood of orchid presence (low, medium, high) was made at each inspection point, based solely on composition and condition of grassland and woodland areas. An inspection was also made of the *Diuris* Conservation Area near to the Ancillary Disturbance Area (Ponding) 1, first reported in Cumberland Ecology (2012).



**Figure 15** Terrestrial orchids (excluding *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis chaetophora*) reported for the Project Area and immediate surrounds (15 km radius shown). Note the aggregation of records for the Mangoola Coal area in the top left corner. Data sourced from NSW BioNet, extracted 13 July 2020.

99. My initial impression of the grasslands within the Project Area during my inspection were that, although in good condition following the recent rains, they supported a different floristic composition to that occurring elsewhere where *Diuris tricolor*, *Prasophyllum petilum* and *Pterostylis chaetophora* occur. Dominant grass species across most areas included *Chloris truncata*, *Dichanthium sericeum*, *Panicum effusum*, *Chloris ventricosa*, *Bothriochloa decipiens* and *Microlaena stipoides* var. *stipoides* (Figure 16), with occasional patches of *Bothriochloa biloba*. Within this matrix was an array of flowering herbs and forbs, including *Calotis lappulacea*, *Glycina tabacina*, *Vittadinia muelleri*, *Erodium crinitum*, *Sida corrugata*, *Eremophila debilis*, *Fimbristylis dichotoma* and *Chrysocephalum apiculatum*. Some areas also supported a dominance of *Aristida ramosa*, more typical of habitat I am familiar with that supports *Diuris tricolor* and *Prasophyllum petilum*, however co-occurring with this species were several other grass taxa typical of richer clay-based soils. I saw no areas where the grasses *Entolasia stricta*, *Themeda triandra* and *Aristida vagans* dominated, such as occurs in areas frequented by *Pterostylis chaetophora*. Weed species were generally sparse or localised around former stock camps or along some drainage lines (including Ancillary Disturbance Areas 1 and 2, where *Juncus acutus* in particular predominated).



**Figure 16** Grassland within the Project Area showing grassland dominated by *Bothriochloa decipiens*, *Dichanthium sericeum* and *Chloris truncata*.

100. As summarised earlier, grassland habitats supporting *Diuris tricolor* and *Prasophyllum petilum* elsewhere in the Hunter Valley are most commonly derived from landscapes of *Eucalyptus crebra*, and tend to be dominated by *Aristida ramosa*, *Cymbopogon refractus*, *Bothriochloa decipiens* and *Sporobolus creber* (species all present in the Project Area, but rarely dominating), with other grass species present but subdominant (Figure 17, cf. Figure 16). Herbs and forbs do occur in those habitats, but they are rarely conspicuous or dominant. The grasslands observed within the Project Area appeared more diverse, in keeping with observations made in other parts of the Hunter that landscapes derived from former *Eucalyptus 'albemol'* or *Eucalyptus moluccana* woodland support more species than those derived from *Eucalyptus crebra* (e.g. such as seen at Rixs Creek; Bell 2012b). Indeed, observations of remnant canopy species made during my inspection revealed that almost all of the Project Area appeared to have once supported a woodland of *Eucalyptus 'albemol'* or *Eucalyptus moluccana*, except for some areas where *Eucalyptus dawsonii*, *Eucalyptus blakelyi* or *Eucalyptus conica* occurred.
101. Similarly, *Pterostylis chaetophora* is known predominantly from forested habitats with a grassy ground layer, with *Entolasia stricta*, *Themeda triandra* and *Aristida vagans* important. While the last of these grass species is common in the upper Hunter Valley (but rare within the Project Area), *Themeda triandra* is now rarely encountered, and *Entolasia stricta* is highly localised normally in sandstone habitats (pers. obs.).
102. Despite these observations, I am conscious of the possibility that these perceived differences in floristic composition may be a reflection on the recent growth following good falls of rain. However, apart from a few areas within the Surface Disturbance Area (Figure 18) where the hardy and grazing-tolerant *Aristida ramosa* dominated some sections, this species formed only a small component of the bulk of lands inspected; it was nearly always present but sub-ordinate to other grasses and herbs. Given the long history of grazing across the Project Area, it may be expected that *Aristida ramosa* would have been the dominant grass species, however clearly soil type rather than past land management appears to drive species composition in these landscapes.



Figure 17 Grassland supporting *Diuris tricolor* and *Prasophyllum petilum* at Mangoola Coal, dominated by *Aristida ramosa* and *Cymbopogon refractus*.



Figure 18 Grassland within the proposed Surface Area Disturbance area, showing (predominantly) swards of *Aristida ramosa* with *Dichanthium sericeum* and *Chloris truncata*.

103. On the GIS, I then examined the vegetation community mapping presented in Hunter Eco (2019) to determine if it accurately portrayed the diversity and distribution of vegetation that I inspected, and found it to represent well the vegetation patterns within the Project Area. I was satisfied, therefore, that I could use the Hunter Eco vegetation community mapping to create maps of likely orchid habitat, so that if necessary, estimates of the number of hectares anticipated to support viable orchid populations could be calculated (**Section 5**). Additional guidance was provided by my own field notes and data analyses, the plot data collected by Hunter Eco and aerial imagery.
104. In determining the suitability of the Project Area as orchid habitat, I drew on my experience from surveying for *Diuris tricolor* and *Prasophyllum petilum* in the Mangoola area over ten years, and for *Pterostylis chaetophora* in the mid-lower Hunter Valley across four sites over three years. Part of this experience included observations of orchids growing in somewhat surprising situations, which may otherwise be glossed over as unsuitable habitat. For example, observations of *Diuris tricolor* growing on agricultural contour banks (**Figure 19**), in heavily weed-infested derived grasslands where no other native species were apparent (**Figure 20**), and proliferating on the manicured lawns of farm homesteads (**Figure 21**). I have also observed *Diuris tricolor* growing within a former vineyard on raised garden beds, and along the margins of management trails. Likewise, *Pterostylis chaetophora* also commonly grows along trail and road edges (**Figure 22**) and in other areas where some ground disturbance has occurred. Collectively, observations of orchids growing in such disturbed habitats suggest that perceived low condition of potentially appropriate habitat should not be immediately dismissed as unlikely to support orchids.



**Figure 19** *Diuris tricolor* and other orchids growing over a constructed contour bank, Mangoola.



Figure 20 *Diuris tricolor* growing with exotic weeds in low quality grassland near Mangoola Coal.



Figure 21 *Diuris tricolor* proliferating in mown lawns of a farm homestead near Mangoola Coal.



Figure 22 *Pterostylis chaetophora* growing in roadside gutter, Columbey National Park.

#### 4.5 Concluding Opinion: Are the Target Species Present?

105. Based on the various floristic and environmental analyses discussed earlier in this section, I concluded that there is limited potential for *Diuris tricolor* or *Prasophyllum petilum* to be present within the Project Area, and that it is unlikely for *Pterostylis chaetophora* to occur at all. I based this opinion largely on the preceding analysis of floristic composition of available habitat, geological unit, soil landscape, annual rainfall and the inferred absence or paucity of mycorrhizal fungi (using reported orchid presence and diversity as a surrogate) (Table 13).
106. The two communities potentially supporting *Diuris tricolor* and/or *Prasophyllum petilum* (Units 9a and 11 of Hunter Eco 2019) occur only at Ancillary Disturbance Area 1 and the Product Stockpile Extension area respectively. Since these conclusions were made, Hunter Eco was commissioned to undertake targeted surveys of these two communities and no threatened flora species were recorded within the Project Area. Consequently, *Diuris tricolor* and *Prasophyllum petilum* (syn. *Prasophyllum* sp. Wybong) are unlikely to be present anywhere in the Project Area.



**Table 13 Concluding Reasoning for Orchid Presence or Absence.**

Species	Factor	Reasoning
<i>Diuris tricolor</i>	floristic composition	<p>Analysis of floristic data (using the upper 95% of species comprising defined vegetation communities within the Project Area) revealed only one community (<b>Unit 9a: Ironbark - Grey Box Grassy Woodland DNG</b>) as being closely related to my own defined communities supporting <i>Diuris tricolor</i> at Mangoola. This community has been mapped by Hunter Eco (2019) for 0.3 ha of the Project Area, at Ancillary Disturbance Area 1.</p> <p>Additionally, the close proximity (c. 400 m) of a population of <i>Diuris tricolor</i> to the Product Stockpile Extension area, and in vegetation similar to that present there, suggests that <b>Unit 11: Grey Box – Spotted Gum – Narrow-leaved Ironbark woodland</b>, mapped by Hunter Eco (2019) across 1.3 ha of the Project Area, also potentially supports this species.</p>
	geological unit	There is only <b>minor correlation</b> (6%) between geological units within the Project Area and those supporting <i>Diuris tricolor</i> elsewhere in the region.
	soil landscape	There is only <b>minor correlation</b> (5%) between soil landscapes within the Project Area and those supporting <i>Diuris tricolor</i> elsewhere in the region.
	mycorrhizal fungi	Based on few records of all terrestrial orchid species within and around the Project Area, it appears that there are <b>very limited and/or highly localised mycorrhizal fungi</b> present in soils, reducing the likelihood of <i>Diuris tricolor</i> or <i>Prasophyllum petilum</i> being present.
<i>Prasophyllum petilum</i>	floristic composition	<p>Analysis of floristic data (using the upper 95% of species comprising defined vegetation communities within the Project Area) revealed only one community (<b>9a - Ironbark - Grey Box Grassy Woodland DNG</b>) as being closely related to my own defined communities supporting <i>Prasophyllum petilum</i> at Mangoola. This community has been mapped by Hunter Eco (2019) for 0.3 ha of the Project Area, at Ancillary Disturbance Area 1.</p> <p>Additionally, the relatively close proximity (c. 1300 m) of an historical population of <i>Prasophyllum petilum</i> to the Product Stockpile Extension area, and in vegetation similar to that present there, suggests that <b>Unit 11: Grey Box – Spotted Gum – Narrow-leaved Ironbark Woodland</b>, mapped by Hunter Eco (2019) across 1.3 ha of the Project Area, also potentially supports this species.</p>
	geological unit	There is only <b>very minor correlation</b> (0.2%) between geological units within the Project Area and those supporting <i>Prasophyllum petilum</i> elsewhere in the region.

Species	Factor	Reasoning
	soil landscape	There is only <b>very minor correlation</b> (0.2%) between soil landscapes within the Project Area and those supporting <i>Prasophyllum petilum</i> elsewhere in the region.
	mycorrhizal fungi	Based on few records of all terrestrial orchid species within and around the Project Area, it appears that there are very limited and/or highly localised mycorrhizal fungi present in the available soils, reducing the likelihood of <i>Prasophyllum petilum</i> being present.
<i>Pterostylis chaetophora</i>	floristic composition	Analysis of floristic data (using the upper 95% of species comprising defined vegetation communities within the Project Area) revealed <b>no relationship</b> with my own defined communities supporting <i>Pterostylis chaetophora</i> at Columbey National Park. Despite superficial similarities of community 11 (Grey Box - Spotted Gum - Narrow-leaved Ironbark Woodland) from within the Project Area to elements of the Columbey vegetation, there are few co-occurring dominants.
	geological unit	There is only <b>minor correlation</b> (3%) between geological units within the Project Area and those supporting <i>Pterostylis chaetophora</i> elsewhere in the region.
	soil landscape	There is only <b>very minor correlation</b> (0.8%) between soil landscapes within the Project Area and those supporting <i>Pterostylis chaetophora</i> elsewhere in the region.
	annual rainfall	Annual mean rainfall across all locations supporting <i>Pterostylis chaetophora</i> (n=128) is 1005 mm/yr (stdev 98.4), while the annual mean rainfall of the Project Area is <b>considerably drier</b> at 640 mm/yr (stdev 20.8).
	mycorrhizal fungi	Based on few records of all terrestrial orchid species within and around the Project Area, it appears that there are very limited and/or highly localised mycorrhizal fungi present in the available soils, reducing the likelihood of <i>Pterostylis chaetophora</i> being present.

## 5. Criterion (d) – Size of Population or Habitat

107. In order to determine the extent of potential habitat for *Diuris tricolor*, *Prasophyllum petilum* and/or *Pterostylis chaetophora* within the Project Area, I considered the elements outlined above in [Table 13](#) and used the mapping of Hunter Eco (2019) to spatially represent that habitat. Following my analyses, habitat for *Diuris tricolor* and *Prasophyllum petilum* was considered to potentially occur within vegetation mapped as Ironbark - Grey Box Grassy Woodland DNG (Unit 9a) or Grey Box – Spotted Gum – Narrow-leaved Ironbark Woodland (Unit 11) by Hunter Eco (2019). These units have been mapped across 0.3 ha of Ancillary Disturbance Area 1, and 1.3 ha for the Product Stockpile Extension area respectively. No other grasslands or woodlands in the Project Area carry vegetation similar to this, supporting both by my field inspection and the comparative numerical floristic analysis I performed on Hunter Eco (2019)

data. Surveys of these areas by Hunter Eco in 2020 did not record *Diuris tricolor* or *Prasophyllum petilum*, suggesting that these species are unlikely to be present within the Project Area.

108. I do not consider *Pterostylis chaetophora* to be present anywhere within the Project Area. Almost all of the known records of this species occur in more coastal habitats receiving higher rainfall than the Project Area, except for two unconfirmed populations at Mangoola Coal and Wingen Maid Nature Reserve. Those two populations, however, are closely tied (geographically) to remnant Triassic Narrabeen series outcrops, and/or colluvial material derived from those rock types, none of which occur within the Project Area.
109. While rarely recognised in assessments of potential orchid habitat, the occurrence of the required mycorrhizal fungi for each of the three target orchid species (and indeed for all terrestrial orchids) is possibly the most important determiner of orchid presence/absence. Unfortunately, it is the one factor that we know very little about and represents an attribute that cannot be readily determined without significant and comprehensive 'baiting' of soils to map their distribution. As a proxy, in this assessment examination of database records for all orchid species across the Hunter region revealed a clear scarcity of terrestrial orchids from the landscapes in and around the Project Area (irrespective of the extensive flora survey effort that has been expended in this area over many years), implying that mycorrhizal fungi themselves are scarce. With no fungi, there can be no orchids (see **Section 2.4.2**).
110. In an earlier report for the adjacent Bayswater and Liddell Power Station upgrade (Bell 2020c), I conservatively estimated approximately 166 ha of derived grassland as potentially supporting *Diuris tricolor* and no *Prasophyllum petilum*, but based on soil landscapes, floristics and proximate records stated that large populations would be unlikely there. Additional research undertaken since then into soil biology, mycorrhizal fungi and terrestrial orchid diversity in the upper Hunter now supports the contention that only small and highly localised populations appear present in these landscapes. This is reinforced by the known populations of *Diuris tricolor* and *Prasophyllum petilum* along Thomas Mitchell Drive, and in the *Diuris* Conservation Area near Ancillary Disturbance Area 1. These localised populations, all occurring in or close to woodland (rather than in derived grassland) and often in areas supporting several other orchid species, are suggestive of highly localised mycorrhizal fungi. Given the fact that no orchid species of any kind has ever been recorded within the Project Area (and only 7 species detected historically within a 15 km radius of it, most as single records), it seems highly unlikely that sufficient mycorrhizal fungi occur in the soils there to support *Diuris tricolor* or *Prasophyllum petilum*. It is quite plausible that a prolonged period of agricultural grazing, incorporating soil enrichment and compaction, over at least 150 years across most of the Project Area has altered soil chemistry and depleted the reserves of mycorrhizal fungi, to the point now that they (and consequently the orchids that rely on them) are largely absent from grassland areas.
111. Considering all of the preceding discussion, I consider the extent of potential habitat within the Project Area to be as shown in [Figure 23](#), and comprises:
  - 1.6 ha for *Diuris tricolor*
  - 1.6 ha for *Prasophyllum petilum*
  - 0 ha for *Pterostylis chaetophora*




Figure 23 Habitat considered suitable (black shading) for *Diuris tricolor* and *Prasophyllum petilum* within the Project Area. Base mapping from Hunter Eco (2019).

## **6. Criterion (e) – Documents & Data Reviewed**

112. I have been provided with the following reports and datasets from Resource Strategies and DPIE to assist in this review:

- GIS files showing Project boundaries and vegetation community mapping
- flora and vegetation mapping report for the proposed Maxwell Project (Hunter Eco 2019)
- an extract of the NSW BioNet database, under licence ASH20009, detailing as-held records for the three target orchids within the Hunter LLS area

All other published and unpublished reports, papers and maps that form part of this assessment have been cited in the normal way, with publication details contained in **Section 9** or advised within the text.

## 7. Criterion (f) – Expert Credentials

113. Under the requirements of the BAM (DPIE 2020), an expert report can be prepared by an endorsed person in the place of undertaking field survey. This report must include information on the credentials of the expert, including the following:

a. *the expert’s academic qualifications such as relevant degrees, post graduate qualifications;*

I possess three degrees in the science and ecology field: a Bachelor of Science (1988), Bachelor of Science (Honours) (1990) and a Doctor of Philosophy in vegetation science (2013).

b. *their history of experience in the ecological research, habitat assessment and survey method, for the relevant species;*

In regard to the threatened orchid species that are the subject of this expert report (*Diuris tricolor*, *Prasophyllum petilum*, *Pterostylis chaetophora*), I have been surveying and monitoring these species for between three (*Pterostylis chaetophora*) and 11 (*Diuris tricolor*, *Prasophyllum petilum*) consecutive years at various locations in the Hunter Valley, including the annual monitoring of over 3000 translocated specimens since 2010. Targeted surveys have incorporated systematic open-ended transect surveys in appropriate habitat, using GPS devices to record tracks searched and orchids located. Separation distances between adjacent search transects vary in relation to quality of habitat and visibility. Search times have only occurred when other known reference populations have been in flower.

c. *a resume detailing projects pertaining to the survey of the relevant species (including the locations and dates of the work), their employers’ names and periods of employment (where relevant) over the previous 10 years;*

I am the principal and owner of Eastcoast Flora Survey, established in the Hunter Valley in October 1996 and spanning a continual period of dedicated flora consulting of nearly 25 years. Since 2014, I have also been a Conjoint Fellow at the University of Newcastle (School of Environmental and Life Sciences), where I am a member of two research groups: the Centre for Plant Science and the Conservation Biology Research Group. My full Curriculum Vitae are appended as [Appendix 5](#) to this report.

In relation to the relevant species that are the subject of this report (*Diuris tricolor*, *Prasophyllum petilum*, *Pterostylis chaetophora*), the following projects and publications pertain to these:

- Bell, S.A.J. (submitted 1) Floristic community diversity in derived native grasslands: a case study from the upper Hunter Valley of New South Wales. *Cunninghamia* (submitted).
- Bell, S.A.J. (submitted 2) Successful recruitment following translocation of a threatened terrestrial orchid (*Diuris tricolor*) into mining rehabilitation in the Hunter Valley of NSW. *Ecological Management and Restoration* (submitted)
- Bell, S.A.J. (2020) Translocation of threatened terrestrial orchids into non-mined and post-mined lands in the upper Hunter Valley of New South Wales, Australia. *Restoration Ecology* 28 1396-1407.

- Bell, S.A.J. (2020) *Survey and monitoring of the Vulnerable Pterostylis chaetophora (Rusty Greenhood) in the Lower Hunter Valley, NSW: 2019 Results*. Unpublished Report to NSW Office of Environment and Heritage. June 2020. Eastcoast Flora Survey.
- Bell, S.A.J. (2020) *Expert Report - Expected Presence of Threatened Terrestrial Orchids (Diuris tricolor & Prasophyllum petilum): Bayswater Water and Other Associated Operational Works Project*. Unpublished Report to Kleinfelder. May 2020. Eastcoast Flora Survey.
- Bell, S.A.J. (2020) *Monitoring of translocated threatened orchids (Diuris tricolor, Prasophyllum petilum) at Mangoola Coal: 2019 Results*. Unpublished Report to Mangoola Coal. February 2020.
- Bell, S.A.J. & Hillier, P. (2020) Targeted surveys of a poorly conserved threatened orchid (*Pterostylis chaetophora*) in Columbey National Park (Hunter Valley, NSW) reveal substantial populations and elucidate occupied habitat. *Cunninghamia* 20: 199-207.
- Bell, S.A.J., Murray, M., & Sims, R. (2020) *Flora and Fauna Monitoring at Condran, Muswellbrook LGA: 2019 Results*. Unpublished Report to Bulga Surface Operations (Glencore). January 2020. Eastcoast Flora Survey & Forest Fauna Surveys Pty Ltd.
- Bell, S.A.J. (2019) *Expert Report: Expected presence of threatened terrestrial orchids (Diuris tricolor & Prasophyllum petilum), Mangoola Coal Continued Operations Project*. Unpublished Report to Umwelt (Australia) Pty Ltd. December 2019.
- Bell, S.A.J. (2019) *Survey and monitoring of the Vulnerable Pterostylis chaetophora (Rusty Greenhood) in the Lower Hunter Valley, NSW: 2018 Results*. Unpublished Report to NSW Office of Environment and Heritage. April 2019. Eastcoast Flora Survey.
- Bell, S. (2019) Translocation success is all about detection: experiences with two threatened orchids from the Hunter Valley of NSW. *Australasian Plant Conservation* 28: 27-31.
- Bell, S.A.J., Murray, M., & Sims, R. (2019) *Flora and Fauna Monitoring at Condran, Muswellbrook LGA: 2018 Results*. Unpublished Report to Bulga Surface Operations (Glencore). January 2019. Eastcoast Flora Survey & Forest Fauna Surveys Pty Ltd.
- Bell, S.A.J., Murray, M., & Sims, R. (2018) *Flora and Fauna Monitoring at Condran, Muswellbrook LGA: 2017 Results*. Unpublished Report to Bulga Surface Operations (Glencore). March 2018. Eastcoast Flora Survey & Forest Fauna Surveys Pty Ltd.
- Bell, S.A.J. (2018) *Monitoring of translocated threatened orchids (Diuris tricolor, Prasophyllum petilum) at Mangoola Coal: 2017 Results*. Unpublished Report to Mangoola Coal. February 2018.
- Bell, S.A.J. (2017) *Targeted survey for the threatened Diuris tricolor at Persoonia Park, North Rothbury, Hunter Valley*. Unpublished Report to Office of Environment & Heritage. November 2017. Eastcoast Flora Survey.
- Bell, S.A.J. (2017) *Baseline monitoring and survey of the vulnerable Pterostylis chaetophora (Orchidaceae) at North Rothbury, Hunter Valley, NSW*. Unpublished Report to NSW Office of Environment & Heritage. October 2017. Eastcoast Flora Survey.
- Bell, S.A.J. & Murray, M. (2017) *Flora and Fauna Monitoring at Condran, Muswellbrook LGA: 2016 Results*. Unpublished Report to Bulga Surface Operations (Glencore). January 2017. Eastcoast Flora Survey & Forest Fauna Surveys Pty Ltd.
- Bell, S.A.J. (2017) *Targeted Orchid Survey: Addendum to Pre-clearance Surveys, Borehole Explorations Areas, Rix's Creek North Mine*. Unpublished Report to Rix's Creek Pty Limited. October 2017. Eastcoast Flora Survey.

- Bell, S.A.J. & Murray, M. (2016) *Flora and Fauna Monitoring at Condran, Muswellbrook LGA: 2015 Results*. Unpublished Report to Bulga Surface Operations (Glencore). May 2016. Eastcoast Flora Survey & Forest Fauna Surveys Pty Ltd.
- Bell, S.A.J. & Murray, M. (2015) *Flora and Fauna Monitoring at Condran, Muswellbrook LGA: 2014 Results*. Unpublished Report to Bulga Surface Operations (Glencore). January 2015. Eastcoast Flora Survey & Forest Fauna Surveys Pty Ltd.
- Bell, S.A.J. & Driscoll, C (2014) *Assessment and mapping of vegetation in the Bylong Valley: Authorisations 287 & 342*. Unpublished Final Report to Hansen Bailey Pty Ltd. Eastcoast Flora Survey. December 2014.
- Bell, S.A.J. (2013) *Monitoring of translocated threatened orchids (Diuris tricolor, Prasophyllum sp. Wybong C.Phelps ORG 5269) at Mangoola Coal: 2013 Results*. Unpublished Report to Mangoola Coal. November 2013. Eastcoast Flora Survey.
- Bell, S.A.J. & Murray, M. (2013) *Flora and Fauna Monitoring at Condran, Muswellbrook LGA*. Unpublished Report to Bulga Surface Operations (Glencore). November 2013. Eastcoast Flora Survey & Forest Fauna Surveys Pty Ltd.
- Bell, S.A.J. (2013) *Monitoring of translocated threatened orchids (Diuris tricolor, Prasophyllum sp. Wybong C.Phelps ORG 5269) at Mangoola Coal: Status Report 2012*. Unpublished Report to Mangoola Coal. Eastcoast Flora Survey, January 2013.
- Bell, S.A.J. (2012) *Targeted terrestrial orchid surveys at Mangoola Coal, Upper Hunter Valley: Spring 2011*. Unpublished Report to Mangoola Coal. Eastcoast Flora Survey, January 2012.
- Bell, S.A.J. & Copeland, L. (2010) *Targeted terrestrial orchid surveys at Mangoola Coal, Upper Hunter Valley: Spring 2010*. Unpublished Report to Mangoola Coal, October 2010. Eastcoast Flora Survey.
- Bell, S.A.J. & Copeland, L. (2010) *A strategy for the translocation of threatened terrestrial orchids at Mangoola Coal, Upper Hunter Valley*. Unpublished Report to Mangoola Coal, September 2010. Eastcoast Flora Survey.
- Bell, S.A.J. & Copeland, L. (2009) *Targeted terrestrial orchid survey, Mangoola, Upper Hunter Valley*. Spring 2009. Unpublished Report to Mangoola Coal. Eastcoast Flora Survey, November 2009.
- Bell, S.A.J. (2009) *Targeted terrestrial orchid survey of the ex-Nipol property, near Denman, Upper Hunter Valley*. Unpublished report to Mangoola Coal. Eastcoast Flora Survey, November 2009.

d. *peer-reviewed publications on the species or other evidence that the person is a well-known authority on the species to which the survey relates;*

I have published two papers specifically addressing *Diuris tricolor* and *Prasophyllum petilum*: Bell (2019a, assessing translocation success in these species) and Bell (2020d, comparing translocation efforts in mined and non-mined lands). A third published paper details population size and occupied habitat of *Pterostylis chaetophora* in the lower Hunter (Bell & Hillier 2020), while a fourth presents the results of new recruitment in *Diuris tricolor* in mine rehabilitation (Bell submitted 2).

I have also published on several other threatened orchid species (e.g. *Cryptostylis hunteriana*: Bell 2001a, de Lacey *et al.* 2012a, b, de Lacey *et al.* 2013; *Thelymitra adorata*: Bell *et al.* 2005; *Diuris praecox*: Yare *et al.* 2020) and non-orchid threatened taxa (e.g. *Acacia dangarensis*: Bell & Elliott 2013; *Acacia pendula*: Bell 2018; Bell *et al.* 2007, Bell & Driscoll 2014, 2016; *Acacia wollarensis*: Bell & Driscoll 2017, Bell & Kodela 2018; *Angophora inopina*: Bell 2004b; *Banksia conferta*: Bell 2017b; *Commersonia rosea*: Bell & Copeland 2004, Bell &



Holzinger 2015; *Dracophyllum macranthum*: Bell & Sims 2018; *Eucalyptus expressa*: Bell & Nicolle 2012; *Eucalyptus calidissima*; Bell & Klaphake 2020; *Eucalyptus dealbata* subsp. *aperticola*: Bell & Nicolle 2020; *Hibbertia procumbens*: Bell 2002, Bell & Driscoll 2005b; *Leionema lamprophyllum* subsp. *fractum*: Bell & Walsh 2015; *Monotaxis macrophylla*: Bell & Holzinger 2015; *Senecio linearifolius* var. *dangarensis* and *S. spathulatus* var. *attenuatus*: Mickaill *et al.* 2020), together with those examining a range of significant and threatened species in sandstone habitats of the Hunter Valley (23 taxa; Bell 2001b) and those present in Wollemi National Park (110 taxa; Bell 2008, 2019c). I am also the lead author of a recently published book with CSIRO Publications (Bell *et al.* 2019), detailing 54 of the endemic plant species of the Hunter Region, many of which are threatened species. One other recent paper, of which I am co-author, examined the conservation status of 822 eucalypt taxa from across Australia (Fensham *et al.* 2020).

I have been surveying and monitoring two of the target species for over 10 years in the Hunter Valley (and the third for three years) and am acutely aware of their habitat requirements and variability in flowering from year to year. Additionally, Dr Lachlan Copeland (Eco Logical Australia & orchid taxonomist) has endorsed me as a recognised authority on the field ecology of *Diuris tricolor* and *Prasophyllum petilum* (see letter attached in [Appendix 6](#)).

## 8. Conclusion

114. Maxwell Ventures (Management) Pty Ltd propose to establish an underground coal mine within EL5460, which will require limited ground disturbance across the full Project Area. Nevertheless, proposed disturbances associated with planned expansion of the Product Stockpile Extension area, establishment of Surface Development Areas to accommodate underground entrance surface facilities and a transport/services corridor, proposed Edderton Rd Realignment to avoid potential road damage from subsidence, and Ancillary Disturbance Areas (Ponding) in response to modelled ground subsidence, may impact on terrestrial orchids, if present.
115. No known records of *Diuris tricolor*, *Prasophyllum petilum* or *Pterostylis chaetophora* exist for the Project Area, and after assessment I considered that approximately 1.6 ha of the proposed 320 ha (0.5%) of ground disturbance area may provide habitat for two of these species (*Diuris tricolor*, *Prasophyllum petilum* [syn. *Prasophyllum* sp. Wybong]). I do not consider that the third target species, *Pterostylis chaetophora*, is likely to occur anywhere within the Project Area.
116. *Diuris tricolor* and *Prasophyllum petilum* occupy extensive geographical ranges outside of the Hunter region, however *Pterostylis chaetophora* is endemic here. Records for *Diuris tricolor* exist close to the Project Area, near Ancillary Disturbance Area 1 and along Thomas Mitchell Drive, and *Prasophyllum petilum* has historically been recorded at the eastern end of Thomas Mitchell Drive but is presumed extinct (B. Holzinger pers. comm.). There are no validated populations of *Pterostylis chaetophora* west of North Rothbury (52 km to the south-east), although two records at Mangoola Coal (21 km north-west) and Wingen Maid Nature Reserve (55 km north) remain unvalidated.
117. Following a single day site inspection of the Project Area in July 2020, I noted observable differences in the floristic composition of habitats to other areas where I know the three target orchids from, and the predominance of former Box (*Eucalyptus 'albemol'* and *Eucalyptus moluccana*) landscapes within these largely cleared lands conflicts with my experience of orchid habitat elsewhere. Differences were supported by a review of ecological information prepared by Hunter Eco (2019) for the Project Area, the mapping of which I found to be accurate and acceptable. Numerical analysis of defined vegetation communities revealed clear differences to habitat from elsewhere in the region known to support *Pterostylis chaetophora*, but one community similar to habitats typical of *Diuris tricolor* and *Prasophyllum petilum*. Additionally, the close proximity of known populations of *Diuris tricolor* (c. 400 m) and *Prasophyllum petilum* (1300 m) to the northern section of the Project Area suggests that potential habitat for these may also occur in a second community. I therefore consider that the vegetation units defined by Hunter Eco (2019) as Ironbark - Grey Box Grassy Woodland DNG (Unit 9a) and Grey Box – Spotted Gum – Narrow-leaved Ironbark Woodland (Unit 11) provide the only potential habitat for *Diuris tricolor* and *Prasophyllum petilum*, which together comprise 1.6 ha (0.5%) of the Project Area. Dr Colin Driscoll (Hunter Eco) was commissioned by Malabar Resources to conduct a targeted survey of this area during suitable survey conditions in 2020, and none of the target species were found. Consequently, *Diuris tricolor* and *Prasophyllum* (syn. *Prasophyllum* sp. Wybong) are unlikely to be present within the Project Area.
118. In support, a detailed assessment of environmental attributes across all Hunter populations of *Diuris tricolor* (n=983), *Prasophyllum petilum* (n=485) and *Pterostylis chaetophora* (n=128) found differences in geological units, soil landscapes, and (for *Pterostylis chaetophora*) annual

rainfall when compared to the Project Area. Of these, soil landscapes were seen as particularly important given they best encapsulate the likely distribution of mycorrhizal fungi across the region, without which no orchids would germinate and prosper. As a surrogate for fungi distribution, an assessment of all terrestrial orchid observation records for the upper Hunter Valley demonstrated a scarcity of orchids in and around the Project Area, suggesting that mycorrhizal fungi were therefore also scarce or localised. This lends further credence to my conclusion that *Diuris tricolor* or *Prasophyllum petilum* (syn. *Prasophyllum* sp. Wybong) are unlikely to be present within the Project Area.

119. The combination of geology, soil landscape, floristic, rainfall, and general orchid distribution (as a surrogate for mycorrhizal fungi) suggest that the Project Area is unlikely to support extensive populations of *Diuris tricolor* or *Prasophyllum petilum* (syn. *Prasophyllum* sp. Wybong), and none of *Pterostylis chaetophora*. The areas where I consider the former two species may occur occupy <1% of the total disturbance footprint, but targeted searches in these areas did not record these species.

## 9. References

- Abel Ecology (2005) *Flora and fauna report for 168 Bells Lane, Denman in the upper Hunter Valley, New South Wales*. Unpublished Report prepared for Anvil Hill Project Watch Association.
- Adams, P.B. & Lawson, S.D. (1993) Pollination in Australian orchids: A critical assessment of the literature 1882-1992. *Australian Journal of Botany* 41: 553-575.
- Ba, L., Ning, J., Wang, D., Facelli, E., Facelli, J.M., Yang, Y. & Zhang, L. (2012) The relationship between the diversity of arbuscular mycorrhizal fungi and grazing in a meadow steppe. *Plant and Soil* 352: 143-156.
- Backhouse, G. & Cameron, D. (2005) Application of IUCN 2001 Red List categories in determining the conservation status of native orchids of Victoria, Australia. *Selbyana* 26(1,2): 58-74.
- Backhouse, G.N., Bates, R.J., Brown, A.P. & Copeland, L.M. (2019) *A Checklist of the Orchids of Australia Including its Island Territories*. Second Edition. Melbourne, Australia. ISBN 978-0-9946489-3-8.
- Backhouse, G., Kosky, B., Rouse, D. & Turner, J. (2016) *Bush Gems: A Guide to the Wild Orchids of Victoria, Australia*. Published by Authors, Melbourne, August 2016. ISBN: 978-0-9946489-0-7.
- Bell, S.A.J. (2001a) Notes on population size and habitat of the vulnerable *Cryptostylis hunteriana* Nicholls (Orchidaceae) from the Central Coast of New South Wales. *Cunninghamia* 7(2): 195-204.
- Bell, S.A.J. (2001b) Notes on the distribution and conservation status of some restricted plant species from sandstone environments of the upper Hunter Valley, New South Wales. *Cunninghamia* 7(1): 77-88.
- Bell, S.A.J. (2002) Habitat of the endangered *Hibbertia procumbens* (Labill.) DC (Dilleniaceae) from the Central Coast of New South Wales. *Victorian Naturalist* 119(2): 69-74.
- Bell, S.A.J. (2004a) *The vegetation of the Hunter Economic Zone (HEZ), Cessnock LGA, New South Wales*. Unpublished Report to Harper Somers O'Sullivan, January 2004. Eastcoast Flora Survey.
- Bell, S.A.J. (2004b) Distribution and habitat of the vulnerable tree species, *Angophora inopina* (Myrtaceae), on the Central Coast of New South Wales. *Cunninghamia* 8(4): 477-484.
- Bell, S.A.J. (2008) Rare or threatened vascular plant species of Wollemi National Park, central eastern New South Wales. *Cunninghamia* 10(3): 331-371.
- Bell, S.A.J. (2009) Vegetation and floristics of Columbey National Park, lower Hunter Valley, New South Wales. *Cunninghamia* 11(2): 241-275.
- Bell, S.A.J. (2012a) *Diversity and condition of derived grasslands at Mangoola Coal, Upper Hunter Valley: Stage 3 - 2011 Surveys*. Unpublished Report to Mangoola Coal, June 2012.

- Bell, S.A.J. (2012b) *Assessment of vegetation for expansion of Rix's Creek Mine, Hunter Valley*. Unpublished Report to Rix's Creek Pty Limited. November 2012.
- Bell, S.A.J. (2016) *Monitoring of translocated threatened orchids (Diuris tricolor, Prasophyllum petilum) at Mangoola Coal: 2015 Results*. Unpublished Report to Mangoola Coal. April 2016. Eastcoast Flora Survey.
- Bell, S.A.J. (2017a) *Targeted survey for the threatened Diuris tricolor at Persoonia Park, North Rothbury, Hunter Valley*. Unpublished Report to Office of Environment and Heritage. November 2017. Eastcoast Flora Survey.
- Bell, S. (2017b) New insights into the ecology of the critically endangered *Banksia conferta* (Proteaceae) from the mid-north coast of NSW. *Australasian Plant Conservation* 26(1): 15-18.
- Bell, S.A.J. (2018) Fate of a rare flowering event in an endangered population of *Acacia pendula* (Weeping Myall) from the Hunter Valley, New South Wales. *Cunninghamia* 18: 79-88.
- Bell, S. (2019a) Translocation 'success' is all about detection: experiences with two threatened orchids from the Hunter Valley of NSW. *Australasian Plant Conservation* 28: 27-31.
- Bell, S.A.J. (2019b) *Expert Report: Expected presence of threatened terrestrial orchids (Diuris tricolor & Prasophyllum petilum), Mangoola Coal Continued Operations Project*. Unpublished Report to Umwelt (Australia) Pty Ltd. December 2019.
- Bell, S.A.J. (2019c) Additions and amendments to the rare or threatened vascular plants of Wollemi National Park, central eastern New South Wales. *Cunninghamia* 19: 43-56.
- Bell, S.A.J. (2019d) *Survey and monitoring of the Vulnerable Pterostylis chaetophora (Rusty Greenhood) in the Lower Hunter Valley, NSW: 2018 Results*. Unpublished Report to NSW Office of Environment and Heritage. May 2019. Eastcoast Flora Survey.
- Bell, S.A.J. (2020a) *Monitoring of translocated threatened orchids (Diuris tricolor, Prasophyllum petilum) at Mangoola Coal: 2019 Results*. Unpublished Report to Mangoola Coal. February 2020. Eastcoast Flora Survey.
- Bell, S.A.J. (2020b) *Survey and monitoring of the Vulnerable Pterostylis chaetophora (Rusty Greenhood) in the Lower Hunter Valley, NSW: 2019 Results*. Unpublished Report to NSW Office of Environment and Heritage. June 2020. Eastcoast Flora Survey.
- Bell, S.A.J. (2020c) *Expert Report - Expected Presence of Threatened Terrestrial Orchids (Diuris tricolor & Prasophyllum petilum): Bayswater Water and Other Associated Operational Works Project*. Unpublished Report to Kleinfelder. May 2020. Eastcoast Flora Survey.
- Bell, S.A.J. (2020d) Translocation of threatened terrestrial orchids into non-mined and post-mined lands in the upper Hunter Valley of New South Wales, Australia. *Restoration Ecology* 28: 1396-1407.
- Bell, S.A.J. (submitted 1) Floristic community diversity in derived native grasslands: a case study from the upper Hunter Valley of New South Wales. *Cunninghamia* submitted.

- Bell, S.A.J. (submitted 2) Successful recruitment following translocation of a threatened terrestrial orchid (*Diuris tricolor*) into mining rehabilitation in the Hunter Valley of NSW. *Ecological Management and Restoration* (submitted)
- Bell, S.A.J. & Copeland, L.M. (2004) *Commersonia rosea* (Malvaceae s.l.: Lasiopetaleae): a new, rare fire-ephemeral species from the upper Hunter Valley, New South Wales. *Telopea* 10(2): 581-587.
- Bell, S.A.J. & Driscoll, C. (2005a) *Vegetation survey of 'Sweetwater', North Rothbury, mid Hunter Valley, New South Wales*. Unpublished Report to Harper Somers O'Sullivan, October 2005. Eastcoast Flora Survey.
- Bell, S. & Driscoll, C. (2005b) New records of the endangered *Hibbertia procumbens* from the Central Coast of NSW. *Australasian Plant Conservation* 13(4): 24-25.
- Bell, S. & Driscoll, C. (2014) *Acacia pendula* (Weeping Myall) in the Hunter Valley of New South Wales: early explorers' journals, database records and habitat assessments raise doubts over naturally occurring populations. *Cunninghamia* 14: 179–200.
- Bell, S.A.J. & Driscoll, C. (2016) Hunter Valley Weeping Myall Woodland – is it really definable and defensible with and without Weeping Myall (*Acacia pendula*)? *Cunninghamia* 16: 15-30.
- Bell, S.A.J. & Driscoll, C. (2017) *Acacia wollarensis* (Fabaceae, Mimosoideae sect. Botrycephalae), a distinctive new species endemic to the Hunter Valley of New South Wales, Australia. *Telopea* 20: 125-136.
- Bell, S. & Elliott, M. (2013) Preliminary results suggest fire is required to maintain *Acacia dangarensis*, a threatened single-population endemic from the Hunter Valley of NSW. *Australasian Plant Conservation* 22(1): 9-10.
- Bell, S.A.J. & Hillier, P. (2020) Targeted surveys of a poorly conserved threatened orchid (*Pterostylis chaetophora*) in Columbey National Park (Hunter Valley, NSW) reveal substantial populations and elucidate occupied habitat. *Cunninghamia* 20: 199-207.
- Bell, S. & Holzinger, B. (2015) Wildfire reveals new populations of the endangered *Commersonia rosea* and *Monotaxis macrophylla* in northern Wollemi National Park, NSW. *Australasian Plant Conservation* 23: 2-4.
- Bell S.A.J. & Kodela P.G. (2018) *Acacia wollarensis*. In: *Flora of Australia*. Australian Biological Resources Study, Department of the Environment and Energy, Canberra. <https://profiles.ala.org.au/opus/foa/profile/Acacia%20wollarensis>.
- Bell, S.A.J. & Klaphake, V. (2020) *Eucalyptus calidissima* (Myrtaceae), a new ironbark species from the Hunter Valley of New South Wales, Australia. *Telopea* 23: 73-87.
- Bell, S.A.J. & Murray, M. (2013) *Flora and Fauna Monitoring at Condran, Muswellbrook LGA*. Unpublished Report to Bulga Surface Operations (Glencore). November 2013. Eastcoast Flora Survey & Forest Fauna Surveys Pty Ltd.

- Bell, S.A.J. & Nicolle, D. (2012) *Eucalyptus expressa* (Myrtaceae): a new and distinctive species from the sandstone ranges north-west of Sydney, New South Wales. *Telopea* 14: 69-76.
- Bell, S.A.J. & Nicolle, D. (2020) Glen Gallic Mallee (*Eucalyptus dealbata* subsp. *aperticola*, Myrtaceae), a new taxon from the sandstone escarpment of the Hunter Valley, New South Wales. *Telopea* 23: 141-150.
- Bell, S. & Sims, R. (2018) Extensive populations of *Dracophyllum macranthum* (Ericaceae) in Coorabakh NP suggest a review of threat status. *Australasian Plant Conservation* 27: 11-14.
- Bell, S.A.J. & Walsh, N. (2015) *Leionema lamprophyllum* subsp. *fractum* (Rutaceae); a new and highly restricted taxon from the Hunter Valley of New South Wales. *Telopea* 18: 505-512.
- Bell, S., Branwhite, B. & Driscoll, C. (2005) *Thelymitra 'adorata'* (Orchidaceae): population size and habitat of a highly restricted terrestrial orchid from the Central Coast of New South Wales. *The Orchadian* 15(1): 6-10.
- Bell, S.A.J., Murray, M. & Sims, R. (2020) *Flora and Fauna Monitoring at Condran VCA, Muswellbrook LGA: 2019 Results*. Unpublished Report to Bulga Surface Operations (Glencore). January 2020. Eastcoast Flora Survey & Forest Fauna Surveys Pty Ltd.
- Bell, S., Peake, T. & Driscoll, C. (2007) Dealing with taxonomic uncertainty in Weeping Myall *Acacia pendula* from the Hunter catchment, New South Wales. *Australasian Plant Conservation*. 16(1): 14-15.
- Bell, S., Rockley, C. & Llewellyn, A. (2019) *Flora of the Hunter Region: Endemic Trees and Larger Shrubs*. CSIRO Publishing and University of Newcastle, Australia.
- Bernhardt, P. (1995) Notes on the anthecology of *Pterostylis curta* (Orchidaceae) *Cunninghamia* 4(1): 1-8.
- Bernhardt, P. & Burns-Balogh, P. (1986) Observations of the floral biology of *Prasophyllum odoratum* (Orchidaceae, Spiranthoideae). *Plant Systematics and Evolution* 153: 65-76.
- Bishop, T. (2000) *Field Guide to the Orchids of New South Wales and Victoria*. University of New South Wales Press, Sydney. Second Edition.
- Bradshaw, C.J.A. (2012) Little left to lose: deforestation and forest degradation in Australia since European colonization. *Journal of Plant Ecology* 5(1): 109-120.
- Brundrett, M.C. (2006) Role of symbiotic relationships in Australian terrestrial orchid conservation. *Australasian Plant Conservation* 15(2): 2-7.
- Brundrett, M.C. (2016) Using vital statistics and core-habitat maps to manage critically endangered orchids in the Western Australian wheatbelt. *Australian Journal of Botany* 64: 51-64.
- Brundrett, M.C. (2019) A comprehensive study of orchid seed production relative to pollination traits, plant density and climate in an urban reserve in Western Australia. *Diversity* 11, 123; doi:10.3390/d11080123.

- Brundrett, M.C., Scade, A., Batty, A.L., Dixon, K.W. & Sivasithamparam, K. (2003) Development of in situ and ex situ seed baiting techniques to detect mycorrhizal fungi from terrestrial orchid habitats. *Mycological Research* 107: 1210-1220.
- Burley, T.M. (1962) The evolution of the agricultural pattern in the Hunter Valley, New South Wales, *Australian Geographer* 8(5): 221-231. DOI: 10.1080/00049186208702383.
- Burrows, G.E. (1999) A survey of 25 remnant vegetation sites in the South Western Slopes, New South Wales. *Cunninghamia* 6 (2): 283-314.
- Cameron, N., Jurskis, V., Peacock, R. & Wilson, R. (2014) *Submission on draft report (released June 2014) on active and adaptive cypress management in the Brigalow and Nandewar State Conservation Areas*. Submission by the Institute of Foresters of Australia, New South Wales Division. August 2014.
- Clarke, K.R. & Gorley, R.N. (2006) *PRIMER v6: User Manual/ Tutorial*. PRIMER-E: Plymouth.
- Clements, M.A. (1989) Catalogue of Australian Orchidaceae. *Australian Orchid Research* 1: 1-160.
- Colquhoun G.P., Hughes K.S., Deysing L., Ballard J.C., Folkes C.B, Phillips G., Troedson A.L. & Fitzherbert J.A. (2020) *New South Wales Seamless Geology dataset, version 2.0 [Digital Dataset]*. Geological Survey of New South Wales, Department of Regional NSW, Maitland.
- Cumberland Ecology (2014) *Warkworth Continuation 2014: Ecological Impact Assessment*. Unpublished Report to EMGA Mitchell McLennan. Report No. 14001RP1, June 2014.
- Cumberland Ecology (2015) *Biodiversity Assessment Report for Drayton South Coal Project*. A report prepared for Hansen Bailey.
- Cunningham, G., Mulham, W.E., Milthorpe, P.L. & Leigh, J.L. (2011) *Plants of Western New South Wales*. CSIRO Publications, Canberra.
- de Lacey, C., Bell, S., Chamberlain, S. & Bossard, K. (2012a) Habitat of the Leafless Tongue Orchid *Cryptostylis hunteriana* Nicholls throughout its known Australian distribution. *The Orchadian* 17(4): 162-174.
- de Lacey, C., Bell, S. & Chamberlain, S. (2012b) Habitat of the Leafless Tongue Orchid *Cryptostylis hunteriana* Nicholls throughout its known Australian distribution. *Australasian Plant Conservation* 20(4): 23-25.
- de Lacey, C., Bell, S., Chamberlain, S. & Bossard, K. (2013) Finding the leafless tongue orchid '*Cryptostylis hunteriana*' Nicholls. *Nature New South Wales* Vol. 57 (1) Autumn 2013: 24-25. [online].
- Department of Environment, Climate Change and Water (DECCW) (2010) *National Recovery Plan for Prasophyllum petilum*. Department of Environment and Climate Change and Water (NSW), Hurstville. Available at: <http://www.environment.gov.au/biodiversity/threatened/recovery-plans/national-recovery-plan-prasophyllum-petilum>.



- Department of Planning, Industry and Environment (2020) *Biodiversity Assessment Method*. Published by the New South Wales Environment, Energy and Science Department of Planning, Industry and Environment, October 2020.
- Duncan, M. & Moloney, P.D. (2018) Comparing wild and reintroduced populations of the threatened orchid *Diuris fragrantissima* (Orchidaceae) in south-eastern Australia. *Australian Journal of Botany* 66: 459-467.
- Duncan, M., Pritchard, A. & Coates, F. (2005) Major threats to endangered orchids of Victoria, Australia. *Selbyana* 26(1/2): 189-195. Retrieved from <http://www.jstor.org/stable/41760190>.
- Eco Logical Australia (2015) *MCO OC4 South-west modification: Flora and fauna impact assessment*. Unpublished Report to Moolarben Coal Operations. April 2015.
- Ecovision Consulting (2008) *Ecological Impact Assessment, Stage 2 of the Moolarben Coal Project, Ulan, New South Wales*. Unpublished Report to Moolarben Coal Mines Pty Limited. December 2008.
- Faast, R. & Facelli, J.M. (2009) Grazing orchids: impact of florivory on two species of *Caladenia* (Orchidaceae). *Australian Journal of Botany* 57(4): 361-372.
- Fensham, R.J., Laffineur, B., Collingwood, T.D., Beech, E., Bell, S., Hopper, S.D., Phillips, G., Rivers, M.C., Walsh, N. & White, M. (2020) Rarity or decline: Key concepts for the Red List of Australian eucalypts. *Biological Conservation* 243 108455-108455.
- FloraSearch (2014) *North Wambo Underground – Longwall 10A modification*. Unpublished Report to Wambo Coal Pty Limited. August 2014.
- Gallagher, R.V. (2016) Correlates of range size variation in the Australian seed-plant flora. *Journal of Biogeography* 43: 1287-1298.
- Gillman, M.P. & Dodd, M.E. (1998) The variability of orchid population size. *Botanical Journal of the Linnean Society* 126: 65-74.
- Hawkeswood, T.J. (2006) Comments on the orchid pollination review paper by P.B. Adams and S.D. Lawson (1993) in the Australian Journal of Botany and a listing of some more recent literature. *Calodema* 7: 1-4.
- Heinrich, A. & Dowling, W. (2000) *Rare and threatened plant surveys of Barrington Tops National Park – plateau area*. Unpublished reports prepared for the NSW National Parks and Wildlife Service, Hunter District.
- Herd, S. & Herd, D. (2005) ANOS Newcastle, Central Coast Groups lend a hand to the Anvil Hill Project Watch, Denman. *The Orchadian* 15(1): 18-21.
- Hosking, J.R. & James, T.A. (1998) Native and exotic flora of the north western slopes upstream of the junction of the Peel and Namoi Rivers, New South Wales. *Cunninghamia* 5(3): 721-766.

- Hunter, C. (2010) *People, Property, Power: Plashett, Jerrys Plains*. Anglo American Metallurgical Coal Pty Ltd: Brisbane.
- Hunter, J.T. (2008) *Vegetation and floristics of Warrabah National Park*. Unpublished Report to the New South Wales National Parks and Wildlife Service, July 2008. doi: 10.13140/RG.2.1.4446.0649.
- Hunter, J.T. (2010) *Vegetation and Floristics of Cubbo, Etoo and Dewsons Lease sections of the Pilliga SCA, Pilliga NP and Pilliga West SCA*. Unpublished Report to Department of Environment, Climate Change and Water. May 2010.
- Hunter Eco (2013) *Mt Arthur Coal Open Cut Modification Ecological Assessment*. Unpublished Report to BHP Billiton, January 2013.
- Hunter Eco (2019) *Maxwell Project Biodiversity Development Assessment Report*. Unpublished Report to Malabar Coal, July 2019.
- Hunter Eco (2020) *Maxwell Project Targeted Orchid Surveys*. Unpublished Report to Malabar Resources.
- Jacobs (2017) *Biodiversity Assessment Report: Hunter Water Corporation Proposed Biobank Site Irrawang Swamp – Grahamstown Dam*. Unpublished Report. August 2017.
- Jacquemyn, H., Duffy, K.J. & Selosse, M-A. (2017) Biogeography of Orchid Mycorrhizas. Chapter 8 IN L. Tedersoo (ed.) *Biogeography of Mycorrhizal Symbiosis* Ecological Studies 230. DOI 10.1007/978-3-319-56363-3\_8.
- Janes, J.K. & Duretto, M.F. (2010) A new classification for subtribe Pterostylidinae (Orchidaceae), reaffirming *Pterostylis* in the broad sense. *Australian Systematic Botany* 23: 260-269.
- Janes, J.K., Steane, D.A. & Vaillancourt, R.E. (2010) An investigation into the ecological requirements and niche partitioning of Pterostylidinae (Orchidaceae) species. *Australian Journal of Botany* 58(5): 335-341.
- Jeanes, J. (2015a) *Prasophyllum* aff. *petilum*. VicFlora Flora of Victoria. [Online]. Royal Botanic Gardens Melbourne. Available from: <http://data.rbg.vic.gov.au/vicflora/>.
- Jeanes, J. (2015b) *Diuris tricolor*. VicFlora Flora of Victoria. [Online]. Royal Botanic Gardens Melbourne. Available from: <http://data.rbg.vic.gov.au/vicflora/>.
- Jersáková, J. & Malinová, T. (2007) Spatial aspects of seed dispersal and seedling recruitment in orchids. *New Phytologist* 176: 237–241.
- Jones, D.L. (1991) New taxa of Australian Orchidaceae. *Australian Orchid Research* 2: 1-208.
- Jones, D.L. (1993) *Native Orchids of Australia*. Reed Australia.
- Jones, D.L. & Rouse, D.T. (2018) *Prasophyllum argillaceum* (Orchidaceae: Prasophyllinae), a new species in the *Prasophyllum campestre* R.J.Bates & D.L.Jones/ *Prasophyllum petilum* D.L.Jones & R.J.Bates complex. *Australian Orchid Review* 83(5): 50-53.

- Jusaitis, M. & Sorensen, B. (1993) Germination of *Pterostylis arenicola* – an endangered greenhood from South Australia. *The Orchadian* 11(1): 18-22.
- Kindlmann, P. & Balounova, Z. (2001) Irregular flowering patterns in terrestrial orchids: theories vs empirical data. *Web Ecology* 2: 75-82.
- Kleinfelder (2017) *Vegetation Mapping Report. Bayswater and Liddell Power Stations*. Unpublished Report to AGL Macquarie Pty Ltd. September 2017.
- Kovac, M. & Lawrie, J.W. (1991) *Soil Landscapes of the Singleton 1:250 000 Sheet*. Soil Conservation Service of New South Wales, Sydney.
- Kuiter, R.H. & Findlater-Smith, M. (2017) Overview of *Pterostylis* pollination (Orchidaceae) in Victoria. *Aquatic Photographics, Seaford* Short Paper 4, July 2017. Available at: [https://www.researchgate.net/publication/318282443\\_Overview\\_of\\_Pterostylis\\_Pollination\\_Orchidaceae\\_in\\_Victoria](https://www.researchgate.net/publication/318282443_Overview_of_Pterostylis_Pollination_Orchidaceae_in_Victoria).
- Light, M.H.S. & MacConnail, M. (2011) Potential impact of insect herbivores on orchid conservation. *European Journal of Environmental Sciences* 1(2): 115-124.
- McCormick, M.K. & Jacquemyn, H. (2014) What constrains the distribution of orchid populations? *New Phytologist* 202: 392–400.
- McCormick, M.K., Whigham, D.F. & Canchani-Viruet, A. (2018) Mycorrhizal fungi affect orchid distribution and population dynamics. *New Phytologist* 219: 1207-1215.
- McQualter, E., Cross, R., McLean, C.B. & Ladiges, P.Y. (2007) *Prasophyllum* and its associated mycorrhizal fungi. *Lankesteriana* 7(3): 497-501.
- Mickail, L.N.N., Bell, S.A.J. & Beranek, C.T. (2020) Dispersal potential in two restricted and five wide-ranging *Senecio* (Asteraceae) taxa from central eastern New South Wales, Australia. *Australian Journal of Botany* 68: 333-344.
- Nurfadilah, S., Swarts, N., Dixon, K., Lambers, H. & Merritt, D. (2013) Variation in nutrient-acquisition patterns by mycorrhizal fungi of rare and common orchids explains diversification in a global biodiversity hotspot. *Annals of Botany* 111(6): 1233-1241.
- Oja, J., Vahtra, J., Bahram, M., Kohout, P., Kull, T., Rannap, R., Koljalg, U. & Tedersoo, L. (2017) Local-scale spatial structure and community composition of orchid mycorrhizal fungi in semi-natural grasslands. *Mycorrhiza* 27: 355-367.
- Paget, A. (2008) *Results of searches for the Ruddy Hood Pterostylis chaetophora M.A.Clem. & D.L.Jones, syn Oligochaetochilus chaetophorus (M.A.Clem. & D.L.Jones), Spring 2008*. Unpublished Report, Hunter-Central Rivers Catchment Management Authority, Taree.
- Perry, T.M. (1955) The spread of rural settlement in New South Wales, 1788–1826. *Historical Studies: Australia and New Zealand* 6(24): 377-395. DOI: 10.1080/10314615508595010.

- Pfeifer, M., Wiegand, K., Heinrich, W. & Jetschke, G. (2006) Long-term demographic fluctuations in an orchid species driven by weather: implications for conservation planning. *Journal of Applied Ecology* 43: 313–324.
- Phillips, R.D., Peakall, R., Retter, B.A., Montgomery, K., Menz, M.H.M., Davis, B.J., Hayes, C., Brown, G.R., Swartz, N.D. & Dixon, K.W. (2015) Pollinator rarity as a threat to a plant with a specialized pollination system. *Botanical Journal of the Linnean Society* 179: 511–525.
- Phillips, R.D., Reiter, N. & Peakall, R. (2020) Orchid conservation: from theory to practice. *Annals of Botany* XX: 1-18. DOI: 10.1093/aob/mcaa093.
- Phillips, R.D., Scaccabarozzi, D., Retter, B.A., Hayes, C., Brown, G.R., Dixon, K.W. & Peakall, R. (2014) Caught in the act: pollination of sexually deceptive trap-flowers by fungus gnats in *Pterostylis* (Orchidaceae). *Annals of Botany* 113: 629–641.
- Reiter, N., Freestone, M., Brown, G. & Peakall, R. (2019) Pollination by sexual deception of fungus gnats (Keroplastidae and Mycetophilidae) in two clades of *Pterostylis* (Orchidaceae). *Botanical Journal of the Linnean Society* 190(1): 101–116.
- Rouse, D.T. (2002) *Report on the taxonomic, ecological and conservation status of taxa in the genus Prasophyllum R.Br. in south-eastern Australia*. Unpublished Report. May 2002.
- Scaccabarozzi, D., Cozzolino, S., Guzzetti, L., Galimberti, A., Milne, L., Dixon, K.W. & Phillips, R.D. (2018) Masquerading as pea plants: behavioural and morphological evidence for mimicry of multiple models in an Australian orchid. *Annals of Botany* 122(6): 1061–1073.
- Schiestl, F.P. (2005) On the success of a swindle: pollination by deception in orchids. *Naturwissenschaften* 92: 255–264.
- Seaton, P.T. (2007) Orchid conservation: Where do we go from here? *Lankestriana* 7: 13-16.
- Shefferson, R.P., Jacquemyn, H., Kull, T. & Hutchings, M.J. (2020) The demography of terrestrial orchids: life history, population dynamics and conservation. *Botanical Journal of the Linnean Society*. 192: 315–332.
- Smith, Z.F., James, E.A. & McLean, C.B. (2010) Mycorrhizal specificity of *Diuris fragrantissima* (Orchidaceae) and persistence in a reintroduced population. *Australian Journal of Botany* 58(2): 97-106.
- Stanley, T.D. & Ross, E.M. (1989) *Flora of South-eastern Queensland. Volume 3*. Queensland Department of Primary Industries, Brisbane.
- Su, Y-Y. & Guo, L-D. (2007) Arbuscular mycorrhizal fungi in non-grazed, restored and over-grazed grassland in the Inner Mongolia steppe. *Mycorrhiza* 17: 689-693.
- Thalwitzer, L., Kelly, D., Smissen, R.D., Butler, R., Suckling, D.M. & El-Sayed, A. (2018) Species-specific male pollinators found for three native New Zealand greenhood orchids (*Pterostylis* spp.) suggest pollination by sexual deception. *Australian Journal of Botany* 66: 243-254.

- Thompson, W.A. & Eldridge, D.J. (2005) White cypress pine (*Callitris glaucophylla*): a review of its roles in landscape and ecological processes in eastern Australia. *Australian Journal of Botany* 53: 555–570.
- Tierney, D.A., Sommerville, K.D., Tierney, K.E., Fatemi, M. & Gross, C.L. (2017) Trading populations—can biodiversity offsets effectively compensate for population losses? *Biodiversity and Conservation* 26: 2115-2131.
- Umwelt (2006) *Ecological Assessment Anvill Hill Project*. Report prepared for Centennial Hunter Pty Limited, June 2006.
- Umwelt (2010) *Ravensworth Operations Project Ecological Assessment*. Unpublished Report Prepared for Ravensworth Operations Pty Limited, February 2010.
- Umwelt (2011a) *Baseline ecological studies of potential biodiversity offset site – Highfields, near Manobalai, NSW*. Unpublished Report to Xstrata Coal. October 2011.
- Umwelt (2011b) *Baseline ecological studies of potential biodiversity offset site – Wybong Heights, near Manobalai, NSW*. Unpublished Report to Xstrata Coal. October 2011.
- Umwelt (2012) *Ecological values and condition of Northern and Western Corridors, as part of Biodiversity Offset Areas for Mangoola Mine*. Unpublished Report to Xstrata Mangoola Pty Limited. January 2012.
- Umwelt (2013) *Ecological Impact Statement: Bulga Coal Optimisation Project*. Unpublished Report to Bulga Coal Management Pty Ltd. April 2013.
- Umwelt (2014) *Ecological assessment; Mount Owen Continued Operations Project*. Unpublished Report on behalf of Glencore, October 2014.
- URS (2009) *Manildra – Parkes 132kV Transmission Line Project: Species Impact Statement*. Unpublished Report to TransGrid. July 2009.
- Vizer, C. (2013) *Contributions to the understanding of the ecology and biology of D. tricolor and Prasophyllum sp. Wybong in the Upper Hunter Region, NSW*. Unpublished BEnv.Sc. & Mgmt (Hons) thesis, University of Newcastle, May 2013.
- Vogel, S. (1973) Fungus gnat flowers and fungus mimesis. IN Brantjes, N.B.M. & Linskens, H.F. (eds.), *Pollination and Dispersal*. Nijmegen: Netherlands.
- Voyron, S., Ercole, E., Ghignone, S., Perotto, S. & Girlanda, M. (2016) Fine-scale spatial distribution of orchid mycorrhizal fungi in the soil of host-rich grasslands. *New Phytologist* 213: 1428-1439.
- Walker, K. (1997) *Natural history information on the presumed pollinator (Lasioglossum (Chilactus) lanarium) of the Small Golden Moth Orchid (Diuris sp. aff. lanceolata)*. Museum of Victoria unpublished report to the Department of Natural Resources and Environment, Victoria.
- Waterman, R.J. & Bidartondo, M.I. (2008) Deception above, deception below: linking pollination and mycorrhizal biology of orchids. *Journal of Experimental Botany* 59(5): 1085-1096.

- Waud, M., Busschaert, P., Lievens, B. & Jacquemyn, H. (2016) Specificity and localised distribution of mycorrhizal fungi in the soil may contribute to co-existence of orchid species. *Fungal Ecology* 20: 155-165.
- Wells, T.C.E., Rothery, P., Cox, R. & Bamford, S. (1998) Flowering dynamics of *Orchis morio* L. and *Herminium monorchis* (L.) R.Br. at two sites in eastern England. *Botanical Journal of the Linnean Society* 126: 39-48.
- Weston, P.H., Perkins, A.J. & Entwisle, T.J. (2005) More than symbioses: orchid ecology, with examples from the Sydney region. *Cunninghamia* 9: 1-15.
- Wildthing (2011) *Seven part test on threatened flora and fauna for a proposed subdivision at Lot 122 DP750924 Almond Street Denman, NSW*. Unpublished Report to Highfields NSW Pty Ltd. October 2012.
- Wilson, N., Seddon, J. and Baines, G. (2016) *Factors influencing the flowering of the Tarengo Leek Orchid (Prasophyllum petilum)*. Technical Report 36. Environment, Planning and Sustainable Development Directorate, ACT Government, Canberra. Available at: [https://www.environment.act.gov.au/data/assets/pdf\\_file/0018/1026342/TR36-Factors-influencing-the-flowering-of-the-Tarengo-Leek-Orchid.pdf](https://www.environment.act.gov.au/data/assets/pdf_file/0018/1026342/TR36-Factors-influencing-the-flowering-of-the-Tarengo-Leek-Orchid.pdf)
- Wraith, J. & Pickering, C. (2018) Quantifying anthropogenic threats to orchids using the IUCN Red List. *Ambio* 47: 307-317.
- Yare, B., Bell, S. & Hunter, N. (2020) Phenology of the threatened *Diuris praecox* (Orchidaceae), a range-restricted terrestrial orchid from central eastern New South Wales. *Cunninghamia* 20: 105-113.
- Zoete, T. (2000) Vegetation Survey of the Barrington Tops and Mount Royal National Parks for use in Fire Management. *Cunninghamia* 6(3): 511-578.

## Appendix 1 – Dr Stephen Bell DPIE Accreditation



DOC18/211176-1

[REDACTED]

Dear [REDACTED]

**Recognition of Dr Stephen Bell as an expert for *Diuris tricolor* and *Prasophyllum petilum* (Syn. *Prasophyllum* sp. 'Wybong')**

Thank you for your e-mail of the 9 April 2018 in which you request that Dr Stephen Bell be recognised as an expert for *Diuris tricolor* (Pine Donkey Orchid) and *Prasophyllum petilum* (Syn. *Prasophyllum* sp. 'Wybong') (Tarengo Leek Orchid) for the [REDACTED].

The Office of Environment and Heritage (OEH) has reviewed the information provided by you in support of this request. OEH is satisfied that Dr Stephen Bell satisfies the definition of species expert for *Diuris tricolor* and *Prasophyllum petilum* (Syn. *Prasophyllum* sp. 'Wybong'), in accordance with Section 6.5.2.3. of the Biodiversity Assessment Method.

Please note that this formal recognition of Dr Stephen Bell as an expert only applies to *Diuris tricolor* and *Prasophyllum petilum* (Syn. *Prasophyllum* sp. 'Wybong').

If you require any further information regarding this matter, please contact Steven Cox, Senior Team Leader, Planning Hunter Central Coast, on 4927 3140.

Yours sincerely

*Sharon Molloy 14/5/2018*

**SHARON MOLLOY**  
**Director Hunter Central Coast Branch**  
**Regional Operations Division**

Contact officer: STEVEN COX  
02 4927 3140

Locked Bag 1002 Dangar NSW 2309  
Level 4, 26 Honeysuckle Drive Newcastle  
NSW 2300  
ABN 30 841 387 271  
[www.environment.nsw.gov.au](http://www.environment.nsw.gov.au)



Planning,  
Industry &  
Environment

Our ref: DOC20/443557-1

Your ref: n/a

Dr Stephen Bell

Vegetation Ecologist  
East Coast Flora Survey  
[sajbell@bigpond.com](mailto:sajbell@bigpond.com)

Dear Dr Bell

**Approval of Dr Stephen Bell as an expert for *Pterostylis chaetophora***

I refer to your correspondence dated 9 June 2020 in which you requested that Dr Stephen Bell be considered as an expert for *Pterostylis chaetophora*.

The Biodiversity and Conservation Division (BCD) of the Department of Planning, Industry and Environment has reviewed the information provided in support of this request. BCD is satisfied that Dr Stephen Bell satisfies the definition of a species expert for *Pterostylis chaetophora* in accordance with Section 6.5.2.3 of the Biodiversity Assessment Method.

Please note that this formal recognition of Dr Stephen Bell as an expert only applies to the Hunter Central Coast Region of NSW which comprises the following local government areas; Central Coast, Lake Macquarie, Newcastle, Port Stephens, Mid Coast, Dungog, Maitland, Cessnock, Singleton, Muswellbrook and Upper Hunter.

If you have any further questions in relation to this matter, please contact Steve Lewer, Acting Senior Team Leader Planning, on 4927 3158 or via email at [rog.hcc@environment.nsw.gov.au](mailto:rog.hcc@environment.nsw.gov.au)

Yours sincerely

A handwritten signature in black ink that reads 'Joe Thompson'.

**JOE THOMPSON**  
Director Hunter Central Coast Branch  
Biodiversity and Conservation Division

Date: 25 June 2020



## Appendix 2 – Floristic Composition of Grassland Habitat (Bell 2012a)

The derivation of diagnostic species for each defined floristic group has been defined using the SIMPER routine in *PRIMER* on available full floristic plot data. SIMPER analysis provides the relative contributions of each species to the Bray-Curtis similarity within each of the defined vegetation communities. Only those species contributing to a total cumulative contribution of 99% of the average similarity (i.e. the value shown at the top of each floristic table) for each community are listed. These species can be described of as *typical* of that community, and have a consistently large presence within the data as reflected in the ratio of their contribution to the standard deviation (the Sim/SD field in each table) across the within-group similarities (the average similarity). Key canopy species are highlighted.

In the tables:

- Average similarity is the within-group similarity for all pairs of sample plots comprising the community. Higher average similarity indicates a better-defined community.
- Av.Abund is the average cover abundance of that species within sample plots comprising the community
- Av.Sim is the average similarity (contribution) made by each species to the within-group similarity (the overall average similarity).
- Sim/SD is the ratio of average similarity to standard deviation for each species across all pairs of samples. A high ratio represents a good discriminating species. At least three samples are required for this ratio to be calculated (not available for four communities).
- Contrib% is the percentage contribution of each species to the overall average similarity for the community.
- Cum.% is the cumulative percentage contribution of each species, up to a maximum of 99%.

### Unit 1a: *Dichanthium/ Sporobolus/ Chloris* Grassland - Key Diagnostic Species [based on 63 plots]:

Group 1a: *Dichanthium/ Sporobolus/ Chloris*

Average similarity: 45.72

Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
<i>Dichanthium sericeum</i> subsp. <i>sericeum</i>	2.92	2.68	1.09	5.87	12.92
<i>Senecio madagascariensis</i> *	1.89	2.58	3.58	5.64	18.56
<i>Sporobolus creber</i>	2.02	2.22	1.79	4.87	23.42
<i>Anagallis arvensis</i> *	1.75	2.13	1.86	4.66	28.09
<i>Chrysocephalum semipapposum</i>	1.71	1.92	1.48	4.20	32.29
<i>Centaurium tenuiflorum</i> *	1.67	1.88	1.40	4.10	36.39
<i>Bothriochloa decipiens</i> var. <i>decipiens</i>	2.02	1.82	1.06	3.98	40.37
<i>Glycine tabacina</i>	1.56	1.78	1.47	3.90	44.27
<i>Chloris truncata</i>	1.79	1.41	0.93	3.09	47.36
<i>Gamochoeta americana</i> *	1.38	1.38	1.04	3.02	50.39
<i>Cyclosporum leptophyllum</i> *	1.35	1.22	1.19	2.67	53.05
<i>Fimbristylis dichotoma</i>	1.30	1.21	0.88	2.66	55.71

<i>Aristida ramosa</i> var. <i>ramosa</i>	1.52	1.21	0.89	2.64	58.35
<i>Vittadinia muelleri</i>	1.41	1.20	0.84	2.63	60.99
<i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>	1.27	1.12	0.86	2.44	63.43
<i>Dichelachne micrantha</i>	1.59	1.09	0.76	2.38	65.80
<i>Vulpia muralis</i> *	1.38	1.05	0.77	2.30	68.10
<i>Hypochaeris radicata</i> *	1.21	0.90	0.73	1.97	70.08
<i>Trifolium arvense</i> *	0.97	0.83	0.93	1.81	71.88
<i>Petrorhagia dubia</i> *	1.08	0.81	0.73	1.78	73.66
<i>Asperula conferta</i>	1.06	0.78	0.68	1.70	75.36
<i>Plantago debilis</i>	1.03	0.77	0.67	1.69	77.05
<i>Hypochaeris microcephala</i> var. <i>albiflora</i> *	1.00	0.74	0.62	1.61	78.66
<i>Dichondra repens</i>	0.94	0.61	0.64	1.33	80.00
<i>Oxalis perenans</i>	0.94	0.61	0.61	1.33	81.33
<i>Carthamus lanatus</i> *	0.81	0.39	0.50	0.86	82.19
<i>Briza minor</i> *	0.76	0.38	0.46	0.84	83.02
<i>Eulalia aurea</i>	0.92	0.37	0.36	0.81	83.83
<i>Wahlenbergia communis</i>	0.62	0.35	0.54	0.77	84.61
<i>Convolvulus erubescens</i>	0.62	0.35	0.49	0.76	85.36
<i>Cymbopogon refractus</i>	0.63	0.31	0.46	0.68	86.04
<i>Daucus glochidiatus</i>	0.65	0.31	0.40	0.67	86.71
<i>Sida corrugata</i>	0.65	0.31	0.39	0.67	87.38
<i>Austrodanthonia tenuior</i>	0.65	0.30	0.36	0.65	88.03
<i>Polycarpon tetraphyllum</i> *	0.62	0.28	0.39	0.62	88.65
<i>Triptilodiscus pygmaeus</i>	0.62	0.28	0.33	0.62	89.27
<i>Calocephalus citreus</i>	0.78	0.27	0.33	0.58	89.85
<i>Brunoniella australis</i>	0.57	0.23	0.31	0.51	90.36

## Unit 2: *Aristida*/ *Cymbopogon* Grassland - Key Diagnostic Species [based on 44 plots]:

### Group 2: *Aristida*/ *Cymbopogon*

Average similarity: 39.82




Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
<i>Aristida ramosa</i> var. <i>ramosa</i>	3.43	4.60	2.17	11.55	11.55
<i>Linum trigynum</i> *	2.18	3.01	2.04	7.56	19.11
<i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>	2.07	2.84	2.01	7.14	26.25
<i>Anagallis arvensis</i> *	1.70	2.42	1.73	6.09	32.34
<i>Senecio madagascariensis</i> *	1.66	2.32	1.65	5.84	38.18
<i>Aristida vagans</i>	1.95	1.83	0.90	4.60	42.78
<i>Hypochaeris radicata</i> *	1.75	1.77	1.00	4.44	47.22
<i>Cymbopogon refractus</i>	1.48	1.73	1.19	4.35	51.58
<i>Glycine tabacina</i>	1.14	1.32	1.25	3.32	54.90
<i>Bothriochloa decipiens</i> var. <i>decipiens</i>	1.43	1.23	0.69	3.08	57.98
<i>Vulpia muralis</i> *	1.27	1.20	0.97	3.02	61.00
<i>Sporobolus creber</i>	1.14	0.99	0.68	2.48	63.48
<i>Briza minor</i> *	1.07	0.96	0.79	2.41	65.89
<i>Chrysocephalum apiculatum</i>	1.02	0.81	0.54	2.03	67.92
<i>Triptilodiscus pygmaeus</i>	0.84	0.58	0.50	1.47	69.39




<i>Vittadinia muelleri</i>	0.93	0.58	0.44	1.45	70.83
<i>Dichondra repens</i>	0.77	0.54	0.53	1.35	72.18
<i>Gamochaeta americana</i> *	0.80	0.53	0.52	1.34	73.52
<i>Dichelachne micrantha</i>	0.82	0.52	0.49	1.31	74.83
<i>Taraxacum officinale</i> *	0.80	0.50	0.43	1.26	76.08
<i>Lomandra confertifolia subsp. pallida</i>	0.75	0.48	0.53	1.21	77.30
<i>Tolpis barbata</i> *	0.77	0.46	0.44	1.16	78.46
<i>Lachnagrostis filiformis</i>	0.75	0.44	0.39	1.10	79.56
<i>Centaurium tenuiflorum</i> *	0.70	0.41	0.41	1.03	80.59
<i>Oxalis perenans</i>	0.68	0.39	0.41	0.97	81.56
<i>Richardia stellaris</i> *	0.66	0.38	0.41	0.94	82.51
<i>Chrysocephalum semipapposum</i>	0.77	0.37	0.38	0.94	83.44
<i>Fimbristylis dichotoma</i>	0.68	0.37	0.37	0.93	84.38
<i>Cyclospermum leptophyllum</i> *	0.66	0.36	0.44	0.90	85.27
<i>Petrorhagia dubia</i> *	0.68	0.35	0.37	0.88	86.15
<i>Asperula conferta</i>	0.59	0.31	0.35	0.77	86.93
<i>Sida corrugata</i>	0.57	0.30	0.39	0.75	87.67
<i>Linaria pelisseriana</i> *	0.57	0.25	0.33	0.64	88.31
<i>Glycine clandestina</i>	0.41	0.23	0.41	0.58	88.89
<i>Murdannia graminea</i>	0.50	0.21	0.31	0.53	89.42
<i>Centaurium erythraea</i> *	0.50	0.20	0.25	0.50	89.92




**Unit 4: Bothriochloa biloba/ Carthamus/ Danthonia Grassland - Key Diagnostic Species [based on 7 plots]:**

Group 4: Bothriochloa biloba/ Carthamus/ Danthonia					
Average similarity: 50.03					
Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
<i>Bothriochloa biloba</i>	5.14	13.03	5.61	26.04	26.04
<i>Carthamus lanatus</i> *	2.57	6.41	2.45	12.82	38.86
<i>Chloris truncata</i>	1.86	4.86	4.58	9.72	48.57
<i>Austrodanthonia tenuior</i>	2.14	4.54	1.32	9.08	57.65
<i>Einadia nutans subsp. linifolia</i>	1.71	4.16	3.83	8.31	65.97
<i>Lolium perenne</i> *	1.57	3.31	1.35	6.61	72.58
<i>Austrostipa aristiglumis</i>	1.57	2.20	0.74	4.40	76.97
<i>Vittadinia cuneata var. cuneata</i>	0.86	1.55	0.90	3.11	80.08
<i>Oxalis perenans</i>	1.14	1.34	0.62	2.68	82.76
<i>Senecio madagascariensis</i> *	0.86	1.22	0.92	2.43	85.19
<i>Sporobulus creber</i>	1.00	1.07	0.59	2.13	87.32
<i>Medicago truncatula</i> *	0.86	0.95	0.60	1.90	89.22
<i>Carex inversa</i>	0.86	0.92	0.58	1.84	91.05



### Appendix 3 – Biophysical Attributes of the Project Area














## Appendix 4 – Biophysical Attributes of Orchid Locations











## Appendix 5 – Resume: Dr Stephen Bell



### Principal

*Eastcoast Flora Survey*, PO Box 216 Kotara Fair NSW 2289

Telephone: (02) 4953 6523

Mobile: (0407) 284 240

e-mail: [sajbell@bigpond.com](mailto:sajbell@bigpond.com)

### Profile:

<http://www.stephenbell.com.au/>



### Conjoint Fellow

*School of Environmental & Life Sciences*, University of Newcastle,  
Callaghan NSW 2308

e-mail: [stephen.bell@newcastle.edu.au](mailto:stephen.bell@newcastle.edu.au)

### Profiles:

<http://www.newcastle.edu.au/profile/stephen-bell>

[https://www.researchgate.net/profile/Stephen\\_Bell10](https://www.researchgate.net/profile/Stephen_Bell10)

---

## PRÉCIS

Stephen has been involved in native vegetation survey, classification and mapping in the Greater Sydney and Hunter Regions since 1990. During this time, he has undertaken comprehensive surveys for the National Parks and Wildlife Service in over 30 conservation reserves, and has been contracted to the NSW Office of Environment & Heritage (OEH) as Senior Botanist and Team Leader for several large scale regional projects within the Sydney Basin bioregion. Under contract to local Councils, Stephen has co-ordinated and completed LGA-wide vegetation classification and mapping projects for Wyong, Gosford, Cessnock, Pittwater and Lake Macquarie LGAs, and has assisted in similar mapping projects for Blue Mountains LGA. Stephen has also completed several studies on Threatened Ecological Communities and threatened plant species, and published the results of some of these in the scientific literature.

On behalf of the Ecological Society of Australia, Stephen was the ecological expert on the Hunter Regional Vegetation Committee (2003), and from 2017 represents that organization on the NSW Threatened Species Scientific Committee (administering the *Biodiversity Conservation Act 2016*). Stephen was also a past member of the Hunter Threatened Flora Recovery Team, a founding member of the Hunter Rare Plants Committee (a sub-committee of the Hunter Region Botanic Gardens), and since 2014 has been a member of the OEH Species Technical Group which oversees management and expenditure of threatened species throughout NSW via its *Saving our Species* initiative. He is also often called upon by Government for advice regarding the significance of vegetation communities and plant species within the northern Sydney Basin bioregion, and has sat on numerous expert panels in this regard. Stephen has been called upon as an Expert Witness for several cases heard in the NSW Land and Environment Court, where his knowledge on the vegetation of the Sydney Basin bioregion has been used to argue contentious land-use decisions.

Stephen has published several scientific papers on various aspects of the vegetation of the Sydney Basin, including classifications of vegetation within conservation reserves, threatened and rare plant species, and the description of new plant taxa. Stephen has completed over 4500 standard full floristic sampling plots within the Sydney Basin, which are stored and used in vegetation classification analyses. Other skills include extensive multivariate data analysis experience, and GIS mapping. Stephen's PhD thesis, completed on a part-time basis through the University of Newcastle, presented improvements in the recognition, identification and classification of restricted and significant vegetation communities, such as Threatened Ecological Communities (TECs).

In October 1996, Stephen established *Eastcoast Flora Survey*, a specialist botanical consultancy providing high quality services to government and the private sector. Since June 2014, Stephen has also been a Conjoint Fellow in the School of Environmental & Life Sciences at the University of Newcastle (NSW), seeking to raise the output of ecological research on plants and vegetation within the Hunter region.

---

## ACADEMIC QUALIFICATIONS

Doctor of Philosophy (PhD), 2013	<i>Defining and mapping rare vegetation communities: Improving techniques to assist land-use planning and conservation</i> (University of Newcastle)
Bachelor of Science (Honours), 1991	<i>Effects of the weed Scotch Broom on bird communities in open forests on Barrington Tops</i> (University of Newcastle)
Bachelor of Science, 1989	<i>Majors in Geography and Biology</i> (University of Newcastle)

---

## EMPLOYMENT HISTORY

University of Newcastle	Conjoint Fellow (Plant Sciences Group)	June 2014 - Present
Eastcoast Flora Survey	Consultant Botanist (Principal)	Oct. 1996 - Present
Ecotone Ecological Consultants Pty Ltd	Manager - Flora Studies	Jan. 1996 - Oct. 1996
Private Ecological Consultant	Sole trader	Jan. 1991 - Dec. 1995
NSW National Parks and Wildlife Service	Project Officer	Sept. 1993 - Jan. 1994
University of Newcastle, Geography Dept.	Field Tutor (Scientific)	July 1993 - Aug. 1993
NSW National Parks and Wildlife Service	Project Officer	Jan. 1993 - June 1993
University of NSW, School of Biol. Sciences	Research Assistant (Bird ecology)	Sept. 1992 - Jan. 1993
NSW National Parks and Wildlife Service	Technical Officer (Scientific)	Jan. 1992 - June 1992
RZ Mines (Newcastle)	Environmental Research Officer	Oct. 1990 - Dec. 1991
Wayne Perry & Associates P/L	Environmental Officer (Casual)	June 1990 - Oct. 1990

---

## RESEARCH INTERESTS

- Vegetation classification and mapping, at local and regional scales
- Definition and mapping of rare and threatened vegetation communities
- Restoration of threatened grassy woodlands from derived grasslands
- Improving data sampling methods for monitoring and classification
- Re-constructing vegetation distribution using information from historical botanical explorers
- Population ecology and habitat of rare and threatened plants
- Taxonomy and significance of Hunter Region plants

---

## MINISTERIAL APPOINTMENTS

- Committee Member (ESA Rep.), NSW Threatened Species Scientific Committee (July 2017-present)
- Committee Member, NSW Species Technical Group, Flora (*Save Our Species Program*) (2014-present)
- Committee Member (ESA Rep.), Hunter Regional Vegetation Committee (2001-2003)

---

## CONFERENCE & WORKSHOP PRESENTATIONS

- Australian Plant Society (NSW) Annual Conference, August 2019, Newcastle: "*Endemic Plants of the Hunter Region: Trees and Larger Shrubs*".
- Best Practice Mine Rehabilitation Conference, September 2014, Singleton, NSW; The Tom Farrell Institute for the Environment, University of Newcastle: "*Effective Biodiversity Offsets: Improving planning, valuation and monitoring practice*" (with Martin Fallding).

- Plant Identification for Flora of the Hunter Valley, 7<sup>th</sup> - 8<sup>th</sup> April 2014, Kurri Kurri, Australian Network for Plant Conservation: “*Introduction to the flora of the Hunter Valley - history, diversity and ecology*”.
- HOTSPOTS Fire Project: Awabakal and Worimi Fire Forum, 27<sup>th</sup> July 2011, Williamstown, Never Never Resources: “*Vegetation of the Worimi Conservation Lands*”.
- HOTSPOTS Fire Project: Wanaruah Fire Forum, 17<sup>th</sup> – 19<sup>th</sup> August 2010, Sandy Hollow, Upper Hunter Valley, Nature Conservation Council: “*Vegetation of Wanaruah Lands, Sandy Hollow*”.
- Coastal Groundwater Dependent Ecosystems Workshop, 3<sup>rd</sup> – 4<sup>th</sup> September 2009, South West Rocks, NSW (Geoscience Australia): “*Surveying, classifying and mapping vegetation on the Tomago Sandbeds*”.
- Vegetation Management and Biodiversity Conservation in the Hunter Region, May 2000, Singleton, NSW (Hunter Environment Lobby Inc.): “*An evaluation of vegetation survey and threatened plant species listings in the Hunter Region*”

---

## PROFESSIONAL MEMBERSHIPS

- Ecological Society of Australia (ESA)
- Australian Network for Plant Conservation Inc. (ANPC)
- International Association for Vegetation Science (IAVS)
- International Association for Vegetation Science Vegetation Classification Working Group (IAVS VSWG)
- Australasian Systematic Botany Society (ASBS)

---

## PUBLICATION REVIEWER

- *Diversity* (MDPI, Switzerland)
- *Forests* (MDPI, Switzerland)
- *International Journal of Environmental Research and Public Health* (MDPI, Switzerland)
- *Journal of Vegetation Science* (International Association for Vegetation Science)
- *Pacific Conservation Biology* (CSIRO Publishing)
- *Resources* (MDPI, Switzerland)
- *Sustainability* (MDPI, Switzerland)
- *Telopea* (National Herbarium of New South Wales)
- *Vegetation Classification and Survey* (International Association for Vegetation Science)

---

## BOARD MEMBERSHIPS

- 2019 - *Vegetation Classification and Survey* (Editorial Board)
- 2019 - *Sustainability* (Review Board)
- 2019 – *Australian Network for Plant Conservation* (Committee Member)

---

## ACCREDITED BAM SPECIES EXPERT (NSW DPIE)

- *Cryptostylis hunteriana* (Orchidaceae)
- *Diuris praecox* (Orchidaceae)
- *Diuris tricolor* (Orchidaceae)
- *Hibbertia procumbens* (Dilleniaceae)
- *Prasophyllum petilum* (Orchidaceae)
- *Prostanthera junonis* (Lamiaceae)
- *Pterostylis chaetophora* (Orchidaceae)
- *Thelymitra adorata* (Orchidaceae)

## PUBLICATIONS (PEER REVIEWED)

- Bell, S.A.J. (submitted) Floristic community diversity in derived native grasslands: a case study from the upper Hunter Valley of New South Wales. *Cunninghamia* (submitted)
- Bell, S.A.J. (submitted) Successful recruitment following translocation of a threatened terrestrial orchid (*Diuris tricolor*) into mining rehabilitation in the Hunter Valley of NSW. *Ecological Management and Restoration* (submitted)
- Bell, S.A.J. (in prep) A strategy for assessing population size in threatened plant surveys using a classification of detectability based on key life-form traits, seasonality and disturbance response. *Diversity* (in prep)
- Bell, S.A.J. & Hillier, P. (2020) Targeted surveys of a poorly conserved threatened orchid (*Pterostylis chaetophora*) in Columbe National Park (Hunter Valley, NSW) reveal substantial populations and identify occupied habitat. *Cunninghamia* 20: 199-207.
- Bell, S.A.J. & Nicolle, D. (2020) Glen Gallic Mallee (*Eucalyptus dealbata* subsp. *aperticola*, Myrtaceae), a new taxon from the sandstone escarpment of the Hunter Valley, New South Wales. *Telopea* 23: 141-150.
- Bell, S.A.J. & Driscoll, C. (2020) Data-informed Sampling and Mapping: A new approach to ensure plot-based classifications locate, classify and map rare and restricted vegetation types. *Australian Journal of Botany* doi: 10.1071/bt20024
- Mickaill, L., Bell, S., & Beranek, C. (2020) Dispersal potential in two restricted and five wide-ranging *Senecio* (Asteraceae) taxa from central eastern New South Wales, Australia. *Australian Journal of Botany* 68: 333-344.
- Bell, S.A.J. (2020) Translocation of threatened terrestrial orchids into non-mined and post-mined lands in the upper Hunter Valley of New South Wales, Australia. *Restoration Ecology* 28: 1396-1407.
- Yare, B., Bell, S., & Hunter, N. (2020) Phenology of the threatened *Diuris praecox* (Orchidaceae), a range-restricted terrestrial orchid from central eastern New South Wales. *Cunninghamia* 20:105-113.
- Bell, S.A.J. & Klaphake, V. (2020) *Eucalyptus calidissima* (Myrtaceae), a new ironbark species from the Hunter Valley of New South Wales, Australia. *Telopea* 23: 73-87.
- Fensham, R., Laffineur, B., Collingwood, T., Beech, E., Bell, S., Hopper, S., Phillips, G., Rivers, M., Walsh, N. & White, M. (2020) Rarity or decline: Key concepts for the Red List of Australian eucalypts. *Biological Conservation* 243 108455
- Bell, S.A.J. (2019) Additions and amendments to the rare or threatened vascular plants of Wollemi National Park, central eastern New South Wales. *Cunninghamia* 19: 43-56.
- Bell, S.A.J. (2019) *Macrozamia flexuosa* C. Moore (Zamiaceae): a review of distribution, habitat and conservation status of this endemic cycad from the Hunter Region of New South Wales. *Cunninghamia* 19: 7-27.
- Bell, S., Rockley, C., & Llewellyn, A. (2019) *Flora of the Hunter Region: Endemic Trees and Larger Shrubs*. CSIRO Publishing. 136 pp. ISBN: 9781486311026
- DeLacey, C., Bell, S., Chamberlain, S., & Bossard, K. (in review) Prediction of and realised habitat for a cryptic plant species: the Leafless Tongue Orchid *Cryptostylis hunteriana* Nicholls. *Cunninghamia* (in review)
- Bell, S.A.J. (2018) Fate of a rare flowering event in a population of the endangered *Acacia pendula* (Weeping Myall) from the Hunter Valley of New South Wales. *Cunninghamia* 18: 79-88.
- Bell, S.A.J. & Driscoll, C. (2017) *Acacia wollarensis* (Fabaceae, Mimosoideae sect. Botrycephalae), a distinctive new species endemic to the Hunter Valley of New South Wales, Australia. *Telopea* 20: 125-136.
- Bell, S.A.J. & Driscoll, C. (2016) Hunter Valley Weeping Myall Woodland – is it really definable and defensible with and without Weeping Myall (*Acacia pendula*)? *Cunninghamia* 16: 15-30.

- Bell, S.A.J. & Walsh, N. (2015) *Leionema lamprophyllum* subsp. *fractum* (Rutaceae); a new and highly restricted taxon from the Hunter Valley of New South Wales. *Telopea* 18: 505-512.
- Bell, S.A.J. & Driscoll, C. (2014) *Acacia pendula* (Weeping Myall) in the Hunter Valley of New South Wales: early explorers' journals, database records and habitat assessments raise doubts over naturally occurring populations. *Cunninghamia* 14: 179-200.
- Bell, S.A.J. & Nicolle, D. (2012) *Eucalyptus expressa* (Myrtaceae): a new and distinctive species from the sandstone ranges north-west of Sydney, New South Wales. *Telopea* 14: 69-76.
- Bell, S.A.J. & Stables, M. (2012) Floristic variability, distribution and an extension of range for the endangered Pittwater Spotted Gum Forest, Central Coast, New South Wales. *Cunninghamia* 12(2): 143-152.
- Bell, S.A.J. (2009) Vegetation and floristics of Columbey National Park, lower Hunter Valley, New South Wales. *Cunninghamia* 11(2): 241-275.
- Bell, S.A.J. (2008) Rare or threatened vascular plant species of Wollemi National Park, central eastern New South Wales. *Cunninghamia* 10(3): 331-371.
- Bell, S., Branwhite, B., & Driscoll, C. (2005) *Thelymitra 'adorata'* (Orchidaceae): population size and habitat of a highly restricted terrestrial orchid from the Central Coast of New South Wales. *The Orchadian* 15(1): 6-10.
- Bell, S.A.J. (2004) Distribution and habitat of the vulnerable tree species, *Angophora inopina* (Myrtaceae), on the Central Coast of New South Wales. *Cunninghamia* 8(4): 477-484.
- Bell, S.A.J. (2004) Vegetation of Werakata National Park, Hunter Valley, New South Wales. *Cunninghamia* 8(3): 331-347.
- Bell, S.A.J. & Copeland, L.M. (2004) *Commersonia rosea* (Malvaceae s.l.: Lasiopetaleae): a new, rare fire-ephemeral species from the upper Hunter Valley, New South Wales. *Telopea* 10(2): 581-587.
- Bell, S.A.J. (2002) Habitat of the endangered *Hibbertia procumbens* (Labill.) DC (Dilleniaceae) from the Central Coast of New South Wales. *Victorian Naturalist* 119(2): 69-74.
- Bell, S.A.J. (2001) Notes on population size and habitat of the vulnerable *Cryptostylis hunteriana* Nicholls (Orchidaceae) from the Central Coast of New South Wales. *Cunninghamia* 7(2): 195-204.
- Bell, S.A.J. (2001). Notes on the distribution and conservation status of some restricted plant species from sandstone environments of the upper Hunter Valley, New South Wales. *Cunninghamia* 7(1): 77-88.
- Bell, S. (2000) An evaluation of vegetation survey and threatened plant species listings in the Hunter Region. Pp. 19-34 IN *Vegetation Management and Biodiversity Conservation in the Hunter Region - Where to from here?* Ed. by M.Fallding. Proceedings of the Public Workshop. Hunter Environment Lobby. Singleton, 12 May 2000.

---

## PUBLICATIONS (OTHERS)

- Bell, S. (in prep) Voucher specimens vs observation records: why 'collecting' is better than 'observing' for plant conservation. *Australasian Plant Conservation* (in prep)
- Bell, S. (2020) Check those IDs... the importance of confirming threatened plant identifications before implementing management. *Australasian Plant Conservation* 28: 18-20.
- Bell, S. (2019) Translocation 'success' is all about detection: experiences with two threatened orchids from the Hunter Valley of NSW. *Australasian Plant Conservation* 28: 27-31.
- Bell, S. (2018) The responsibilities of ecological consultants in disseminating outcomes from threatened species surveys. *Australasian Plant Conservation* 27: 3-6.
- Bell, S. & Sims, R. (2018) Extensive populations of *Dracophyllum macranthum* (Ericaceae) in Coorabakh NP suggest a review of threat status. *Australasian Plant Conservation* 27: 11-14.
- Bell S.A.J. & Kodala P.G. (2018) *Acacia wollarensis*. In: *Flora of Australia*. Australian Biological Resources Study, Department of the Environment and Energy, Canberra. <https://profiles.ala.org.au/opus/foa/profile/Acacia%20wollarensis>
- Bell, S. (2017) New insights into the ecology of the critically endangered *Banksia conferta* (Proteaceae) from the mid-north coast of NSW. *Australasian Plant Conservation* 26(1): 15-18.

- Bell, S. & Holzinger, B. (2015) Wildfire reveals new populations of the endangered *Commersonia rosea* and *Monotaxis macrophylla* in northern Wollemi National Park, NSW. *Australasian Plant Conservation* 23: 2-4.
- Bell, S. & Elliott, M. (2013) Preliminary results suggest fire is required to maintain *Acacia dangarensis*, a threatened single-population endemic from the Hunter Valley of NSW. *Australasian Plant Conservation* 22(1): 9-10.
- de Lacey, C, Bell, S, Chamberlain, S. & Bossard, K. (2013) Finding the leafless tongue orchid '*Cryptostylis hunteriana*' Nicholls. *Nature New South Wales* Vol. 57 (1) Autumn 2013: 24-25. [online]
- de Lacey, C., Bell, S., & Chamberlain, S. (2012) Habitat of the Leafless Tongue Orchid *Cryptostylis hunteriana* Nicholls throughout its known Australian distribution. *Australasian Plant Conservation* 20(4): 23-25.
- de Lacey, C., Bell, S., Chamberlain, S., & Bossard, K. (2012) Habitat of the Leafless Tongue Orchid *Cryptostylis hunteriana* Nicholls throughout its known Australian distribution. *The Orchadian* 17(4): 162-174.
- Bell, S.A.J. (2010) Defining and mapping an endangered ecological community within Lake Macquarie Local Government Area, New South Wales. *Australasian Plant Conservation* 18(3): 18-19.
- Bell, S., Peake, T. & Driscoll, C. (2007) Dealing with taxonomic uncertainty in Weeping Myall *Acacia pendula* from the Hunter catchment, New South Wales. *Australasian Plant Conservation*. 16(1): 14-15.
- Bell, S. & Driscoll, C. (2005) New records of the endangered *Hibbertia procumbens* from the Central Coast of NSW. *Australasian Plant Conservation* 13(4): 24-25.
- Bell, S.A.J., Parsons, J., & Meldrum, R. (2005) Towards the protection and management of hanging swamps on the Somersby Plateau, Central Coast, New South Wales. *Australasian Plant Conservation* 13(3): 10-11.
- Bell, S. (2003) Another new and highly restricted mallee from the Hunter Valley, *Eucalyptus castrensis*. *Hunter Flora* 11: 2.
- Peake, T., Bell, S., Tame, T., Simpson, J., & Curran, T. (2003) *The Hunter Rare Plants Database: Identification and listing of regionally significant flora for the Hunter Region, New South Wales*. Poster Presentation at the Ecological Society of Australia Annual Conference 2003, Armidale NSW.
- Peake, T., Bell, S., Tame, T., Simpson, J., & Curran, T. (2002) *Warkworth Sands Woodland – An Endangered Ecological Community: Distribution, Ecological Significance and Conservation Status*. Hunter Region Botanic Gardens Technical Paper [www.huntergardens.org.au/]
- Bell, S. (2002) Plant profile: The Leafless Tongue Orchid, *Cryptostylis hunteriana*. *Hunter Flora* 9: 2.

---

## SELECTED UNPUBLISHED TECHNICAL REPORTS (1993 – 2020)

- Bell, S.A.J. (2020) *Survey and monitoring of the Vulnerable Pterostylis chaetophora (Rusty Greenhood) in the Lower Hunter Valley, NSW: 2019 Results*. Unpublished Report to NSW Office of Environment and Heritage. June 2020. Eastcoast Flora Survey.
- Bell, S.A.J. (2020) *Expert Report - Expected Presence of Threatened Terrestrial Orchids (Diuris tricolor & Prasophyllum petilum): Bayswater Water and Other Associated Operational Works Project*. Unpublished Report to Kleinfelder. May 2020. Eastcoast Flora Survey.
- Bell, S.A.J. & Oppen, M. (2020) *Management of habitat for the Critically Endangered Persoonia pauciflora (Proteaceae) at 'Georgia' (Lot 56 DP755211), North Rothbury, NSW*. Unpublished Report to NSW Biodiversity Conservation Trust. April 2020. Eastcoast Flora Survey.
- Bell, S.A.J. (2020) *Identifying potential pollinators in the critically endangered Banksia conferta (Proteaceae), Coorabakh National Park, NSW*. Unpublished Report to NSW National Parks and Wildlife Service. March 2020. Eastcoast Flora Survey.
- Bell, S.A.J. (2019) *Expert Report: Hibbertia procumbens (Labill.) DC. (Dilleniaceae) at a Proposed Industrial Warehouse Development, Gindurra Road, Somersby*. Unpublished Report to Anderson Environment & Planning.
- Bell, S.A.J. (2019) *A Re-assessment of Foreshore Vegetation along Tuggerah Lake, Chittaway Bay to The Entrance, Central Coast LGA*. Report to Central Coast Council. November 2019.



- Bell, S.A.J. (2019) *Baseline surveys for two threatened mallees (Eucalyptus castrensis and Eucalyptus pumila) in the lower Hunter Valley of New South Wales*. Unpublished Report to NSW Office of Environment and Heritage. April 2019. Eastcoast Flora Survey.
- Bell, S.A.J. (2019) *Peer Review: Assessment of the vulnerable Tetratheca juncea at a proposed development site at Gwandalan, Central Coast Council*. Unpublished Report to Resolve Urban Planning. April 2019. Eastcoast Flora Survey.
- Bell, S.A.J. (2019) *Survey and monitoring of the Vulnerable Pterostylis chaetophora (Rusty Greenhood) in the Lower Hunter Valley, NSW: 2018 Results*. Unpublished Report to NSW Office of Environment and Heritage. April 2019. Eastcoast Flora Survey.
- Bell, S.A.J. (2019) *A Revised Interim Vegetation Classification of the Central Coast Local Government Area*. Unpublished Report to Central Coast Council. July 2019. Eastcoast Flora Survey.
- Bell, S.A.J. (2019) *Verification Survey of the Critically Endangered Pomaderris reperta (Denman Pomaderris) at Aarons Pass Road, Central Tablelands, NSW*. Unpublished Report to Office of Environment & Heritage. February 2019. Eastcoast Flora Survey. [peer review survey, *Pomaderris cotoneaster* rather than *Pomaderris reperta*]
- Bell, S.A.J. (2019) *Monitoring of translocated threatened orchids (Diuris tricolor, Prasophyllum petilum) at Mangoola Coal: 2018 Results*. Unpublished Report to Mangoola Coal. February 2019. Eastcoast Flora Survey.
- Bell, S.A.J. (2018) *Review of significant vegetation at 100 Bakali Road, Forresters Beach (Lot 9, DP 8857)*. Unpublished Report to Darcy Smith. November 2018. Eastcoast Flora Survey.
- Bell, S.A.J. (2018) *Monitoring of the endangered Pterostylis gibbosa (Orchidaceae) at Milbrodale, Hunter Valley: Year 3 Results*. Unpublished Report to Office of Environment & Heritage. November 2018. Eastcoast Flora Survey.
- Bell, S.A.J. (2018) *Monitoring of Acacia pendula in the Weeping Myall Management Areas, Bulga Coal, Singleton LGA: 2018 Results*. Unpublished Report to Bulga Coal. October 2018. Eastcoast Flora Survey.
- Bell, S.A.J. (2018) *Monitoring and survey of the Critically Endangered Pomaderris reperta (Rhamnaceae) and Vulnerable Lasiopetalum longistamineum (Malvaceae), Upper Hunter Valley, New South Wales: 2017-2018 (Year 2) Results*. Unpublished Report to Office of Environment & Heritage. October 2018. Eastcoast Flora Survey.
- Bell, S.A.J. (2018) *Mt Dangar Multi-species Saving our Species Project: Year 2 (2017-2018) Summary Report*. Unpublished Report to Office of Environment & Heritage. October 2018. Eastcoast Flora Survey.
- Bell, S.A.J. (2018) *Monitoring of Acacia pendula flowering in the Weeping Myall Management Area, Singleton LGA*. Unpublished Report to Bulga Coal. August 2018. Eastcoast Flora Survey.
- Bell, S.A.J. (2018) *Vegetation and floristics of the Huntlee Offset Lands, Hunter Valley, NSW: Sweetwater (North Rothbury)*. Unpublished Report to Office of Environment & Heritage. June 2018. Eastcoast Flora Survey.
- Bell, S.A.J. & Sims, R. (2018) *Targeted Surveys for the Vulnerable Dracophyllum macranthum (Ericaceae) in Coorabakh National Park, Lansdowne Plateau, NSW*. Unpublished Report to NSW Office of Environment & Heritage. 21 June 2018. Eastcoast Flora Survey.
- Bell, S.A.J. (2018) *Baseline demography of a significant new population of the Critically Endangered Banksia conferta (Proteaceae), Coorabakh National Park, NSW*. Unpublished Report to NSW Office of Environment & Heritage. June 2018. Eastcoast Flora Survey. Final Report.
- Bell, S.A.J. (2018) *Vegetation and floristics of the Huntlee Offset Lands, Hunter Valley, NSW: 4. Warrawalong (Cessnock LGA)*. Unpublished Report to Office of Environment & Heritage. April 2018. Eastcoast Flora Survey.
- Bell, S.A.J. (2018) *Vegetation and floristics of the Huntlee Offset Lands, Hunter Valley, NSW: 3. Corrabare (Cessnock LGA)*. Unpublished Report to Office of Environment & Heritage. April 2018. Eastcoast Flora Survey.
- Bell, S.A.J. (2018) *Vegetation and floristics of the Huntlee Offset Lands, Hunter Valley, NSW: 2. Elderslie (Singleton LGA)*. Unpublished Report to Office of Environment & Heritage. April 2018. Eastcoast Flora Survey.
- Bell, S.A.J. (2018) *Monitoring of translocated threatened orchids (Diuris tricolor, Prasophyllum petilum) at Mangoola Coal: 2017 Results*. Unpublished Report to Mangoola Coal. February 2018.

- Bell, S.A.J. (2017) *Monitoring of the endangered Pterostylis gibbosa (Orchidaceae) at Milbrodale, Hunter Valley: Year 2 Results*. Unpublished Report to NSW Office of Environment & Heritage. November 2017. Eastcoast Flora Survey.
- Bell, S.A.J. (2017) *A strategy for assessing population size in threatened plant surveys*. Unpublished Report to NSW Office of Environment and Heritage. November 2017.
- Bell, S.A.J. (2017) *Targeted survey for the threatened Diuris tricolor at Persoonia Park, North Rothbury, Hunter Valley*. Unpublished Report to Office of Environment & Heritage. November 2017. Eastcoast Flora Survey.
- Bell, S.A.J. (2017) *Baseline monitoring and survey of the vulnerable Pterostylis chaetophora (Orchidaceae) at North Rothbury, Hunter Valley, NSW*. Unpublished Report to NSW Office of Environment & Heritage. October 2017. Eastcoast Flora Survey.
- Bell, S.A.J. (2017) *Vegetation and floristics of the Huntlee Offset Lands, Hunter Valley, NSW: 1. Cedar Creek (Cessnock LGA)*. Unpublished Report to NSW Office of Environment and Heritage. October 2017.
- Bell, S.A.J. (2017) *Baseline monitoring and survey of the Critically Endangered Pomaderris reperta (Rhamnaceae) and Vulnerable Lasiopetalum longistamineum (Malvaceae), Upper Hunter Valley, New South Wales*. Unpublished Report to NSW Office of Environment and Heritage. October 2017.
- Bell, S.A.J. (2017) *Field Validation of Wetlands Mapping on the Central Coast of New South Wales*. Unpublished Report to NSW Office of Environment and Heritage.
- Bell, S.A.J. (2017) *Mt Dangar Multi-species Saving our Species Project: Year 1 (2016-2017) Summary Report*. Unpublished Report to NSW National Parks and Wildlife Service. June 2017.
- Bell, S.A.J. (2017) *Monitoring of the Critically Endangered Banksia conferta (Proteaceae) at Coorabakh National Park, NSW*. Unpublished Report to NSW Office of Environment and Heritage.
- Bell S.A.J. (2016) *Monitoring of the endangered Pterostylis gibbosa (Orchidaceae) at Milbrodale, Hunter Valley, NSW*. Unpublished Report to NSW Office of Environment and Heritage.
- Bell S.A.J. (2016) *Distribution and condition of the threatened Melaleuca biconvexa (Myrtaceae) on the Central Coast of NSW*. Unpublished Report to NSW Office of Environment and Heritage.
- Bell, S.A.J. (2016) *Additional notes on distribution and fragmentation of Warkworth Sands Woodland, from the Hunter Valley of New South Wales*. Unpublished Report to Department of Environment, Ecological Communities Section, Protected Species & Communities Branch, Canberra. April 2016.
- Bell, S.A.J. (2016) *An assessment of the vegetation occurring on coastal sands at Pelican Flats, Lake Macquarie LGA*. Lake Macquarie Research Grant Project 2004/2005-01.
- Bell, S.A.J. & Driscoll, C. (2016) *Volume 1: Vegetation Mapping Report, Lake Macquarie Local Government Area. Stages 1 - 6*. Unpublished Report to Lake Macquarie City Council. March 2016. Eastcoast Flora Survey.
- Bell, S.A.J. (2016) *Volume 2: Vegetation Community Profiles, Lake Macquarie Local Government Area. Working Draft v2*. Unpublished Report to Lake Macquarie City Council. March 2016. Eastcoast Flora Survey.
- Bell, S.A.J. (2016) *Vegetation and floristics of the Columbey Conservation Lands (NP & SCA), lower Hunter Valley, New South Wales*. Unpublished Report to Office of Environment & Heritage (Gloucester). January 2016. Eastcoast Flora Survey.
- Bell, S.A.J. (2015) *Review of Offset Multipliers (Tg values) for Selected Threatened Species*. Unpublished Report to NSW Office of Environment and Heritage. November 2015. Eastcoast Flora Survey.
- Bell, S.A.J. (2015) *Distribution, habitat and conservation status of Macrozamia flexuosa (Zamiaceae) in Lake Macquarie LGA and the lower Hunter Valley of New South Wales*. Lake Macquarie Research Grant Project 1999/2000-02.
- Bell, S.A.J. (2015) *External Review: Draft Threatened Plant Survey Guidelines*. Unpublished Report to Office of Environment & Heritage, NSW Department of Premier and Cabinet. May 2015. Eastcoast Flora Survey.
- Bell, S.A.J. (2015) *Lower Hunter Spotted Gum-Ironbark Forest: Verification Survey, Proposed Newcastle Inner City Bypass (Rankin Park to Jesmond), Newcastle LGA*. Unpublished Report to Roads and Maritime Services. March 2015. Eastcoast Flora Survey.

- Bell, S.A.J. (2014) *Tablelands Snow Gum TEC: Field Survey & Classification Analysis, Southern Tablelands, New South Wales. Tallaganda & Badja State Forest Trial Areas*. Unpublished DRAFT Report to NSW Office of Environment & Heritage. Eastcoast Flora Survey. November 2014.
- Bell, S.A.J. (2014) *Preliminary data analysis of Slaty Gum (Eucalyptus dawsonii) Forests & Woodlands in the Hunter Valley*. Unpublished Report to Department of Environment, Ecological Communities Section, Protected Species & Communities Branch, Canberra. August 2014. Prepared as part of deliberations emanating from the Expert Panel assessing the proposed Central Hunter Valley Eucalyptus Forest and Woodland Complex endangered ecological community.
- Bell, S.A.J. (2014) *Submission on Warkworth Sands Woodland EEC: Warkworth / Mt Thorley Continuation Project*. Unpublished Submission to NSW Department of Planning and Environment. Eastcoast Flora Survey. July 2014.
- Bell, S.A.J. (2013) *Ecological studies on Acacia dangarensis: Baseline data to inform management*. Unpublished Report to NSW Office of Environment & Heritage, Mudgee. October 2013. Eastcoast Flora Survey.
- Bell, S.A.J. (2013) *Assessment of Alluvial Riparian Blackbutt Forest in Wyong LGA against River Flat Eucalypt Forest EEC*. Unpublished Report to Wyong Shire Council. June 2013. Eastcoast Flora Survey.
- Bell, S.A.J. (2013) *A strategy for monitoring revegetation of rehabilitation after coal mining, Ravensworth Operations, Hunter Valley*. Unpublished Report to Ravensworth Operations, Xstrata Coal, April 2013. Eastcoast Flora Survey.
- Bell, S.A.J. (2013) *Summary Report of Baseline Pomaderris reperta Monitoring Data, Mangoola Coal*. Unpublished Report to Mangoola Coal. Eastcoast Flora Survey.
- Bell, S.A.J. (2012) *Update of Listing Advice and Fact Sheet for Weeping Myall - Coobah - Scrub Wilga Woodland/Low Forest of the Hunter Valley*. Unpublished Report to the Commonwealth Department of Sustainability, Environment, Water, Populations and Communities. Eastcoast Flora Survey, June 2012.
- Bell, S.A.J. (2012) *Comment on potential Blue Gum High Forest CEEC at Jesmond Park, Newcastle LGA*. Report to Newcastle City Council. Eastcoast Flora Survey. April 2012.
- Bell, S.A.J. & Carty, A. (2012) *Vegetation mapping of the Singleton Military Area*. Unpublished report to Commonwealth Department of Defence. Eastcoast Flora Survey & SKM, March 2012.
- Bell, S.A.J. (2011) *Implementing the NSW Vegetation Classification and Assessment (VCA) Scheme in the Hunter-Central Rivers Region: Attribution of Priority Vegetation Types*. Unpublished Database to Hunter-Central Rivers Catchment Management Authority. Eastcoast Flora Survey.
- Bell, S.A.J. (2011) *Review of Greater Hunter Vegetation Mapping: Version 0.1*. Unpublished Report to Office of Environment & Heritage, Department of Premier & Cabinet. August 2011. Eastcoast Flora Survey.
- Bell, S.A.J. (2011) *Diversity and condition of derived grasslands at Mangoola Coal, Upper Hunter Valley: Stage 2 – 2010 Surveys*. Unpublished Report to Mangoola Coal, August 2011. Eastcoast Flora Survey.
- DeLacey, C., Bell, S.A.J., & Chamberlain, S. (2011) *Pittwater Vegetation Classification, Vegetation Mapping, pre-1750 Vegetation Mapping and Vegetation Profiles*. Unpublished Report prepared for Pittwater Council.
- DeLacey, C., Bell, S., Chamberlain, S., & Bossard, K. (2010) *Prediction of habitat for cryptic plant species: The Leafless Tongue Orchid *Cryptostylis hunteriana* Nicholls as a case study*. A research project funded by the NSW Environmental Trust (2006/RR/0003). October 2010.
- Bell, S.A.J. (2010) *Lower Hunter Spotted Gum-Ironbark Forest EEC in the Warnervale area, Wyong LGA*. Unpublished Report to Wyong Shire Council, September 2010. Eastcoast Flora Survey.
- Bell, S.A.J. (2010) *Vegetation and floristics of Columbey National Park, lower Hunter Valley, New South Wales: Update with Duns Creek addition*. Unpublished Report to Department of Environment & Climate Change (Gloucester Area). June 2010. Eastcoast Flora Survey.
- Bell, S.A.J. (2010) *Lower Hunter Spotted Gum – Ironbark Forest: Distribution and composition in Cessnock, Maitland, Newcastle & Wyong LGAs*. Report to Department of Environment & Climate Change (Coffs Harbour). Eastcoast Flora Survey.

- Bell, S.A.J. & Driscoll, C. (2010) *Vegetation of the Worimi Conservation Lands, Port Stephens, New South Wales: Worimi NP, Worimi SCA & Worimi RP*. Unpublished Report to Department of Environment, Climate Change & Water, November 2010. Eastcoast Flora Survey.
- Bell, S.A.J. & Copeland, L. (2010) *A strategy for the translocation of threatened terrestrial orchids at Mangoola Coal, Upper Hunter Valley*. Unpublished Report to Mangoola Coal, September 2010. Eastcoast Flora Survey.
- Bell, S.A.J. (2009) *Lower Hunter Spotted Gum – Ironbark Forest: Distribution and composition in Lake Macquarie LGA*. Unpublished Draft Report to Lake Macquarie City Council. Eastcoast Flora Survey. May 2009.
- Bell, S.A.J. (2009) *Equivalence analysis of Hunter and Central Coast vegetation communities for the Hunter VCA Project*. Report to EcoLogical Australia & Hunter-Central Rivers Catchment Management Authority. Eastcoast Flora Survey.
- Bell, S.A.J. (2009) *The natural vegetation of the Gosford local government area, Central Coast, New South Wales. Volumes 1 & 2. Fully Revised*. Unpublished Report to Gosford City Council. 2009. Eastcoast Flora Survey.
- Bell, S.A.J. (2009) *Vegetation and floristics of Murrurundi Pass and Crawney Pass National Parks, upper Hunter Valley, New South Wales*. Unpublished Report to Department of Environment & Climate Change (Scone Area). February 2009.
- Bell, S.A.J. & Driscoll, C. (2009) *Vegetation survey and mapping of Sugarloaf State Conservation Area, Lake Macquarie*. Unpublished report and map to Department of Environment & Climate Change.
- Driscoll, C. & Bell, S.A.J. (2009) *The experimental translocation of *Tetratheca juncea* Sm. (Tremandraceae) at Gwandalan, Wyong Shire. Final Report*. Unpublished Report to Crighton Properties Pty Ltd, Wyong Shire Council, and Department of Environment and Climate Change. April 2009.
- Bell, S.A.J. (2008) *Review of flora issues relating to proposed Coal & Allied development on the Wallarah Peninsula*. Unpublished report to Department of Environment & Climate Change, Newcastle. March 2008.
- Bell, S. & Driscoll, C. (2008) *Revised Vegetation Mapping of Wyong LGA: Stage 1 – West of F3 Freeway*. Unpublished report and map to Wyong Shire Council.
- Bell, S.A.J. (2007) *The Vegetation of Eraring Power Station, Lake Macquarie, New South Wales*. Unpublished Report to Eraring Energy and HLA ENSR. Eastcoast Flora Survey.
- Bell, S. & Driscoll, C. (2007) *Vegetation of the Cessnock-Kurri Region, Cessnock LGA, New South Wales: Survey, Classification and Mapping*. Unpublished report and map prepared for Department of Environment and Conservation.
- Bell, S.A.J. & Driscoll, C. (2006) *Vegetation of the Tomago and Anna Bay Sandbeds, Port Stephens, New South Wales: Management of Groundwater Dependent Ecosystems. Part 1 – Vegetation Classification*. Unpublished Report to Hunter Water. Eastcoast Flora Survey. September 2006.
- Bell, S.A.J. & Driscoll, C. (2006) *Vegetation of the Salt Ash Air Weapons Range, Medowie, New South Wales*. Unpublished Final Report to Department of Defence. May 2006. Eastcoast Flora Survey.
- Bell, S.A.J. & Driscoll, C. (2006) *Vegetation Mapping of Watagans National Park and Jilliby State Conservation Area*. Summary Report to Parks & Wildlife Division, Department of Environment and Conservation. January 2006.
- Bell, S.A.J. (2006) *Eucalyptus parramattensis subsp. decadens: Status, distribution and habitat*. Unpublished Report prepared for the Department of Environment & Conservation, Newcastle. Eastcoast Flora Survey. June 2006.
- Bell, S.A.J. (2006) *Review of floristic and structural vegetation baseline data for Biometrics, Hunter & Central Rivers CMA*. Department of Environment & Conservation/ CSIRO.
- Bell, S.A.J. & Driscoll, C. (2005) *Assessment of littoral rainforest and coastal headland vegetation, Treachery Head, Myall Lakes National Park*. Unpublished Report to Parks and Wildlife Division, Department of Environment and Conservation. Eastcoast Flora Survey. March 2005.
- Bell, S.A.J. (2004) *Umina Coastal Sandplain Woodland: A regional analysis with reference to the vegetation present in Hillview Street, Woy Woy*. Unpublished Report to Gosford City Council. November 2004.

- Bell, S.A.J. (2004) *Data audit of vegetation survey within the Central Coast LGA's: 2000 - 2004*. Eastcoast Flora Survey - Report to Hunter Councils. July 2004.
- Bell, S.A.J. (2004) *The vegetation of the Hunter Economic Zone (HEZ), Cessnock LGA, New South Wales*. Unpublished Report to Harper Somers O'Sullivan. January 2004. Eastcoast Flora Survey.
- Driscoll, C., & Bell, S. (2004) *A taxonomic review of the Grevillea linearifolia complex as it occurs in the Central and Lower North Coast of NSW, with particular reference to the threatened Grevillea parviflora subsp. parviflora*. Unpublished Report to Lake Macquarie City Council, December 2004.
- Bell, S.A.J. & Driscoll, C. (2004) *Population count and habitat assessment of Rutidosis heterogama (Asteraceae), Lower Hunter and Central Coast*. Unpublished Report to Wyong Shire Council, April 2004. Eastcoast Flora Survey.
- Douglas, S.D. & Bell, S.A.J. (2003) *Vegetation survey and mapping of Clyde River National Park, NSW South Coast*. Unpublished Final Report to NSW NPWS South Coast Region. ESP Ecological Surveys & Planning P/L & Eastcoast Flora Survey. April 2003.
- Douglas, S.J. & Bell, S.A.J. (2003) *Vegetation survey and mapping, Bimberamala National Park, NSW South Coast*. Unpublished Final Report to NSW National Parks and Wildlife Service, South Coast Region. ESP Ecological Surveys & Planning P/L & Eastcoast Flora Survey. January 2003.
- Bell, S.A.J. (2002) *The natural vegetation of the Wyong Local Government Area, Central Coast, New South Wales. Parts 1 & 2*. Report to Wyong Shire Council. Eastcoast Flora Survey.
- Bell, S.A.J. & Driscoll, C. (2002) *Population size and habitat of the endangered Acacia bynoeana Benth. (Fabaceae: Mimosoideae) in Wyong Shire*. Unpublished Report to Wyong Shire Council. September 2002. Eastcoast Flora Survey.
- Bell, S.A.J. & Fallding, M. (2002) *Tomago Sandbeds: Guidelines for weed and bush fire management*. Final Draft Report to Department of Land and Water Conservation. Unpublished. Eastcoast Flora Survey.
- Peake, T., Bell, S., Tame, T., Simpson, J., & Curran, T. (2002) *Warkworth Sands Woodland – An Endangered Ecological Community: Distribution, Ecological Significance and Conservation Status*. Hunter Region Botanic Gardens Technical Paper [www.huntergardens.org.au/]
- Murray, M., Bell, S., & Hoye, G. (2001) *Lower Hunter and Central Coast Flora and Fauna Survey Guidelines. Version 1.0*. Prepared for the Lower Hunter and Central Coast Regional Environmental Management Strategy. November 2001.
- Bell, S.A.J. (2001) *Survey and Assessment of Angophora inopina in the Lower Hunter and Central Coast*. Eastcoast Flora Survey - Report to NSW National Parks and Wildlife Service, Threatened Species Unit, Central Directorate. June 2001.
- Bell, S.A.J. (2001) *Distribution, conservation & management of the vulnerable Angophora inopina. technical report and conservation management plan*. Eastcoast Flora Survey - Final Draft Report to Wyong Shire Council.
- Bell, S.A.J. & Murray, M. (2001) *The Ecological Significance of Bow Wow Creek Gorge, Mulbring, lower Hunter Valley, New South Wales: A Nationally Significant Site*. Unpublished Report to Cessnock City Council. Eastcoast Flora Survey & Forest Fauna Surveys Pty Ltd. June 2001.
- Douglas, S. & Bell, S. (2001) *Native vegetation mapping of Areas 1 to 5 in Blue Mountains City local government area*. Unpublished Report to Blue Mountains City Council. ESP Ecological Surveys & Planning Pty Ltd.
- Hill, L., Peake, T., Bell, S., & Raine, A. (2001) *The vegetation of Towarri National Park, Cedar Brush Nature Reserve & Wingen Maid Nature Reserve for fire management purposes*. Unpublished Report to NSW National Parks and Wildlife Service, Hunter District. Eastcoast Flora Survey.
- Bell, S.A.J. (1998) *Popran National Park vegetation survey. A fire management document. Volumes 1 - 4*. Eastcoast Flora Survey - Report to NSW National Parks and Wildlife Service (Central Coast Districts). December 1998.
- Bell, S.A.J. (1998) *Glenrock SRA and Awabakal NR vegetation survey. A fire management document. Volumes 1 & 2*. Eastcoast Flora Survey - Report to NSW National Parks and Wildlife Service (Hunter District). August 1998.

- Bell, S.A.J. (1998) *Lake Macquarie SRA, Pulbah Island NR, and Tingira Heights NR vegetation survey. A fire management document*. Volumes 1 & 2. Eastcoast Flora Survey - Report to NSW National Parks and Wildlife Service (Hunter and Central Coast Districts). April 1998.
- Bell, S.A.J. (1998) *Wollemi National Park vegetation survey. A fire management document*. Volumes 1 & 2. Eastcoast Flora Survey - Report to NSW National Parks and Wildlife Service (Upper Hunter District).
- Bell, S.A.J. (1997) *Vegetation Survey and Mapping of Crown Land, South of Manobalai Nature Reserve, Upper Hunter Valley*. Report to the Department of Land and Water Conservation and the NSW National Parks and Wildlife Service Upper Hunter District.
- Bell, S.A.J. (1997) *Tomaree National Park vegetation survey. A fire management document*. Eastcoast Flora Survey - Report to NSW National Parks and Wildlife Service (Hunter District). May 1997.
- Bell, S. & Russell, D. (1993) *A brief survey of the vegetation of the Hunter Valley floor. Report for Inquiry into Proposed Open-cut Mining in Part of Ravensworth State Forest, Near Singleton*. NSW National Parks and Wildlife Service.
- Bell, S., Vollmer, J. & Gellie, N. (1993) *Yengo National Park and Parr State Recreation Area. Vegetation Survey for Use in Fire Management*. Report prepared for NSW National Parks and Wildlife Service. Unpublished.

---

### **NSW LAND & ENVIRONMENT COURT (EXPERT WITNESS)**

- Bell, S.A.J. (2016) *Review of Biodiversity Issues: United Wambo Open Cut Coal Mine Project*. Unpublished Report to EDO NSW, September 2016.
- Bell, S.A.J. (2012) *Expert Report: Bulga Milbrodale Progress Association v Minister for Planning and Infrastructure and Warkworth Mining Limited*. Land and Environment Court Proceedings No: 10224 of 2012. Unpublished Report to EDO NSW, July 2012.
- Bell, S.A.J. (2009) *Affidavit: Assessment of vegetation at Lots 3 & 4 (DP399581) Quorrobolong, from field data collected in May 2006*. Unpublished Report to Department of Environment, Climate Change & Water, November 2009. Eastcoast Flora Survey.
- Bell, S.A.J. (2009) *Expert Report: Colongra Swamp Nature Reserve*. Unpublished Report to Department of Environment & Climate Change. Eastcoast Flora Survey. June 2009.
- Bell, S.A.J. (2006) *Expert Report: Providence Projects Pty Ltd v Gosford City Council*. Land and Environment Court Proceedings No: 11626 of 2004; 10101 of 2005. Unpublished Report to Gosford City Council.
- Bell, S.A.J. (2005) *Assessment of vegetation, 37 Laycock Street Carey Bay: Lake Macquarie City Council ats. First Cape Management Pty Ltd. L & E Court Proceedings 11475/04*. Expert Report to Land & Environment Court: April 2005.
- Bell, S.A.J. (2001) *Expert Report: Hunter Resort v Cessnock City Council*. Land and Environment Court Proceedings.

## Appendix 6 – Endorsement: Dr Lachlan Copeland



5<sup>th</sup> April, 2018

### Reference for orchid expertise of Dr Stephen Bell

#### **To whom it may concern**

This letter serves to introduce Dr Stephen Bell of Eastcoast Flora Survey as worthy to fill the role of 'expert' for matters concerning the threatened terrestrial orchids *Diuris tricolor* and *Prasophyllum petilum*. I myself have an extensive knowledge of the orchid flora of New South Wales, Victoria and Queensland gained over several decades of field survey. With Gary Backhouse, Robert Bates and Andrew Brown, I have co-authored the "Checklist of the Orchids of Australia including its Island Territories". In separate publications I have also described 15 new orchid species. I am personally very familiar with *Diuris tricolor* and *Prasophyllum petilum* from throughout their known distributional ranges, and have worked with Stephen on these two species previously.

Stephen has a wealth of experience and professional ability in botany and threatened species management, having worked in the consultancy field since the mid 1990's. In 1996, he established Eastcoast Flora Survey, a small consultancy focusing on vegetation surveys, classification, mapping and threatened species research and management. Over a 20 year period, Stephen has shown himself to be dedicated to presenting an accurate portrayal of the distribution and abundance of native plant species, particularly threatened plant species and ecological communities. He has researched a number of threatened plant species, including several orchids, and has published regularly in the scientific literature on these. In the past, I have personally approached Stephen on numerous occasions to discuss the distribution, abundance and threats of a wide variety of threatened plant species including several orchid taxa.

I first worked with Stephen on surveys for *Diuris* and *Prasophyllum* in the upper Hunter Valley in 2009, and since that time I am aware that he has undertaken annual surveys for these species across a number of offset and development sites, including overseeing and conducting a large translocation program at one mine site requiring detailed monitoring of individual plants, and the monitoring of natural populations elsewhere. Over a nine year period of surveying and monitoring for these species, Stephen is clearly highly regarded as an expert for these taxa in the Hunter Valley. I believe that Stephen would have personally seen more individuals of these two species and know their ecology more intimately than anyone else.

Given the experience detailed above, I have no hesitation in recommending Stephen as an expert in any matters concerning *Diuris tricolor* and *Prasophyllum petilum*, particularly with regard to habitats and status within the Hunter Valley of NSW.

Yours sincerely

A handwritten signature in black ink, appearing to read 'Lachlan Copeland'.

Dr Lachlan Copeland  
Senior Botanist,  
Eco Logical Australia

ATTACHMENT D  
THREATENED FLORA AND FAUNA SPECIES KNOWN OR PREDICTED TO OCCUR IN THE  
LOCALITY



**Table D-1  
Threatened Flora and Fauna Species Known or Predicted to occur in the Locality**

Common Name	Scientific Name	Conservation Status			Database Records				Recorded in Previous Studies and/or Recent Surveys <sup>8</sup>
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>	Potentially Associated with PCTs in the Biodiversity Assessment Development Footprint <sup>4</sup>	EPBC Act Protected Matters Search <sup>5</sup>	BioNet Atlas <sup>6</sup>	ALA <sup>7</sup>	
<b>Flora</b>									
White-flowered Wax Plant	<i>Cynanchum elegans</i>	E	E	S	Yes	Predicted	Yes	-	-
-	<i>Ozothamnus tessellatus</i>	V	V	S	Yes	-	Yes	Yes	-
Large-leafed Monotaxis	<i>Monotaxis macrophylla</i>	-	E	S	Yes	-	-	-	-
Acacia pendula population in the Hunter Catchment	<i>Acacia pendula</i> – endangered population	-	E	S	Yes	-	Yes	-	K*
Singleton Mint Bush	<i>Prostanthera cineolifera</i>	V	V	S	Yes	-	-	-	-
Wollemi Mint-bush	<i>Prostanthera cryptandroides</i> subsp. <i>cryptandroides</i>	V	V	S	Yes	Predicted	-	-	-
Slaty Red Gum	<i>Eucalyptus glaucina</i>	V	V	S	Yes	Predicted	Yes	-	-
Leafless Tongue-orchid	<i>Cryptostylis hunteriana</i>	V	V	S	Yes	Predicted	-	-	-
<i>Cymbidium canaliculatum</i> in the Hunter Catchment	<i>Cymbidium canaliculatum</i> – endangered population	-	E	S	Yes		Yes	-	K, L
Pine Donkey Orchid	<i>Diuris tricolor</i>	-	V	S	Yes	-	Yes	-	K
Pine Donkey Orchid population in the Muswellbrook local government area	<i>Diuris tricolor</i> – endangered population	-	E	S	Yes	-	Yes	-	K
Tarengo Leek Orchid	<i>Prasophyllum petilum</i> (sp. Wybong)	E	E	S	Yes	Predicted	-	-	-
-	<i>Pterostylis chaetophora</i>	-	V	S	Yes	-	-	-	-
Bodalla Pomaderris	<i>Pomaderris bodalla</i>	-	V	S	Yes	-	-	-	-

Common Name	Scientific Name	Conservation Status			Database Records				Recorded in Previous Studies and/or Recent Surveys <sup>8</sup>
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>	Potentially Associated with PCTs in the Biodiversity Assessment Development Footprint <sup>4</sup>	EPBC Act Protected Matters Search <sup>5</sup>	BioNet Atlas <sup>6</sup>	ALA <sup>7</sup>	
Scant Pomaderris	<i>Pomaderris queenslandica</i>	-	E	S	Yes	-	-	-	-
Denman Pomaderris	<i>Pomaderris reperta</i>	CE	CE	S	Yes	-	-	-	-
Austral Toadflax	<i>Thesium australe</i>	V	V	S	-	Predicted	-	-	-
<b>Amphibians</b>									
Green and Golden Bell Frog	<i>Litoria aurea</i>	V	E	S	Yes	Predicted	Yes	Yes	-
Booroolong Frog	<i>Litoria booroolongensis</i>	E	E	S	-	Predicted	-	-	-
Green-thighed Frog	<i>Litoria brevipalmata</i>	-	V	S	Yes	-	-	-	-
<b>Reptiles</b>									
Pink-tailed Legless Lizard	<i>Aprasia parapulchella</i>	V	V	S	-	-	-	-	M
Striped Legless Lizard	<i>Delma impar</i>	V	V	S	Yes	-	-	-	M
Pale-headed Snake	<i>Hoplocephalus bitorquatus</i>	-	V	S	Yes	-	-	-	-
<b>Birds</b>									
Freckled Duck	<i>Stictonetta naevosa</i>	-	V	E	-	-	-	Yes	-
Australasian Bittern	<i>Botaurus poiciloptilus</i>	E	E	E	-	Predicted	-	-	-
Black Falcon	<i>Falco subniger</i>	-	V	E	-	-	-	Yes	-
Square-tailed Kite	<i>Lophoictinia isura</i>	-	V	S/E	Yes	-	Yes	Yes	M
White-bellied Sea-Eagle	<i>Haliaeetus leucogaster</i>	MA	V	S/E	Yes	-	Yes	Yes	M
Spotted Harrier	<i>Circus assimilis</i>	-	V	E	Yes	-	Yes	Yes	A, M
Red Goshawk	<i>Erythrotriorchis radiatus</i>	V	CE	S	-	Predicted	-	-	-
Little Eagle	<i>Hieraaetus morphnoides</i>	-	V	S/E	Yes	-	Yes	Yes	A, I
Bush Stone-curlew	<i>Burhinus grallarius</i>	-	E	S	Yes	-	-	Yes	-
Australian Painted Snipe	<i>Rostratula australis</i>	E	E	E	-	Predicted	-	-	-

Common Name	Scientific Name	Conservation Status			Database Records				Recorded in Previous Studies and/or Recent Surveys <sup>8</sup>
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>	Potentially Associated with PCTs in the Biodiversity Assessment Development Footprint <sup>4</sup>	EPBC Act Protected Matters Search <sup>5</sup>	BioNet Atlas <sup>6</sup>	ALA <sup>7</sup>	
Eastern Curlew	<i>Numenius madagascariensis</i>	CE	-	S/E	-	Predicted	-	-	-
Curlew Sandpiper	<i>Calidris ferruginea</i>	CE	E	S/E	-	Predicted	-	-	-
Glossy Black-Cockatoo	<i>Calyptorhynchus lathami</i>	-	V	S/E	Yes	-	-	-	M
Gang-gang Cockatoo	<i>Callocephalon fimbriatum</i>	-	V	S/E	Yes	-	-	Yes	-
Little Lorikeet	<i>Glossopsitta pusilla</i>	-	V	E	Yes	-	Yes	Yes	J, M
Turquoise Parrot	<i>Neophema pulchella</i>	-	V	E	Yes	-	-	Yes	-
Swift Parrot	<i>Lathamus discolor</i>	CE	E	S/E	Yes	Predicted	-	-	A
Eastern Grass Owl	<i>Tyto longimembris</i>	-	V	E	Yes	-	-	-	-
Masked Owl	<i>Tyto novaehollandiae</i>	-	V	S/E	Yes	-	-	-	-
Powerful Owl	<i>Ninox strenua</i>	-	V	S/E	Yes	-	Yes	Yes	-
Barking Owl	<i>Ninox connivens</i>	-	V	S/E	Yes	-	Yes	Yes	B
Brown Treecreeper (eastern subspecies)	<i>Climacteris picumnus victoriae</i>	-	V	E	Yes	-	Yes	Yes	A, M
Speckled Warbler	<i>Chthonicola sagittata</i>	-	V	E	Yes	-	Yes	Yes	A, M
Black-chinned Honeyeater (eastern subspecies)	<i>Meliphreptus gularis gularis</i>	-	V	E	Yes	-	Yes	-	M
Regent Honeyeater	<i>Anthochaera phrygia</i>	CE	CE	S/E	Yes	Predicted	-	-	-
Painted Honeyeater	<i>Grantiella picta</i>	V	V	E	Yes	Predicted	-	-	-M
Hooded Robin (south-eastern form)	<i>Melanodryas cucullata cucullata</i>	-	V	E	Yes	-	Yes	-	-
Flame Robin	<i>Petroica phoenicea</i>	-	V	E	Yes	-	-	Yes	M
Scarlet Robin	<i>Petroica boodang</i>	-	V	E	Yes	-	Yes	-	A, M

Common Name	Scientific Name	Conservation Status			Database Records				Recorded in Previous Studies and/or Recent Surveys <sup>8</sup>
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>	Potentially Associated with PCTs in the Biodiversity Assessment Development Footprint <sup>4</sup>	EPBC Act Protected Matters Search <sup>5</sup>	BioNet Atlas <sup>6</sup>	ALA <sup>7</sup>	
Grey-crowned Babbler (eastern subspecies)	<i>Pomatostomus temporalis temporalis</i>	-	V	E	Yes	-	Yes	-	A, M
Varied Sittella	<i>Daphoenositta chrysoptera</i>	-	V	E	Yes	-	Yes	Yes	C, M
Dusky Woodswallow	<i>Artamus cyanopterus cyanopterus</i>	-	V	E	Yes	-	Yes	Yes	M
Diamond Firetail	<i>Stagonopleura guttata</i>	-	V	E	Yes	-	Yes	Yes	A, B, J
<b>Mammals</b>									
Spotted-tailed Quoll	<i>Dasyurus maculatus maculatus (south-eastern mainland population)</i>	E	V	E	Yes	Predicted	Yes	Yes	D, E
Brush-tailed Phascogale	<i>Phascogale tapoatafa</i>	-	V	S	Yes	-	Yes	-	-
Common Planigale	<i>Planigale maculata</i>	-	V	S	Yes	-	-	-	-
Koala	<i>Phascolarctos cinereus</i>	V	V	S/E	Yes	Predicted	Yes	-	-
Eastern Pygmy-possum	<i>Cercartetus nanus</i>	-	V	S	Yes	-	-	-	-
Yellow-bellied Glider	<i>Petaurus australis</i>	-	V	E	Yes	-	-	-	-
Squirrel Glider	<i>Petaurus norfolcensis</i>	-	V	S	Yes	-	Yes	-	A, D, E, F, J, M
Greater Glider	<i>Petauroides volans</i>	V	-	S	-	Predicted	-	-	-
Brush-tailed Rock-wallaby	<i>Petrogale penicillata</i>	V	E	S	Yes	Predicted	Yes	-	-
Grey-headed Flying-fox	<i>Pteropus poliocephalus</i>	V	V	S/E	Yes	Predicted	Yes	-	J, M
Yellow-bellied Sheath-tail-bat	<i>Saccolaimus flaviventris</i>	-	V	E	Yes	-	Yes	-	A, G, J, M
Eastern Coastal Free-tailed Bat	<i>Micronomus norfolkensis</i>	-	V	E	Yes	-	Yes	-	A, B, C, E, G, J, M
Northern Freetail-bat	<i>Mormopterus lumsdenae</i>	-	V	E	No	-	-	-	G
Little Bentwing-bat	<i>Miniopterus australis</i>	-	V	S/E	Yes	-	Yes	-	G, M

Common Name	Scientific Name	Conservation Status			Database Records				Recorded in Previous Studies and/or Recent Surveys <sup>8</sup>
		EPBC Act <sup>1</sup>	BC Act <sup>2</sup>	Credit Class <sup>3</sup>	Potentially Associated with PCTs in the Biodiversity Assessment Development Footprint <sup>4</sup>	EPBC Act Protected Matters Search <sup>5</sup>	BioNet Atlas <sup>6</sup>	ALA <sup>7</sup>	
Large Bentwing-bat	<i>Miniopterus orianae oceanensis</i>	-	V	S/E	Yes	-	Yes	-	A, C, D, E, F, G, H, J, M
Corben's Long-eared Bat	<i>Nyctophilus corbeni</i>	V	V	E	Yes	Predicted	Yes	-	B
Large-eared Pied Bat	<i>Chalinolobus dwyeri</i>	V	V	S	Yes	Predicted	Yes	-	A, C, G, M
Eastern False Pipistrelle	<i>Falsistrellus tasmaniensis</i>	-	V	E	Yes	-	Yes	-	E, F
Southern Myotis	<i>Myotis macropus</i>	-	V	S	Yes	-	Yes	-	A, B, G, M
Greater Broad-nosed Bat	<i>Scoteanax rueppellii</i>	-	V	E	Yes	-	Yes	-	B, D, E, J
Eastern Cave Bat	<i>Vespadelus troughtoni</i>	-	V	S	Yes	-	Yes	-	A, G, J
New Holland Mouse	<i>Pseudomys novaehollandiae</i>	V	-	E	-	Predicted	-	-	-

Shaded species are species with records in the locality.

<sup>1</sup> Conservation status under the EPBC Act (current as at August 2021). V = Vulnerable; E = Endangered; CE = Critically Endangered; MA = Migratory.

<sup>2</sup> Conservation status under the BC Act (current as at August 2021). V = Vulnerable; E = Endangered; CE = Critically Endangered.

<sup>3</sup> Biodiversity credit class under the TBDC (DPIE 2021a) (current as at August 2021). E = Ecosystem; S = Species.

<sup>4</sup> DPIE (2021a).

<sup>5</sup> DEE (2018a).

<sup>6</sup> DPIE (2021c).

<sup>7</sup> Atlas of Living Australia (2018).

<sup>8</sup> A – Cumberland Ecology (2009a) and/or Cumberland Ecology (2012).

B – Ecotone (2000).

C – Eco Logical Australia (2015).

D – Eco Logical Australia (2016a).

E – Eco Logical Australia (2016b).

F – Eco Logical Australia (2014).

G – Eco Logical Australia (2017).

H – Umwelt Environmental Consultants (Umwelt) (2006).

I – Umwelt (2007).

J – Hansen Bailey (2007).

K – Cumberland Ecology (2015).

L = Hunter Eco (2019).

M = Future Ecology (2019) (Attachment A).

\* Note that the location of the *Acacia pendula* reported by Cumberland Ecology (2015) was re-surveyed by Hunter Eco (2019) and was found to be *Acacia melvillei*.

ATTACHMENT E  
BAM BIODIVERSITY CREDIT REPORT - BIODIVERSITY ASSESSMENT DEVELOPMENT  
FOOTPRINT

## Proposal Details

Assessment Id	Proposal Name	BAM data last updated *
00026727/BAAS17004/21/00026728	Malabar Coal Maxwell Mine Entry Area	10/06/2021
Assessor Name	Report Created	BAM Data version *
Colin Driscoll	05/08/2021	45
Assessor Number	BAM Case Status	Date Finalised
BAAS17004	Finalised	05/08/2021
Assessment Revision	Assessment Type	
0	Major Projects	

\* Disclaimer: BAM data last updated may indicate either complete or partial update of the BAM calculator database. BAM calculator database may not be completely aligned with Bionet.

## Ecosystem credits for plant communities types (PCT), ecological communities & threatened species habitat

Zone	Vegetation zone name	TEC name	Current Vegetation integrity score	Change in Vegetation integrity (loss / gain)	Area (ha)	BC Act Listing status	EPBC Act listing status	Species sensitivity to gain class (for BRW)	Biodiversity risk weighting	Potential SAI	Ecosystem credits
<b>Bull Oak grassy woodland of the central Hunter Valley</b>											
2	1692_Mod erate_good	Not a TEC	48.5	48.5	0.5			High Sensitivity to Potential Gain	1.75		11
										<b>Subtotal</b>	<b>11</b>

## White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter

1	1606_Derived_native_grass	White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highla	31.4	31.4	3.8	Critically Endangered Ecological Community	Critically Endangered	High Sensitivity to Potential Gain	2.50	TRUE	75
										<b>Subtotal</b>	<b>75</b>
										<b>Total</b>	<b>86</b>

## Species credits for threatened species

Vegetation zone name	Habitat condition (Vegetation Integrity)	Change in habitat condition	Area (ha)/Count (no. individuals)	BC Act Listing status	EPBC Act listing status	Biodiversity risk weighting	Potential SAI	Species credits
<b><i>Delma impar / Striped Legless Lizard ( Fauna )</i></b>								
1606_Derived_native_grass	31.4	31.4	3.8	Vulnerable	Vulnerable	1.5	False	45
1692_Moderate_good	48.5	48.5	0.5	Vulnerable	Vulnerable	1.5	False	9



								<b>Subtotal</b>	<b>54</b>
<b><i>Petaurus norfolcensis / Squirrel Glider ( Fauna )</i></b>									
1606_Derived_nati ve_grass	31.4	31.4	1.1	Vulnerable	Not Listed	2	False		17
								<b>Subtotal</b>	<b>17</b>

## Proposal Details

Assessment Id 00026727/BAAS17004/21/00026729	Proposal Name Malabar Coal Maxwell Modification 2021 to Water Treatment Plant	BAM data last updated * 10/06/2021
Assessor Name Colin Driscoll	Report Created 05/08/2021	BAM Data version * 45
Assessor Number BAAS17004	BAM Case Status Finalised	Date Finalised 05/08/2021
Assessment Revision 0	Assessment Type Major Projects	

\* Disclaimer: BAM data last updated may indicate either complete or partial update of the BAM calculator database. BAM calculator database may not be completely aligned with Bionet.

## Ecosystem credits for plant communities types (PCT), ecological communities & threatened species habitat

Zone	Vegetation zone name	TEC name	Current Vegetation integrity score	Change in Vegetation integrity (loss / gain)	Area (ha)	BC Act Listing status	EPBC Act listing status	Species sensitivity to gain class (for BRW)	Biodiversity risk weighting	Potential SAI	Ecosystem credits
<b>Bull Oak grassy woodland of the central Hunter Valley</b>											
1	1692_Low	Not a TEC	38.2	38.2	0.1			High Sensitivity to Potential Gain	1.75		2
										<b>Subtotal</b>	<b>2</b>
										<b>Total</b>	<b>2</b>

## Species credits for threatened species

Vegetation zone name	Habitat condition (Vegetation Integrity)	Change in habitat condition	Area (ha)/Count (no. individuals)	BC Act Listing status	EPBC Act listing status	Biodiversity risk weighting	Potential SAI	Species credits
----------------------	--	-----------------------------	-----------------------------------	-----------------------	-------------------------	-----------------------------	---------------	-----------------

ATTACHMENT F  
BAM BIODIVERSITY CREDIT REPORT (LIKE FOR LIKE)



# BAM Biodiversity Credit Report (Like for like)

## Proposal Details

Assessment Id	Proposal Name	BAM data last updated *
00026727/BAAS17004/21/00026728	Malabar Coal Maxwell Mine Entry Area	10/06/2021
Assessor Name	Assessor Number	BAM Data version *
Colin Driscoll	BAAS17004	45
Proponent Names	Report Created	BAM Case Status
	05/08/2021	Finalised
Assessment Revision	Assessment Type	Date Finalised
0	Major Projects	05/08/2021

\* Disclaimer: BAM data last updated may indicate either complete or partial update of the BAM calculator database. BAM calculator database may not be completely aligned with Bionet.

## Potential Serious and Irreversible Impacts

Name of threatened ecological community	Listing status	Name of Plant Community Type/ID
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highla	Critically Endangered Ecological Community	1606-White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter
Species		

## BAM Biodiversity Credit Report (Like for like)

Nil

### Additional Information for Approval

PCTs With Customized Benchmarks

PCT

No Changes

Predicted Threatened Species Not On Site

Name

**Climacteris picumnus victoriae** / Brown Treecreeper (eastern subspecies)

**Grantiella picta** / Painted Honeyeater

**Petaurus australis** / Yellow-bellied Glider

**Phascolarctos cinereus** / Koala

**Pteropus poliocephalus** / Grey-headed Flying-fox

**Anthochaera phrygia** / Regent Honeyeater

**Haliaeetus leucogaster** / White-bellied Sea-Eagle

### Ecosystem Credit Summary (Number and class of biodiversity credits to be retired)



## BAM Biodiversity Credit Report (Like for like)

Name of Plant Community Type/ID	Name of threatened ecological community	Area of impact	HBT Cr	No HBT Cr	Total credits to be retired
1606-White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter	White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highla	3.8	0	75	75
1692-Bull Oak grassy woodland of the central Hunter Valley	Not a TEC	0.5	0	11	11

1606-White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter	Like-for-like credit retirement options					
	Name of offset trading group	Trading group	Zone	HBT	Credits	IBRA region
	White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highla This includes PCT's: 74, 75, 83, 250, 266, 267, 268, 270, 274, 275, 276,	-	1606_Derived_native_grass	No	75	Hunter, Ellerston, Karuah Manning, Kerrabee, Liverpool Range, Peel, Tomalla, Upper Hunter, Wyong and Yengo.  or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.



## BAM Biodiversity Credit Report (Like for like)

	277, 278, 279, 280, 281, 282, 283, 284, 286, 298, 302, 312, 341, 342, 347, 350, 352, 356, 367, 381, 382, 395, 401, 403, 421, 433, 434, 435, 436, 437, 451, 483, 484, 488, 492, 496, 508, 509, 510, 511, 528, 538, 544, 563, 567, 571, 589, 590, 597, 599, 618, 619, 622, 633, 654, 702, 703, 704, 705, 710, 711, 796, 797, 799, 840, 847, 851, 921, 1099, 1103, 1303, 1304, 1307, 1324, 1329, 1330, 1331, 1332, 1333, 1334, 1383, 1401, 1512, 1606, 1608, 1611, 1691, 1693, 1695, 1698					
<b>1692-Bull Oak grassy woodland of the central Hunter Valley</b>	<b>Like-for-like credit retirement options</b>					
	Class	Trading group	Zone	HBT	Credits	IBRA region



## BAM Biodiversity Credit Report (Like for like)

	Coastal Valley Grassy Woodlands This includes PCT's: 116, 618, 622, 623, 760, 761, 762, 829, 830, 834, 837, 838, 849, 850, 1326, 1395, 1603, 1604, 1691, 1692	Coastal Valley Grassy Woodlands >=50% and <70%	1692_Moderate_good	No	11	Hunter, Ellerston, Karuah Manning, Kerrabee, Liverpool Range, Peel, Tomalla, Upper Hunter, Wyong and Yengo.  or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.
--	---	--	--------------------	----	----	---

### Species Credit Summary

Species	Vegetation Zone/s	Area / Count	Credits
<b>Delma impar</b> / Striped Legless Lizard	<b>1606_Derived_native_grass, 1692_Moderate_good</b>	4.3	54.00
<b>Petaurus norfolcensis</b> / Squirrel Glider	<b>1606_Derived_native_grass</b>	1.1	17.00

### Credit Retirement Options

Like-for-like credit retirement options

Species	Spp	IBRA subregion
<b>Delma impar</b> / Striped Legless Lizard	<b>Delma impar</b> / Striped Legless Lizard	Any in NSW



## BAM Biodiversity Credit Report (Like for like)

<b>Petaurus norfolcensis / Squirrel Glider</b>	Spp	IBRA subregion
	<b>Petaurus norfolcensis / Squirrel Glider</b>	Any in NSW



# BAM Biodiversity Credit Report (Like for like)

## Proposal Details

Assessment Id	Proposal Name	BAM data last updated *
00026727/BAAS17004/21/00026729	Malabar Coal Maxwell Modification 2021 to Water Treatment Plant	10/06/2021
Assessor Name	Assessor Number	BAM Data version *
Colin Driscoll	BAAS17004	45
Proponent Names	Report Created	BAM Case Status
	05/08/2021	Finalised
Assessment Revision	Assessment Type	Date Finalised
0	Major Projects	05/08/2021

\* Disclaimer: BAM data last updated may indicate either complete or partial update of the BAM calculator database. BAM calculator database may not be completely aligned with Bionet.

## Potential Serious and Irreversible Impacts

Name of threatened ecological community	Listing status	Name of Plant Community Type/ID
<b>Nil</b>		
Species		
<b>Nil</b>		

## Additional Information for Approval

Assessment Id

00026727/BAAS17004/21/00026729

Proposal Name

Malabar Coal Maxwell Modification 2021 to Water Treatment

Page 1 of 3

## BAM Biodiversity Credit Report (Like for like)

### PCTs With Customized Benchmarks

PCT

No Changes

### Predicted Threatened Species Not On Site

Name

**Haliaeetus leucogaster** / White-bellied Sea-Eagle

### Ecosystem Credit Summary (Number and class of biodiversity credits to be retired)

Name of Plant Community Type/ID	Name of threatened ecological community	Area of impact	HBT Cr	No HBT Cr	Total credits to be retired
1692-Bull Oak grassy woodland of the central Hunter Valley	Not a TEC	0.1	0	2	2

## BAM Biodiversity Credit Report (Like for like)

<b>1692-Bull Oak grassy woodland of the central Hunter Valley</b>	<b>Like-for-like credit retirement options</b>					
	Class	Trading group	Zone	HBT	Credits	IBRA region
	Coastal Valley Grassy Woodlands This includes PCT's: 116, 618, 622, 623, 760, 761, 762, 829, 830, 834, 837, 838, 849, 850, 1326, 1395, 1603, 1604, 1691, 1692	Coastal Valley Grassy Woodlands >=50% and <70%	1692_Low	No	2	Hunter, Ellerston, Karuah Manning, Kerrabee, Liverpool Range, Peel, Tomalla, Upper Hunter, Wyong and Yengo.  or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.

### Species Credit Summary

No Species Credit Data

### Credit Retirement Options

Like-for-like credit retirement options

ATTACHMENT G  
BAM BIODIVERSITY CREDIT REPORT (VARIATIONS)

## Proposal Details

<b>Assessment Id</b>	Proposal Name	BAM data last updated *
00026727/BAAS17004/21/00026728	Malabar Coal Maxwell Mine Entry Area	10/06/2021
Assessor Name	Assessor Number	BAM Data version *
Colin Driscoll	BAAS17004	45
Proponent Name(s)	Report Created	BAM Case Status
	05/08/2021	Finalised
Assessment Revision	Assessment Type	Date Finalised
0	Major Projects	05/08/2021

\* Disclaimer: BAM data last updated may indicate either complete or partial update of the BAM calculator database. BAM calculator database may not be completely aligned with Bionet.

## Potential Serious and Irreversible Impacts

Name of threatened ecological community	Listing status	Name of Plant Community Type/ID
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highla	Critically Endangered Ecological Community	1606-White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter
Species		
<b>Nil</b>		

## Additional Information for Approval

PCTs With Customized Benchmarks

# BAM Biodiversity Credit Report (Variations)

PCT

No Changes

Predicted Threatened Species Not On Site

Name

**Climacteris picumnus victoriae** / Brown Treecreeper (eastern subspecies)

**Grantiella picta** / Painted Honeyeater

**Petaurus australis** / Yellow-bellied Glider

**Phascolarctos cinereus** / Koala

**Pteropus poliocephalus** / Grey-headed Flying-fox

**Anthochaera phrygia** / Regent Honeyeater

**Haliaeetus leucogaster** / White-bellied Sea-Eagle

## Ecosystem Credit Summary (Number and class of biodiversity credits to be retired)

Name of Plant Community Type/ID	Name of threatened ecological community	Area of impact	HBT Cr	No HBT Cr	Total credits to be retired
1606-White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter	White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highla	3.8	0	75	75.00
1692-Bull Oak grassy woodland of the central Hunter Valley	Not a TEC	0.5	0	11	11.00



## BAM Biodiversity Credit Report (Variations)

**1606-White Box - Narrow-leaved Ironbark - Blakely's Red Gum shrubby open forest of the central and upper Hunter**

**Like-for-like credit retirement options**

Class	Trading group	Zone	HBT	Credits	IBRA region
-------	---------------	------	-----	---------	-------------

## BAM Biodiversity Credit Report (Variations)

	<p>White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highla</p> <p>This includes PCT's: 74, 75, 83, 250, 266, 267, 268, 270, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 286, 298, 302, 312, 341, 342, 347, 350, 352, 356, 367, 381, 382, 395, 401, 403, 421, 433, 434, 435, 436, 437, 451, 483, 484, 488, 492, 496, 508, 509, 510, 511, 528, 538, 544, 563, 567, 571, 589, 590, 597, 599, 618, 619, 622, 633, 654, 702, 703, 704, 705, 710, 711, 796, 797, 799, 840, 847, 851, 921, 1099, 1103, 1303, 1304, 1307, 1324, 1329, 1330, 1331, 1332, 1333, 1334, 1383, 1401, 1512, 1606, 1608, 1611, 1691, 1693, 1695, 1698</p>	-	1606_Derived_native_grass	No	75	<p>Hunter, Ellerston, Karuah Manning, Kerrabee, Liverpool Range, Peel, Tomalla, Upper Hunter, Wyong and Yengo.</p> <p>or</p> <p>Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.</p>
--	--	---	---------------------------	----	----	--

## BAM Biodiversity Credit Report (Variations)

<b>1692-Bull Oak grassy woodland of the central Hunter Valley</b>	<b>Like-for-like credit retirement options</b>					
	Class	Trading group	Zone	HBT	Credits	IBRA region
	Coastal Valley Grassy Woodlands This includes PCT's: 116, 618, 622, 623, 760, 761, 762, 829, 830, 834, 837, 838, 849, 850, 1326, 1395, 1603, 1604, 1691, 1692	Coastal Valley Grassy Woodlands >=50% and <70%	1692_Moderate_good	No	11	Hunter, Ellerston, Karuah Manning, Kerrabee, Liverpool Range, Peel, Tomalla, Upper Hunter, Wyong and Yengo. or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.
	<b>Variation options</b>					
	Formation	Trading group	Zone	HBT	Credits	IBRA region
Grassy Woodlands	Tier 3 or higher threat status	1692_Moderate_good	No	11	IBRA Region: Sydney Basin, or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.	

### Species Credit Summary

Species	Vegetation Zone/s	Area / Count	Credits
<b>Delma impar</b> / Striped Legless Lizard	<b>1606_Derived_native_grass, 1692_Moderate_good</b>	4.3	54.00
<b>Petaurus norfolcensis</b> / Squirrel Glider	<b>1606_Derived_native_grass</b>	1.1	17.00

### Credit Retirement Options    Like-for-like options

## BAM Biodiversity Credit Report (Variations)

<b>Delma impar/</b> Striped Legless Lizard	Spp		IBRA region
	<b>Delma impar/</b> Striped Legless Lizard		Any in NSW
	<b>Variation options</b>		
	Kingdom	Any species with same or higher category of listing under Part 4 of the BC Act shown below	IBRA region
Fauna	Vulnerable	Hunter, Ellerston, Karuah Manning, Kerrabee, Liverpool Range, Peel, Tomalla, Upper Hunter, Wyong and Yengo.  or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.	
<b>Petaurus norfolcensis/</b> Squirrel Glider	Spp		IBRA region
	<b>Petaurus norfolcensis/</b> Squirrel Glider		Any in NSW
	<b>Variation options</b>		
	Kingdom	Any species with same or higher category of listing under Part 4 of the BC Act shown below	IBRA region

## BAM Biodiversity Credit Report (Variations)

	Fauna	Vulnerable	<p>Hunter, Ellerston, Karuah Manning, Kerrabee, Liverpool Range, Peel, Tomalla, Upper Hunter, Wyong and Yengo.</p> <p>or</p> <p>Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.</p>
--	-------	------------	--

## Proposal Details

<b>Assessment Id</b>	Proposal Name	BAM data last updated *
00026727/BAAS17004/21/00026729	Malabar Coal Maxwell Modification 2021 to Water Treatment Plant	10/06/2021
Assessor Name	Assessor Number	BAM Data version *
Colin Driscoll	BAAS17004	45
Proponent Name(s)	Report Created	BAM Case Status
	05/08/2021	Finalised
Assessment Revision	Assessment Type	Date Finalised
0	Major Projects	05/08/2021

\* Disclaimer: BAM data last updated may indicate either complete or partial update of the BAM calculator database. BAM calculator database may not be completely aligned with Bionet.

## Potential Serious and Irreversible Impacts

Name of threatened ecological community	Listing status	Name of Plant Community Type/ID
<b>Nil</b>		
Species		
<b>Nil</b>		

## Additional Information for Approval

PCTs With Customized Benchmarks

PCT
No Changes

## Predicted Threatened Species Not On Site

Name
<b>Haliaeetus leucogaster</b> / White-bellied Sea-Eagle

## Ecosystem Credit Summary (Number and class of biodiversity credits to be retired)

Name of Plant Community Type/ID	Name of threatened ecological community	Area of impact	HBT Cr	No HBT Cr	Total credits to be retired
1692-Bull Oak grassy woodland of the central Hunter Valley	Not a TEC	0.1	0	2	2.00

1692-Bull Oak grassy woodland of the central Hunter Valley	Like-for-like credit retirement options					
	Class	Trading group	Zone	HBT	Credits	IBRA region
	Coastal Valley Grassy Woodlands This includes PCT's: 116, 618, 622, 623, 760, 761, 762, 829, 830, 834, 837, 838, 849, 850, 1326, 1395, 1603, 1604, 1691, 1692	Coastal Valley Grassy Woodlands >=50% and <70%	1692_Low	No	2	Hunter, Ellerton, Karuah Manning, Kerrabee, Liverpool Range, Peel, Tomalla, Upper Hunter, Wyong and Yengo. or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.
Variation options						
Formation	Trading group	Zone	HBT	Credits	IBRA region	
Grassy Woodlands	Tier 3 or higher threat status	1692_Low	No	2	IBRA Region: Sydney Basin, or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.	

## Species Credit Summary



# BAM Biodiversity Credit Report (Variations)

---

No Species Credit Data

**Credit Retirement Options**    Like-for-like options